

Long-term land use trends and their legacies in the Carpathian Region of Eastern Europe

by

Catalina Munteanu

A dissertation completed in partial fulfillment of

the requirements for the degree of

Doctor of Philosophy

(Forestry)

at the

UNIVERSITY OF WISCONSIN – MADISON

2016

Date of final oral examination: 05/02/2016

This dissertation is approved by the following committee:

Volker C. Radeloff, Professor, Forest and Wildlife Ecology

Murray Clayton, Professor, Forest and Wildlife Ecology

Holly Gibbs, Assistant Professor, Department of Geography, Nelson Institute of Environmental Sciences

Tobias Kuemmerle, Professor, Department of Geography, Berlin, Germany

Anna M. Pidgeon, Professor, Forest and Wildlife Ecology

Acknowledgements

I would like to express my deepest appreciation for all those who made my PhD endeavor possible. I am grateful to my advisor, Volker Radeloff for his mentorship, for teaching me to develop an inquisitive mind and for his endless excitement for our scholarship. I am thankful to my committee members: Anna Pidgeon for her subtle yet strong support throughout all aspects of grad school, and for sparking my interest in birds; Tobias Kuemmerle for his razor-sharp mind and for exceptional generosity with his time; Holly Gibbs for being a model of success and for boosting my confidence in the first year of my PhD and Murray Clayton for sharing his excitement and love of statistics.

I am indebted to all project partners, collaborators and co-authors for collecting and sharing data, and for their knowledge on the Carpathians. Daniel Müller, thanks for always constructively challenging my thinking. Nick Keuler, thanks for your patience and perseverance in making me understand statistics. Dominik Kaim, Katarzyna Ostapowicz and Juraj Lieskovsky, thanks for topping your hard work in this project with friendship and kindness. Mihai Nita, thanks for sharing your enthusiasm for historical forestry and for being the oh-so-Romanian part of my PhD. Thanks to É. Konkoly-Gyuró, L. Halada, J. Kozak, G. Kiraly, P. Griffiths, U. Gimmi, P. Balasz, A. Shandra, P. Stych, M. Boltiziar, M. Mojses, K. Ostafin, M. Dobosz, E. Grabska, A. Prociak, M. Chmiel, P. Stuglik and K. Kysucká for all your contributions to this endeavor. Our work was possible due to support from the NASA's Land Use Land Cover Change Program and Earth System Science Fellowship.

Life in Madison was a wonderful experience. I feel very lucky to have been part of the SILVIS lab, which was to me a source of friendship, inspiration and feedback over the last four years: thank you Shelley for sharing laughs, tears and office space, and for cheerfully enduring my at

times excessive romanian-ness. Johanna and Ana-Maria, thanks for often reminding me of the little fun ways in which to enjoy being a grad student. Martina thanks for your academic insights and for the marshmallow victories in Brandi-dog. He, thank you for becoming a caring and trusted friend in the shortest time. I also am most grateful to my friends, who kept me sane and happy in my non-academic life: Jamila Sidiqqi, thanks for many delicious cups of tea, spiced with best conversation; Andria Ellis, thanks for craft nights, numerous treats and for always lending an ear; Raluca, thanks for adding color and long bike rides into the mix. Patricia von Maravic and Claudia Drexler – thanks for sending me many wonderful letters over the past years, to which I often went back when feeling overwhelmed.

I am forever grateful to Betty VanLeuven, David Reichenberger and Simon Goldberg for becoming my family here in Madison. Betty, thanks for being my model for active and positive thinking, David, thanks for being my model of kindness and deep contemplation, and Simon, thanks for filling the space of a brother I never had. Back home across the Ocean, I am blessed with the most wonderful family ever. Gigi and Lili Ciobanu, the words ‘unchiu’ and ‘tusi’ are too small to fit all the love, care and support you have shown me over time – thank you. Andrei and Horia, thanks for all of our small and big adventures together, including moss-paintings on T.Ala’s fence. I can hardly express how grateful I feel for my parents Sorin and Mariana Munteanu, who have been by my side in all aspects of life. I am lucky to have you both: Dad thanks for teaching me attention to detail and perseverance, Mom thanks for showing me love and navigating with me through the thickness of life events.

Ben Kraemer, thank you for continual confidence, support and encouragement, from portaging a 75lb canoe to finishing up a 5 chapter dissertation. Thanks for paper submission dances, comforting hugs and endless delicious breakfasts. Thanks for sharing life with me.

Table of Contents

Introduction.....	1
<i>Chapter summaries</i>	<i>7</i>
<i>Related publications</i>	<i>14</i>
<i>Significance and implications</i>	<i>16</i>
<i>References</i>	<i>21</i>
 Chapter 1. Forest and agricultural land change in the Carpathian region – a meta-analysis of long-term patterns and drivers of change	 25
<i>Introduction</i>	<i>25</i>
<i>Methods</i>	<i>27</i>
<i>Results</i>	<i>33</i>
<i>Discussion</i>	<i>36</i>
<i>References</i>	<i>44</i>
<i>Tables and figures</i>	<i>53</i>
<i>Appendix.....</i>	<i>61</i>
 Chapter 2. Legacies of 19th century land use shape contemporary forest cover	 64
<i>Introduction</i>	<i>64</i>
<i>Methods</i>	<i>67</i>
<i>Results</i>	<i>74</i>
<i>Discussion</i>	<i>78</i>

<i>Conclusions</i>	84
<i>References</i>	86
<i>Tables and figures</i>	94
<i>Appendix</i>	101
Chapter 3. Historical forest management in Romania is imposing strong legacies on contemporary forests and their management	106
<i>Introduction</i>	106
<i>Methods</i>	108
<i>Results</i>	116
<i>Discussion</i>	120
<i>Conclusions</i>	126
<i>References</i>	128
<i>Figures and tables</i>	136
Chapter 4. Land use legacies of historic political regimes for contemporary land abandonment	147
<i>Introduction</i>	147
<i>Methods</i>	151
<i>Results</i>	158
<i>Discussion</i>	163
<i>References</i>	171

<i>Tables and figures</i>	177
<i>Appendix</i>	188
Chapter 5. Conservation recommendations for the protection of bird habitat in the Carpathian Ecoregion in light long-term land use trends	193
<i>Introduction</i>	193
<i>Methods</i>	196
<i>Results</i>	201
<i>Discussion</i>	204
<i>Conclusions</i>	209
<i>References</i>	210
<i>Tables and figures</i>	216
<i>Appendix</i>	224

List of Figures

- Figure 1.1 a) Study area, including spatial extent of case studies (grey) and centroids for 102 case study locations (triangles). Country codes: AT: Austria, HU: Hungary, PL: Poland, CZ: Czech Republic, SK: Slovakia, UA: Ukraine, RO: Romania, MD: Moldova, HR: Serbia, SI: Slovenia. b) Spatio-temporal distribution of 102 study locations (one location may include multiple case-studies). 55
- Figure 1.2 Representativeness check of case studies biophysical characteristics for the country's share of the study areas: comparison of a) elevation and b) slope. The grey line indicates the mean value for the country's share of the study area. c) Soil type distribution for the case studies in each country and for the region as a whole. Country codes: CZ: Czech Republic, HU: Hungary, PL: Poland, RO: Romania, SK: Slovakia, UA: Ukraine..... 56
- Figure 1.3 Proportion of studies reporting decreasing ($<-0.1\%$ annually), stable (-0.1% to 0.1% annually) and increasing ($>0.1\%$ annually) cover for each time period for a) forest and c) agriculture and distribution of annual rates of change per time period for b) forest and d) agricultural cover. Country codes: CZ: Czech Republic, HU: Hungary, PL: Poland, RO: Romania, SK: Slovakia, UA: Ukraine..... 57
- Figure 1.4 Spatial and temporal distribution of forest change case studies. Annual rates of change are mapped for each case study and time period. Studies are represented by centroids. The size of the symbols indicate the amount of change, the colors indicate the direction of change (increase/stability/decrease). Shaded colors indicate that annual rates are calculated for more than one time period. Country codes: CZ: Czech Republic, HU: Hungary, PL: Poland, RO: Romania, SK: Slovakia, UA: Ukraine. ... 58

Figure 1.5 Spatial and temporal distribution of agricultural change case studies. Annual rates of change are mapped for each case study and time period. Studies are represented by centroids. The size of the symbols indicate the amount of change, the colors indicate the direction of change (increase/stability/decrease). Shaded colors indicate that annual rates are calculated for more than one time period. Country codes: CZ: Czech Republic, HU: Hungary, PL: Poland, RO: Romania, SK: Slovakia, UA: Ukraine. ... 59

Figure 1.6 Main classes of land change drivers and the relative importance of drives for each land change process in the study area. The proportions are calculated based on the number of times a driver was deemed as important in influencing change. 60

Figure 2.1 Study area in Eastern Europe and forest cover maps for the 1860s, 1960s and 2010s. CZ: Czech Republic, HU: Hungary PL: Poland, RO: Romania, SK: Slovakia, UA: Ukraine..... 96

Figure 2.2 Percentage forest disturbance in the Carpathian Region (overall) and by country (RO: Romania, SK: Slovakia, UA: Ukraine, PL: Poland, CZ: Czech Republic, HU: Hungary) in old forests and new forests. Old forests refer to areas that were forested throughout 1860s-1985, new forests to areas not forested in 1860s, but forested in 1960s. The graph indicates consistently more forest disturbance in new forests. Number of observations per country in brackets. 97

Figure 2.3 Odds of forest disturbance (in %) in new forests (not forested in 1860s) compared to old forests (forested in 1860s), overall and in country models (RO: Romania, SK: Slovakia, UA: Ukraine, PL: Poland, CZ: Czech Republic, HU: Hungary) 98

Figure 2.4 Proportion of forest disturbance (left y axis) in old forests (black line) and new forests (grey line) in relation to elevation, slope, annual mean temperature, annual

precipitation, crop suitability, length of growing season, accessibility, distance to cities, settlements, roads and rivers. The graph represents a data summary of all observed disturbance. The dotted line represents number of forest observations in 1985 (right y axis). Continuous variables were divided in 30 equal interval data bins and the proportion of disturbance as well as the mean number of forest observations was plotted for each bin. For model based results please see Appendix 2.4	99
Figure 2.5 (a) Forest types by year and country since 1860s, (b) 1860s forest types of disturbed forests between 1985-2010, (c)1860s forest types of old forests (d) 2010s forest types in new forests, and (e) 2010s forest types of old forests . RO: Romania, SK: Slovakia, UA: Ukraine, CZ: Czech Republic, HU: Hungary. Poland has been excluded from this graph due to scarcity of forest type data. Note: Figures are not for entire country territory but for the study region depicted in Figure 2.1. Forest type distribution in (a) is not identical to Griffiths et al (2013) because the data presented here is restricted to 1960 forest cover.....	100
Figure 3.1 Location of study area in Europe, imperial boundaries from 1900s, the location of 3 cases studies in the Carpathian Mountains and county borders for 1924 and 2015 .	140
Figure 3.2 Overview of the evolution of forest cover, species composition, disturbance and ownership patterns in Romania between 1870s and 2010, in the context of major land tenure changes.....	141
Figure 3.3. Percentage forest cover and forest disturbance in Romania historically (n= 58 counties) and contemporary (n=42 counties).....	142
Figure 3.4 Historical and contemporary forest disturbance at forest stand level in three case studies: Humor (3500ha), Oituz (9000ha) and Madaras (1500ha)	143

Figure 3.5 Percentage of major coniferous species (<i>Picea abies</i> , <i>Abies alba</i>) and major deciduous species (<i>Fagus sylvatica</i> , <i>Quercus sp</i>) within forest cover of Romanian regions in 1924 (n= 58 regions) and in 2014 (n=42 regions)	144
Figure 3.6 Species composition at local level in Humor (3500ha), Oituz (9000ha) and Madaras (1500ha) at the beginning of the 20th century and the beginning of the 21st century. Stands with species cover higher than 50% are represented in the graphic.....	145
Figure 3.7 Forest age distribution in Romania in at (a) national level for the years 1924, 1964 and 2014 and at local case study level in (b) Oituz (9000ha) between 1930 and 2013, (c) Humor (3500ha) between 1926 and 2006 and (d) Madaras (1500ha) between 1945 and 2006.....	146
Figure 4.1 Study objectives: persistence and strength of land use legacies. For Obj. 1 we developed two models (A,B), both of which compared abandonment of 1930s agricultural land that was already farmed in 1860 with land that was not farmed then, and we modelled the effect of early versus late Habsburg agriculture on both A) socialist and B) post-socialist land abandonment.. For Obj. 2, we modelled the legacy effect of three periods, i.e., the C) Habsburg, D) Interwar and E) Socialist period, on post-socialist abandonment. Models C is identical to model B for objective 2. Model D compares land farmed at any point during the Habsburg era with land farmed at any time during the Socialist period. Model E refers to socialist agricultural expansion and compares abandonment on land farmed prior to 1960, i.e., during the early Socialist period, with land farmed after 1960 (i.e. late Socialist). Please see Table 1 for definitions of terms and time periods	180

Figure 4.2 Study area. Country codes: CZ: Czech Republic, HU: Hungary PL: Poland, RO: Romania, SK: Slovakia, UA: Ukraine	182
Figure 4.3 Agricultural dynamics in the Carpathian region and by country.	183
Figure 4.4 Agricultural cover between 1860 and 2010, socialist and post-socialist land abandonment in the Carpathian region (see Table 4.1 for definitions).....	184
Figure 4.5 Model performance for distribution of agricultural land as a function of environmental variables during the Habsburg, socialist and post-socialist eras. Model performance is reported as area under the receiver operator curve (AUC), where values close to 1 indicate high model performance.....	185
Figure 4.6 Persistence of Habsburg land use legacies. Odds of abandonment (in %) in areas that were not farmed by the Habsburgs in 1860s, compared to areas farmed then. The odds of abandonment on land not farmed in 1860s were higher and the relationship was consistent across countries. For example, in Romania, the odds of socialist abandonment (a) were 63% higher in areas farmed in 1860s, compared to areas not farmed then and the odds of post-socialist abandonment (b) were 86% higher.	186
Figure 4.7 Strength of land use legacies of political regimes. Odds of abandonment (in %) in areas that were not farmed in a given historic period, compared to areas farmed then. The odds of abandonment on land not farmed historically were higher and the relationship was consistent across countries. For example, in Romania, the odds of abandonment were 86% higher in areas farmed in 1860s, compared to areas not farmed then. Legacy effects were strongest for the political transformation legacy, which captures differences in political regimes between the Habsburg and Socialist periods.....	187

Figure 5.1 Study region: Carpathain Ecoregion in Eastern Europe.....	219
Figure 5.2 (a) Evolution of potential major habitat (grassland, agriculture and forest) for a species whose range covers the entire Carpathians. (b) Evolution of agriculture, grassland and forest for all species that use these habitats. Double counting is possible if a species uses more than one habitat. This does not take into account dynamics of other habitat types.	220
Figure 5.3 Percentage of European range in the Carpathians for 170 species, by major habitat type.....	221
Figure 5.4 Potential habitat dynamics for species of high conservation responsibility. The color of the line represents the conservation responsibility class. Red: HHCD Blue: HHLC Green: HMCD.....	222
Figure 5.5 IUCN threats and number of times they occur listed for the 29 species of conservation responsibility.....	223

List of Tables

Table 1.1 Time periods, their duration, the expected land changes and the number of studies that report land change for the specific period. The first number (*) indicates that the annual rate of change has been calculated for only one period. The second number (**) indicates that the case-study spans at least two time periods, and the annual rate of change is calculated based only on land cover at the beginning and end of the considered time span.	53
Table 1.2 Comparison of expected and observed land changes for each time period and the mean annual rates of change, calculated for all the case studies for which change rates were not spanning more than one period (marked * in Table 1.1). For these calculations, only studies that report annual change for single periods were considered.	54
Table 2.1 Maps and satellite images used for forest cover mapping.....	94
Table 2.2 List of predictors used in the forest disturbance models (n=19947), including data sources, measurements in units (Unit), their spatial resolution (SpRes), mean values (Mean), standard deviation (SD), range (Min, Max)	95
Table 3.1 The three case studies, including name, area, historic region, type of management and forest ownership	136
Table 3.2 Data sources for forest extent, composition, age classes and disturbances for three time periods and at two spatial scales.....	137
Table 3.3 Data sets used in our analysis and comparison to values from other sources such as forestry literature, statistical yearbooks and remote sensing estimates for historical (a) and contemporary (b) time periods	138
Table 4.1 Paper terminology.....	177

Table 4.2 Map and remote sensing data sources used to map agricultural land cover in the Carpathians	178
Table 4.3 List of predictor used in logistic regression models, including land use legacies, agro-ecological conditions, accessibility and socio-political variables.	179
Table 5.1 Conservation responsibility classes for 252 bird species in the Carpathians	216
Table 5.2 Major habitat types for species of high conservation responsibility. Highlighted species use more than one habitat type.	217
Table 5.3 IUCN threats related to agricultural and forest systems for species of high conservation responsibility	218

List of Appendices

Appendix 1.1 List of publications included in the meta-analysis	61
Appendix 1.2 List of publications included in the meta-analysis	62
Appendix 2.1 Number of observations for the entire dataset and for each country for each old forests (F) and new forests (NF). Not disturbed = forested in 1960 and never disturbed between 1985-2010, Disturbed = forested in 1960 and disturbed at least once in a five year time step between 1985-2010.....	101
Appendix 2.2 Model outputs for best selected models: regression coefficients, odds ratio (OR) of disturbance on historic non forest vs forest, and model performance as area under ROC curve (AUC). An odds ratio>1 indicates higher rate of disturbance on 1860 non-forest areas.	102
Appendix 2.3 Number of best performing models ($\Delta AIC < 3$), of those number of models that include land use legacy as an explanatory variable, ΔAIC of best model including legacies. The second part of the table shows the odds ratios for each selected variable in the exhaustive search	103
Appendix 2.4 Disturbance in old forests (black line) and new forests (grey line) in relation to elevation, slope, annual mean temperature, annual precipitation, crop suitability, length of growing season, accessibility, distance to cities, settlements, roads and rivers. The graph represents regression results once we controlled for all other spatial determinants of change. For a data-based summary please see Figure 2.4	104
Appendix 2.5. Numbers of observations for forest type data by year and country. Forest type (FT) column indicates the number of points for which forest type information was	

available. No forest type (NoFT) column indicates the number of data points that were identified as forest but for which forest type information could not be retrieved. (RO: Romania, SK: Slovakia, UA: Ukraine, PL: Poland, CZ: Czech Republic, HU: Hungary).....	105
Appendix 4.1 Sample sizes for legacy persistence modelling.....	188
Appendix 4.2 Sample sizes for strength of legacies in relation to political regimes	189
Appendix 4.3 Model utility and performance (AUC) for three legacy models for the Carpathian region as a whole and by country	191
Appendix 4.4 Variables selected in socialist abandonment models and sign of the coefficient indicating the direction of the effect	191
Appendix 4.5 Variables selected in post-socialist abandonment models and sign of the coefficient indicating the direction of the effect.	192
Appendix 5.1 Species of high conservation responsibility for the Carpathians and their species attributes.....	224
Appendix 5.2 Possible Carpathian landscape configuration. Left: habitat distribution in 2010. Right: potential future habitat distribution if all agricultural land would be abandoned, and forest succession would occur in all areas that were forested at any point after 1860s.....	227

Introduction

“Brothers, what we do in life, echoes in eternity!”

- Russell Crowe, *Gladiator* (2000)

We all have a choice between leaving a positive or negative legacy for generations to come. As individuals, we often make this decision consciously, for example, by securing the future of our children. But as a society, we rarely consider the responsibility that we carry for generations ahead. This may be partly because we do not fully understand how our ancestor’s legacies affect our lives and our environment today. My dissertation is looking at a special type of legacies – land use legacies from a century ago – and how they affect contemporary land change and conservation. I ask if and how land use legacies affect current land change, and what is our environmental responsibility for future generations? To answer these questions, I turn to land system science and the tools of scientific inquiry that this emerging discipline provides.

The mosaic of land covers and land uses, both drives and responds to global environmental change (Foley et al., 2005; Turner et al., 2013; Verburg et al., 2013). Land use change is one of the largest contributors to changes in climate, biodiversity, ecosystem services and economies, which in turn cause further land change (Gibbs and Herold, 2007; Ojima et al., 1994; Verburg et al., 2013). Land management and conservation efforts are often focused on changing the magnitude or even reversing the direction of land changes that negatively affect the environment (Ferretti-Gallon and Busch, 2014). Usually this requires a shift in the underlying driving forces (such as economic incentives, policies or nature protection) across a range of spatio-temporal scales (Verburg et al., 2013). Most drivers of recent land transitions are well understood (Bürgi

et al., 2005; Geist and Lambin, 2002; Lambin et al., 2001) and could be adjusted to accommodate our future land plans. However, one aspect of land change that remains poorly understood and largely unquantified is the role of historic land uses in driving land change. This is unfortunate, because disregarding legacies may lead to misinterpretation of land change patterns and to ill-informed land use decisions.

Over long time periods, land-use transition theories predict gradual changes among land covers (such as from cropland to forest), primarily as a function of demographic and economic factors (DeFries et al., 2004; Foley et al., 2005; Lambin and Meyfroidt, 2010). A leading theoretical example is the forest transition theory, which postulates that gradual economic and demographic change lead initially to deforestation, but then to agricultural specialization and reforestation of marginal lands. The forest transition itself is the shift from decreasing forest area in a given country or region to increasing forest area (Barbier et al., 2010; Mather, 1992). A regions' socio-economic context and socio-ecological feedbacks may lead to differences in timing and extent of the forest transition (Foley et al., 2005; Lambin and Meyfroidt, 2010; Meyfroidt and Lambin, 2011). Regions where socio-economic and political changes occur at relatively short intervals are particularly interesting in this respect, because they allow comparisons between different socio-economic, political and demographic components of the land change process. Eastern Europe represents a great study case for transition theories, because “the region reinvented itself every half century” in terms of its economies and politics (Good, 1994).

Land use transitions are explained by a series of causal mechanism, i.e., a set of complex factors involved in causing land change (Lambin and Meyfroidt, 2010; Meyfroidt, 2015). The immediate human activities that affect the environment act at local scales and are the proximate causes of land change; they are in turn driven by underlying forces, or complex social, political,

economic, technological, and cultural factors (Geist and Lambin, 2001; Meyfroidt, 2015). For example, the establishment of a new type of land use is constrained by local and national markets or policies, with magnitudes of change being attenuated or amplified by global forces (Lambin et al., 2001). Deforestation in Vietnam was directly driven by coffee expansion which was in turn a result of increasing global coffee demand (Meyfroidt, 2015). The agricultural expansion in Kazakhstan in the 1950s was driven by the increasing demand for agricultural products in the Soviet Union, which in turn resulted from poor policies (Kraemer et al., 2015). One potential shortcoming of current land use theory is that, despite acknowledging path dependency, and the fact that ecosystems may respond to past changes for decades in the future (Foster et al., 2003; Wallin et al., 1994), land use histories and the legacies they create are rarely quantified in land change models (Perring et al., 2016). In Eastern Europe, the factors causing land change after the collapse of the Soviet Union are well understood (Baumann et al., 2011; Kuemmerle et al., 2007; Müller et al., 2013), but historical land use dynamics remain widely unexplored, despite potential lingering effects of historical events (Good, 1994).

Land change has been an active process for millennia, with forest and agricultural land use experiencing substantial changes in the past century at unprecedented rates (Gibbs et al., 2010; Hansen et al., 2010): more than 3% forest was lost from 2000 to 2005 across the globe (Hansen et al., 2010) while agriculture expanded by 3% from 1985 to 2005 (Foley et al., 2011).

Highlighting that land use changes are complex, though, agricultural land of the size of France was also abandoned globally between 1995 and 2005, mostly in the former Soviet Union and Latin America (Munroe et al., 2013). In Europe alone, an area the size of Spain was abandoned since 2001, most of it in the Former Eastern Bloc (Estel et al., 2015). In parts of the World where long term land use transitions are punctuated by political and socio-economic shifts, the changes

in land use can be abrupt and transformations between land uses can succeed at faster pace under economic or institutional pressures. The history of Eastern Europe since the mid 19th century represent in this respect a great ‘natural experiment’ for investigating the effects of different socio-economic and political contexts on long term land change processes.

Despite acknowledging the magnitude of land changes under political transformations, longitudinal studies addressing questions of the legacies they produce are scarce. Legacies and path dependencies have mostly focused on urban systems (Lambin and Geist, 2006; Seto et al., 2012) although scattered evidence suggests legacies exist in forest and agricultural systems (Börjeson, 2007; Coomes et al., 2011; Radeloff et al., 2001). More evidence on legacies is provided in ecology, where legacies of past land uses are shown to linger in the ecosystems for decades or even centuries, affecting carbon storage, soil structure and vegetation composition (Brudvig et al., 2013; Ficetola et al., 2010; Foster et al., 2003). Long term changes in land use and habitat, can have cascading effects for species abundance (Pidgeon et al., 2011), and may cause extinctions or spread of invasive species (Dullinger et al., 2013; Essl et al., 2011; Tilman et al., 1994). This is why land management and conservation planning need to take into account historic and recent land use changes, if they are to provide sustainable solutions for the future environment.

Long term land use trends are important not just for improving our understanding of contemporary land change, but also for devising appropriate conservation measures. Globally, two strategies are considered when aiming to balance food production and nature conservation. Areas of low intensity farming are considered ideal for implementing a land sharing strategy where both agricultural production and nature conservation coexist (Fischer et al., 2012; Mikulcak et al., 2013). Alternatively, under a land sparing scenario, high yield farming in some

places, could free up land for wild nature in others (Fischer et al., 2014; Grau et al., 2013; Phalan et al., 2011). The Carpathian Mountains of Eastern Europe, have been devised as an ideal place for retaining wildlife friendly farming due to the long history of human-environment interactions and high levels of biodiversity in low-intensity landscapes (Fischer et al., 2012; Hartel et al., 2013). On the other hand, the region has also been a primary target for rewilding due to the high rate of agricultural abandonment, contiguous forest ecosystems, and unique presence of large carnivore and herbivore species (Ceașu et al., 2015; Kuemmerle et al., 2010; Navarro and Pereira, 2012). Despite the existence of empirical evidence about which species could benefit under each of these conservation strategies, an integrated view on the conservation responsibility of the Carpathians in light of the historical land use changes is still missing. This is unfortunate, because conservation efforts could be miss-directed at species or habitats whose conservation could bring only minor benefits at European or global level.

The overarching goal of my dissertation was to understand the role of land use legacies for contemporary land use dynamics and for conservation action. I investigated forest and agricultural land use transitions over the past two centuries in Eastern Europe, in the context of the rise and the collapse of several political systems and explored their implications for conservation. My specific objectives were to:

- 1) Identify and quantify long-term forest and agricultural change patterns and their main driving forces in times of socio-economic and political change.
- 2) Assess if and how much land-use legacies affect contemporary forest disturbance and the abundance of forest types
- 3) Explore potential cause-effect relationships between historic forest characteristics and contemporary forest patterns

- 4) Assess the role of land use legacies of different historic time periods on contemporary agricultural land use abandonment
- 5) Provide conservation recommendations, based on historic and recent bird habitat evolution for species of highest conservation responsibility at European level

To address these goals, I studied land use dynamics since the 18th century in the Carpathian region using a combination of historical land use maps, remote sensing data and statistical records, as well as species range maps for birds whose range at least partly cover my study region. Eastern Europe represents a great ‘natural experiment’ for the study of century long term land use dynamics because it experienced multiple political and land management systems which caused drastic land cover changes over a relatively short time span. The substantial changes that occurred in forest and agricultural management over the past centuries in the Carpathians allow me to analyze legacy effects of multiple historic periods at broad spatial scales. Another advantage of the regions is that under the influence of major political regimes, land use was thoroughly documented in form of historical cartographic and statistical material, allowing me to spatially reconstruct over a century of land use history. Although well documented, the extensive land cover changes of the last decades remain poorly understood in the context of historical changes in policies and land use patterns. Furthermore, despite being considered a conservation hotspot in Europe (Hartel et al., 2013; Knorn et al., 2012), it remains unknown for which species the Carpathians carry highest conservation responsibility at European level. The five chapters of my dissertation examine specific questions related to long term land use dynamics in the Carpathians.

Chapter summaries

Chapter 1. Forest and agricultural land change in the Carpathian region – a meta-analysis of long-term patterns and drivers of change

Despite the availability of many small scale land change studies in the Carpathians, the overall land use transitions and their drivers remain broadly unknown. Furthermore, differences induced by shifts in socio-economic and institutional transformations remain largely unexplored at trans-Carpathian scale.

In Chapter 1, I examined land use transition trajectories (such as the forest transition) in the Carpathians over the past 250 years and the relation to underlying drivers of land change. I conducted a meta-analysis of 66 publications describing 102 case study locations and quantified the main forest and agricultural changes in the Carpathian region since the 18th century. I assessed the heterogeneity of the local-scale studies across the region and identified the importance of the different drivers of land-use change for individual forest and agricultural dynamic processes. The case studies captured gradual changes since the peak of the Austro-Hungarian Empire up to the accession to the European Union of most of the formerly socialist countries in the study region. Agricultural land-use increased during the Austro-Hungarian Empire in 70% of the case studies, but dropped sharply during and especially after the collapse of the Socialism (over 70% of the cases). The Carpathian region experienced forest transition during the Interwar period (93% of the cases), and the forest expansion trend persisted after the collapse of Socialism (70% of the cases). In terms of the drivers, institutional and economic factors were most influential in shaping deforestation and agricultural expansion, while socio-demographics and institutional shifts were the key drivers of land abandonment. My study highlights the drastic effects that socio-economic and institutional changes can have on land-use

and land-cover change, and the value of longitudinal studies of land change to uncover these effects.

Related manuscript: Munteanu C, Kuemmerle T, Boltiziar M, Butsic V, Gimmi U, Halada L, Kaim D, Király G, KonkolyGyuró E, Kozak J, Lieskovsky J, Mojses M, Müller D, Ostafin K, Ostapowicz K, Shandra O, Štych P, Walker S, Radeloff VC (2014) Forest and agricultural land change in the Carpathian region—A metaanalysis of long-term patterns and drivers of change. *Land Use Policy* 38:685–697

Chapter 2. Legacies of 19th century land use shape contemporary forest cover.

Although long-term persistence of legacies for ecosystem structure and composition is relatively well understood in ecology, it remains unclear if land use legacies can shape the extent and pattern of current environmental change, affecting management and conservation decisions.

In Chapter 2, I assessed if and how much land-use legacies affect contemporary forest disturbance. Specifically I quantified (1) the magnitude of contemporary forest disturbance in relation to historic land uses, (2) the relation to spatial determinants of forest disturbance, and (3) changes in main forest types. I modeled contemporary forest disturbance (based on satellite image analysis from 1985 to 2010) as a function of historic land use (based on digitized topographic maps from 1860 and 1960). Contemporary forest disturbance was strongly related to historic land use even when controlling for environmental, accessibility and socio-political variation. Across the Carpathian region, the odds of forest disturbance were about 50% higher in areas that were not forested in 1860 (new forests) compared to areas that were forested then (old forests). These legacies may be explained by extensive plantations outside forest ranges, predominantly spruce, poplar, and black locust, which are prone to natural disturbances. Furthermore, as plantations reach harvestable age of about 70 years for pulp and 120 year for

saw-timber, these are likely to be clear-cut, producing the observed legacy effects. Across the Carpathians, forest types shifted towards less coniferous cover in 2010 compared to the 1860s and 1960s likely due to extensive historic conifer harvest, and to recent natural disturbance events and clear-cuts of forest plantations. My results underscore the importance of land-use legacies, and show that past land uses can greatly affect subsequent forest disturbance for centuries. Given rapid land use changes worldwide, it is important to understand how past legacies affect current management and what the impact of current land management decisions may be for future land use.

Related manuscript: Munteanu C, Kuemmerle T, Keuler NS, Müller D, Balázs P, Dobosz M, Griffiths P, Halada L, Kaim D, Király G, Konkoly-Gyuró E, Kozak J, Lieskovsky J, Ostafin K, Ostapowicz K, Sandra O, Radeloff VC (2015) Legacies of 19th century land use shape contemporary forest cover. *Global Environmental Change*, 34: 83-94

Chapter 3. Historical forest management in Romania is imposing strong legacies on contemporary forests and their management

Historical forest management can heavily affect contemporary forest management and conservation. Yet, relatively little is known about century-long changes in forest composition, structure, ownership and management, and that limits the understanding of how past management and land tenure relate to current forestry practice and forest conservation.

In Chapter 3, I examined the relationship between historical forest management (as depicted by historical forest cover, species composition, age structure and harvesting data) and contemporary forest patterns. I used the case study of Romania, the country with the largest share of the Carpathian region, because detailed data on historic forest management was available at multiple spatial scales, allowing me to explore the relationship between past and current management. I

reviewed forestry literature and statistics since the 19th century and reconstructed a time-series of forest cover, composition, disturbance patterns, and ownership patterns, and interpreted these data in light of institutional changes. I also assessed changes in forest cover, forest harvest, species composition and age structure between two points in time (1920s and 2010s) at the county level and for three case studies for which stand- forest management data was available. I found that forest area increased in Romania since 1924 by 5% and that the annual rate of forest harvest between 2000 and 2013 was half of the annual rate between 1912 and 1922, which indicates high potential for forest biodiversity conservation. However, the composition, distribution, and age structure of contemporary forests is also substantially different from historical forests. I found an overall increase in coniferous species and several deciduous species (such as *Tilia*, *Populus*, *Betula*, *Alnus* sp), a spatial homogenization of species composition, and more even-aged stands. I also found a drop from 14% to 9% in the relative abundance of old forests (>100 years). Spikes in forest harvests coincided with times of widespread forest privatization, and drastic institutional changes, such as agrarian reforms, or the onset and collapse of the Soviet Regime. Overall, my results suggest that effects of past management, land ownership and institutional changes can persist for a long time, and affect forest ecosystem composition, health and structure. My findings are scientifically important because they provide evidence for legacies of past management and for the effects of forest privatization on harvesting rates. My findings are also relevant to forest management and conservation practice, because they highlight that environmentally sound management over long time periods is essential for sustainable forestry and old-growth forest protection in Europe and elsewhere.

Related manuscript: Munteanu C, Nita M-D, Abrudan I V., Radeloff VC (2016) Forest history is repeating itself, imposing strong legacies for Romanian forests and forestry. *Forest Ecology and Management*, 361: 179-193

Chapter 4: Land use legacies of historic political regimes on contemporary land abandonment.

Land-use legacies can shape landscapes for centuries into the future, affecting their structure and function - but for how long legacies may persist and whether they differ with historical political and socio-economic regimes remains unclear.

In Chapter 4, I assessed the effect of land use legacies for contemporary agricultural abandonment and studied the legacy effects of different political regimes after environmental variation was controlled for. The specific research questions were (1) to what extent agro-ecological conditions explained farming choices during major political regimes, (2) if legacy effects existed once agro-ecological variation was controlled for, and whether their effect diminished with time and (3) how legacies differed during different political and institutional regimes? To answer these questions, I modeled socialist (1960-1985) and post-socialist abandonment (1985-2010) in relation to legacies of three historic periods: the Habsburg (1860s), the Interwar (1930s) and the Socialist (1960s) periods. I controlled for agro-ecological, accessibility and socio-political variation. I found that Socialist agricultural expansion took place mostly in agro-ecologically unsuitable areas ($AUC = 0.70$), leading to subsequent abandonment both during the later socialist (34%) and the post-socialist periods (21%). Even after controlling for agro-ecological variability, I found that Habsburg land uses affected abandonment for as long as 100 years into the future, and that their effect diminished over time. However, I found differences when comparing legacies of different political regimes (Habsburg and Soviet). The

differences in land abandonment were greatest when comparing land farmed during the Habsburg versus the Socialist period. My results suggest that land use legacies are largest in times of rampant land use change, and that the underlying drivers of change such as institutional or political factors may contribute to these effects. My results also suggest that major shifts in land management systems can have ripple effects for land change even after a century. As globally land change is happening at ever faster rates, my study stresses the importance of considering our land use responsibility for future generation when making land use decisions.

Related manuscript: Munteanu C, Kuemmerle T, Boltiziar M, Halada L, Kaim D, Király G, Konkoly-Gyuró E, Kozak J, Lieskovsky J, Mojses M, Müller D, Ostafin K, Ostapowicz K, Radeloff VC (in preparation): Political regimes modulate the effect of land use legacies. (to be submitted: Regional Environmental Change).

Chapter 5: Conservation recommendations for the protection of bird habitat in the Carpathian Ecoregion in light long-term land use trends

Bird populations in Europe are declining and conservation action is needed to protect species and their habitats. Suggested conservation strategies include both wildlife friendly farming and rewilding. Despite the existence of empirical evidence on which species could benefit under each of these conservation strategies, the historical landscape context and the conservation responsibility of the Carpathians for different bird species remain largely unexplored, potentially leading to miss-directed conservation efforts.

In Chapter 5, I provide conservation recommendations for the Carpathian Mountains, based on historic and recent habitat evolution for species of highest conservation responsibility at European level. My specific research questions were: 1) How did habitat for different bird

species change in the Carpathian Mountains over the past 150 years? 2) For which of the bird species do the Carpathians carry the highest conservation responsibility at the European level? 3) What bird conservation strategies should be pursued, and what future land use trends would be most desirable for conservation, given the major threats to different species, their past habitat trends and the conservation responsibility of the Carpathians?

To address these questions, I analyzed long term land use trends since 1860 in relation to species range maps for 252 bird species whose European ranges at least partially cover the Carpathian ecoregion. I analyzed the major habitat and their changes in the Carpathians over the past 150 years, for all species present in the Carpathians, and evaluated for which ones the region carries highest conservation responsibility at the European level. Finally, based on observed habitat changes, conservation responsibility and species life history, I discussed conservation strategies and land management.

I found that forest and grassland habitat increased substantially since 1860 within the ranges of all Carpathian species, and that agricultural habitat declined. Overall, the Carpathians carry high conservation responsibility for species that use forests and grasslands as their major habitat and only low conservation responsibility for birds that rely on agricultural fields. Furthermore, the habitat requirements of species of high conservation responsibility indicated that several species would benefit from a mosaic of forest and grassland landscapes. The main concerns at European level for the survival of species of high conservation responsibility were agricultural intensification and natural system modification. Because I found that agricultural land declined substantially in our study region since the 1960s, I suggest that the Carpathians have high potential for conservation of forest and grassland species. I identified a list of 29 species for which the Carpathian ecoregion could become a conservation hotspot and suggest that land

management is focused on providing suitable habitat for these species, such as supporting forest recovery, low-intensity grassland management and promoting forest structure and diversity.

Related manuscript: Munteanu C, Pidgeon A, Radeloff V C (in preparation) Conservation recommendations for the protection of bird habitat in the Carpathian Ecoregion in light of long-term land use trends (to be submitted: Conservation Biology)

Related publications

Aside from this dissertation, I authored and co-authored several other papers related to the larger topic of long term land change in the Carpathian Region:

Munteanu C, Radeloff V, Griffiths P, et al. (2016) Land change in the Carpathian Region before and after major institutional changes. In: Gutman G, Radeloff VC (eds) Land Use and Land Cover Change in Eastern Europe after the Collapse of Socialism. Springer (in press)

In this book chapter, we synthesized the long term drivers of land change and their land-use outcomes in the Carpathian region, with a particular focus on forests, agriculture and grasslands. We provided evidence on how ecosystems respond to political shocks using examples of alternative stable states, time-lags and land-use legacies.

Butsic V, **Munteanu C**, Griffiths P, Knorn J, Radeloff V C, Lieskovský J, Müller D, Kuemmerle T (in review) The impact of protected areas on forest disturbance in the Carpathian Mountains 1985-2010. Conservation Biology

Here, we estimated the effectiveness of protected areas over time and space using forest disturbance data for 1985-2010, matching statistics and a fixed effects estimator. We found heterogeneous results in terms of protection effectiveness and suggest that the

strength of institutions, the differences in forest privatization and management and the timing of accession to the EU may provide explanations for these differences.

Feurdean A, **Munteanu C**, Kuemmerle T, Nielsen AB, Hutchinson SM, Ruprecht E, Persoiu A, Parr K, Hickler T (in preparation) Effects of socio-political transformations on semi-natural grasslands in Transylvania (CE Europe), based on pollen and historical land survey. (to be submitted: Regional Environmental Change)

For a study case in Transylvania, we analyzed land cover from historical maps in conjunction with a pollen record, in order to assess changes in grassland persistence and diversity over time. We found that although grassland extent increased over time, grassland persistence was low and diversity varied over time, likely in relation to land management policies of different political regimes. Our results suggest recent positive changes in grassland extent and diversity and the need for conservation action to maintain these landscapes.

Kaim D, Kozak J, Kolečka N, Ziółkowska E, Ostafin K, Ostapowicz K, Gimmi U, **Munteanu C**, Radeloff V (2016): Broad scale forest cover reconstruction from historical topographic maps. *Applied Geography*, 67: 39-48

In this paper we compare two methods for reconstructing forest cover based on historical maps: a classical wall-to-wall polygon digitizing approach and a regular sampling grid approach, similar to the one used in my dissertation. We found that the point-based reconstruction captured forest cover dynamics with a comparable accuracy to the wall-to-wall mapping, yet was much more time efficient. Our findings demonstrate that historic land change can be effectively assessed over larger areas much further back in time than commonly done.

Griffiths P, Kuemmerle T, Baumann M, Radeloff, VC, Abrudan, IV, Lieskovsky, J, **Munteanu C**, Ostapowicz K, Hostert P (2014) Forest disturbances, forest recover, and changes in forest types across the Carpathian ecoregion from 1985 to 2010 based on Landsat image composites. *Remote Sensing of Environment* 151:72–88.

In this paper, we analyzed changes in forest cover, forest disturbance and forest types for the Carpathian region, based on Landsat image 5-year composites between 1985 and 2010. We found that forest cover increased in the Carpathians by 4.4% and that forest composition shifted towards more deciduous. We also found a time-lag in the timing of disturbance, with Romania, Poland and Czech Republic experiencing disturbance later than other countries in the region.

Significance and implications

With an increasing world population and ever shifting dynamics in global economic and socio-political contexts, the effects of land use decisions for future land change become of increasing land management and conservation concern (Perring et al., 2016). My dissertation addresses broad questions on the role of land use legacies for subsequent land change and conservation, with particular focus on periods of political and socio-economic change. My research is – broadly speaking – relevant to land change science and land management and has wide implications for conservation. The Carpathian region represents the ideal ‘natural experiment’ for investigating effects of land use legacies, socio-political shifts, and the implications for conservation at broad spatial and temporal scales. But the results of my research are relevant to many landscapes with long history of human use and good land use records, such as the Eastern US or Sub-Saharan Africa. Similarly, my methods are straightforward, and can be applied to other regions and extrapolated to continental or global scales.

In this dissertation, I shed new light on the long term land changes and their drivers in the Carpathians, quantify land use legacies and provide conservation recommendations for bird species in light of their habitat changes. I showed that forest transition occurred in the Interwar period, and that contemporary forest cover is higher than historically. Similarly, an agricultural transition from increasing to decreasing cropland occurred in the 1960s and land abandonment was strong since. Historic land uses did affect rates of forest disturbance and agricultural abandonment, even after accounting for the classical determinants of land change. I found strongest legacy effects when comparing land abandonment on land farmed under different political regimes. Last but not least, I showed that decreasing land use intensity provides great conservation opportunities in the Carpathians – specifically for species of high conservation responsibility that require forest and grassland for their survival and for which habitat increased in the last century.

Scientifically, my work makes a major contribution to land change science, by quantifying the role of land use history and of path dependency for contemporary land change. By modeling change over long time periods, I advance the scientific and theoretical understanding of land use legacies as driving forces of land use change that may in turn be affected by underlying forces such as political regimes. Furthermore, the finding that the longer a given land use persisted, the less likely it was to transition to a different state (Chapters 2 and 4) bridges the gap between the theoretical concept of path dependency and the quantitative measure of land use legacy in contemporary landscapes. The concept of path dependency has been mostly tested in urban systems, and my research extends evidence of path dependency to forest and agricultural systems. Furthermore, I show that all else being equal, legacies may diminish over time, as expected from ecological succession theory (Chapter 2 and Chapter 4). However, legacy effects

can be stronger when comparing land change between political and institutional regimes which may abruptly affect historical land use decisions (Chapter 4). My results highlight the importance interactions between environmental and anthropogenic drivers of land use change at various spatial and temporal scales (Chapter 3).

In addition to elucidating legacies, my results verified the applicability of forest transition theory in an area that experienced multiple socio-economic and political shifts and showed that the timing of transition can vary between countries due to local and national differences in policies and institutional transitions (Chapters 1 and 2). My results are of important to land change science, because they show that in parallel to forest transition, a ‘mirrored agricultural transition’ occurred in the Carpathians – agricultural expansion peaked with a delay compared to the forest transition point and continued throughout the contemporary period (Chapter 4).

Last but not least, because Eastern Europe has “reinvented itself every half century” in terms of politics and socio-economics (Good, 1994), my findings from the Carpathians can contribute more broadly to the understanding of land change dynamics of regions affected by multiple socio-economic and socio-ecological changes. My findings showing accelerated rates of change in relation to political shifts are relevant to any regions that are currently hot-spots of human pressure on the environment due to wars, political unrest (e.g., the Middle East) or large development projects (e.g., China). For the Carpathian region specifically, this work represents the first comprehensive land change assessment over broad temporal and spatial scales, integrating and synthesizing most of the previous knowledge on the region.

From a methodological perspective, my work utilized straightforward approaches to answer broad questions relevant to land change science. The meta-analytical approach I used in Chapter

1 allows for harmonization and comparability of historical land use data from different sources. In Chapters 2 and 4, I used a novel approach to quantify and model historic land uses. Because the efficiency in analyzing historical data has been a major barrier to broad-scale historical analysis so far, I employed a point grid approach for mapping land use, which proved to be more time-efficient and equally reliable to traditional approaches (Kaim et al., 2016). This approach could be used at different scales and for any type of spatial datasets. Having at hand a method to assess historic land change based on this point-grid approach provides great new opportunities for scientific inquiry related to historic land use across a large part of the world for which historical spatial data is available. The straightforward methods I applied in Chapter 5 to assess conservation responsibility and outline conservation goals can be easily adapted to other species or eco-regions because they rely on comparable, globally available data.

In addition to new mapping and spatial analysis methods, I developed statistical models that quantified the role of historic land use for recent land change and explored their relation to other drivers of change (Chapter 2, Chapter 4). My models are easily applicable to any area for which spatially explicit long term datasets of land cover are available and can be used to reveal the relative importance of past uses for recent changes. Areas where the implementation of my models would be particularly interesting include regions which underwent multiple land use transitions in the past, driven by changes of political, socio-demographic, economic or environmental pressures such as the Midwestern U.S., the Middle and Far East, and South-Central and Eastern European countries.

Last but not least, my results are relevant in the context of land management and conservation. Three broad management and conservation messages emerge from my work. First the changes in contemporary landscapes are affected by land use history, and thus historic landscape conditions,

should be considered when making contemporary land use and conservation decisions. For instance, prioritizing areas for conservation might consider constraints imposed by historic land use. In turn, knowing the constraints on a specific landscape can help mitigate the legacy effects when making land management decisions. Second, my results highlight the need for making farsighted land management and conservation decisions, because we bear great land use responsibility for generations to come. Furthermore, because land use legacies may not affect just rates of land change into the future, but also species population dynamics (e.g. extinction debt), considering the cascading effects of contemporary land use decision is of utmost importance for land management and conservation. Third, providing conservation goals based on long term habitat dynamics and the conservation responsibility of specific regions, may enhance conservation outcomes. My results suggest that decreasing land use intensity provides great opportunities for conservation, as long as appropriate conservation policies are implemented.

For the Carpathian region, my datasets are directly relevant for management and planning because they represents the first a spatially explicit and cross-border account of land change for the past century. These datasets can inform forestry and agricultural planning and highlight areas suitable for conservation, such as protected areas. In terms of conservation action, my analysis of habitat dynamics for birds of high conservation responsibility suggests that conservation should focus on forest and grassland bird species, for which habitat has been increasing over the last century. Because the Carpathians do not carry conservation responsibility for cropland bird species, we suggest that allowing the remaining agricultural lands to convert to grasslands would benefit biodiversity at European level more than maintaining low intensity farming on the landscape. Last but not least, my study provides an important basis for the emerging cross-border

conservation and management efforts in the Carpathians, because it is consistent and comparable across six countries.

References

- Barbier, E.B., Burgess, J.C., Grainger, A., 2010. The forest transition: Towards a more comprehensive theoretical framework. *Land use policy* 27, 98–107.
- Baumann, M., Kuemmerle, T., Elbakidze, M., Ozdogan, M., Radeloff, V.C., Keuler, N.S., Prishchepov, A. V., Kruhlov, I., Hostert, P., 2011. Patterns and drivers of post-socialist farmland abandonment in Western Ukraine. *Land use policy* 28, 552–562.
- Börjeson, L., 2007. Boserup backwards? Agricultural intensification as “its own driving force” in the Mbulu Highlands, Tanzania. *Geogr Ann Ser B Hum Geogr* 89, 249–267.
- Brudvig, L.A., Grman, E., Habeck, C.W., Orrock, J.L., Ledvina, J.A., 2013. Strong legacy of agricultural land use on soils and understory plant communities in longleaf pine woodlands. *For Ecol Manage* 310, 944–955.
- Bürgi, M., Hersperger, A.M., Schneeberger, N., 2005. Driving forces of landscape change - current and new directions. *Landsc Ecol* 19, 857–868.
- Ceaușu, S., Hofmann, M., Navarro, L.M., Carver, S., Verburg, P.H., Pereira, H.M., 2015. Mapping opportunities and challenges for rewilding in Europe. *Conserv Biol* 00, 1–11.
- Coomes, O.T., Takasaki, Y., Rhemtulla, J.M., 2011. Land-use poverty traps identified in shifting cultivation systems shape long-term tropical forest cover. *Proc Natl Acad Sci U S A* 108, 13925–30.
- DeFries, R.S., Foley, J.A., Asner, G.P., 2004. Land-use choices : balancing human needs and ecosystem function In a nutshell : *Front Ecol* 2, 249–257.
- Dullinger, S., Essl, F., Rabitsch, W., Erb, K.-H., Gingrich, S., Haberl, H., Hülber, K., Jarosík, V., Krausmann, F., Kühn, I., Pergl, J., Pyšek, P., Hulme, P.E., 2013. Europe’s other debt crisis caused by the long legacy of future extinctions. *Proc Natl Acad Sci U S A* 110, 7342–7.
- Essl, F., Dullinger, S., Rabitsch, W., Hulme, P.E., Hülber, K., Jarošík, V., Kleinbauer, I., Krausmann, F., Kühn, I., Nentwig, W., Vilà, M., Genovesi, P., Gherardi, F., Desprez-Loustau, M.-L., Roques, A., Pyšek, P., 2011. Socioeconomic legacy yields an invasion debt. *Proc Natl Acad Sci U S A* 108, 203–7.
- Estel, S., Kuemmerle, T., Alcántara, C., Levers, C., Prishchepov, A., Hostert, P., 2015. Mapping farmland abandonment and recultivation across Europe using MODIS NDVI time series. *Remote Sens Environ* 163, 312–325.
- Ferretti-Gallon, K., Busch, J., 2014. What Drives Deforestation and What Stops It? A Meta-Analysis of Spatially Explicit Econometric Studies. Washington DC.

- Ficetola, G.F., Maiorano, L., Fallucci, A., Dendoncker, N., Boitani, L., Padoa-Schioppa, E., Miaud, C., Thullier, W., 2010. Knowing the past to predict the future: land-use change and the distribution of invasive bullfrogs. *Glob Chang Biol* 16, 528–537.
- Fischer, J., Abson, D.J., Butsic, V., Chappell, M.J., Ekroos, J., Hanspach, J., Kuemmerle, T., Smith, H.G., von Wehrden, H., 2014. Land sparing versus land sharing: moving forward. *Conserv Lett* 7.
- Fischer, J., Hartel, T., Kuemmerle, T., 2012. Conservation policy in traditional farming landscapes. *Conserv Lett* 5, 167–175.
- Foley, J.A., DeFries, R.S., Asner, G.P., Barford, C., Bonan, G., Carpenter, S.R., Chapin, F.S., Coe, M.T., Daily, G.C., Gibbs, H.K., Helkowski, J.H., Holloway, T., Howard, E.A., Kucharik, C.J., Monfreda, C., Patz, J.A., Prentice, I.C., Ramankutty, N., Snyder, P.K., 2005. Global consequences of land use. *Science* (80-) 309, 570–574.
- Foley, J.A., Ramankutty, N., Brauman, K. a, Cassidy, E.S., Gerber, J.S., Johnston, M., Mueller, N.D., O’Connell, C., Ray, D.K., West, P.C., Balzer, C., Bennett, E.M., Carpenter, S.R., Hill, J., Monfreda, C., Polasky, S., Rockström, J., Sheehan, J., Siebert, S., Tilman, D., Zaks, D.P.M., 2011. Solutions for a cultivated planet. *Nature* 478, 337–42.
- Foster, D.R., Swanson, F.J., Aber, J., Burke, I., Brokaw, N., Tilman, D., Knapp, A., 2003. The importance of land-use legacies to ecology and conservation. *Bioscience* 53, 77–88.
- Geist, H.J., Lambin, E.F., 2001. What Drives Tropical Deforestation? A meta-analysis of proximate and underlying causes of deforestation based on subnational case study evidence., LUCC Report Series. Louvain-la-Neuve.
- Geist, H.J., Lambin, E.F., 2002. Proximate Causes and Underlying Driving Forces of Tropical Deforestation. *Bioscience* 52, 143.
- Gibbs, H.K., Herold, M., 2007. Tropical deforestation and greenhouse gas emissions. *Environ Res Lett* 2, 045021.
- Gibbs, H.K., Ruesch, A.S., Achard, F., Clayton, M.K., Holmgren, P., Ramankutty, N., Foley, J.A., 2010. Tropical forests were the primary sources of new agricultural land in the 1980s and 1990s. *Proc Natl Acad Sci U S A* 107, 16732–16737.
- Good, D.F., 1994. Economic transformation in East and Central Europe. Legacies from the Past and Policies for the Future. Routledge, London, New York.
- Grau, R., Kuemmerle, T., Macchi, L., 2013. Beyond “land sparing versus land sharing”: environmental heterogeneity, globalization and the balance between agricultural production and nature conservation. *Curr Opin Environ Sustain* 5, 477–483.
- Hansen, M.C., Stehman, S. V, Potapov, P. V, 2010. Quantification of global gross forest cover loss. *Proc Natl Acad Sci U S A* 107, 8650–5.

- Hartel, T., Dorresteijn, I., Klein, C., Máthé, O., Moga, C.I., Öllerer, K., Roellig, M., von Wehrden, H., Fischer, J., 2013. Wood-pastures in a traditional rural region of Eastern Europe: Characteristics, management and status. *Biol Conserv* 166, 267–275.
- Kaim, D., Kozak, J., Kolečka, N., Ziółkowska, E., Ostafin, K., Ostapowicz, K., Gimmi, U., Munteanu, C., Radeloff, V.C., 2016. Broad scale forest cover reconstruction from historical topographic maps. *Appl Geogr* 67, 39–48.
- Knorn, J., Kuemmerle, T., Radeloff, V.C., Keeton, W.S., Gancz, V., Biriş, I.-A., Svoboda, M., Griffiths, P., Hagatis, A., Hostert, P., 2012. Continued loss of temperate old-growth forests in the Romanian Carpathians despite an increasing protected area network. *Environ Conserv* 40, 182–193.
- Kraemer, R., Prishchepov, A. V., Müller, D., Kuemmerle, T., Radeloff, V.C., Dara, A., Terekhov, A., Frühauf, M., 2015. Long-term agricultural land-cover change and potential for cropland expansion in the former Virgin Lands area of Kazakhstan. *Environ Res Lett* 10, 054012.
- Kuemmerle, T., Hostert, P., Radeloff, V.C., Perzanowski, K., Kruhlov, I., 2007. Post-socialist forest disturbance in the Carpathian border region of Poland, Slovakia, and Ukraine. *Ecol Appl* 17, 1279–95.
- Kuemmerle, T., Perzanowski, K., Chaskovskyy, O., Ostapowicz, K., Halada, L., Bashta, A.-T., Kruhlov, I., Hostert, P., Waller, D.M., Radeloff, V.C., 2010. European Bison habitat in the Carpathian Mountains. *Biol Conserv* 143, 908–916.
- Lambin, E.F., Geist, H.J. (Eds.), 2006. *Land-Use and Land-Cover Change - Local Processes and Global Impacts*, Global Cha. ed. Springer.
- Lambin, E.F., Meyfroidt, P., 2010. Land use transitions: Socio-ecological feedback versus socio-economic change. *Land use policy* 27, 108–118.
- Lambin, E.F., Turner, B.L., Geist, H.J., Agbola, S.B., Angelsen, A., Bruce, J.W., Coomes, O.T., Dirzo, R., Fischer, G., Folke, C., George, P.S., Homewood, K., Imbernon, J., Leemans, R., Li, X., Moran, E.F., Mortimore, M., Ramakrishnan, P.S., Richards, J.F., Skånes, H., Steffen, W., Stone, G.D., Svedin, U., Veldkamp, T.A., Vogel, C., Xu, J., 2001. The causes of land-use and land-cover change: moving beyond the myths. *Glob Environ Chang* 11, 261–269.
- Mather, A.S., 1992. The forest transition. *Area* 24, 367–379.
- Meyfroidt, P., 2015. Approaches and terminology for causal analysis in land systems science. *J Land Use Sci* 4248, 1–27.
- Meyfroidt, P., Lambin, E.F., 2011. Global Forest Transition: Prospects for an End to Deforestation. *Annu Rev Environ Resour* 36, 343–371.
- Mikulcak, F., Newig, J., Milcu, A.I., Hartel, T., Fischer, J., 2013. Integrating Rural Development and Biodiversity Conservation in Central Romania. *Environ Conserv* 40, 129–137.

- Müller, D., Leitão, P.J., Sikor, T., 2013. Comparing the determinants of cropland abandonment in Albania and Romania using boosted regression trees. *Agric Syst* 117, 66–77.
- Munroe, D.K., van Berkel, D.B., Verburg, P.H., Olson, J.L., 2013. Alternative trajectories of land abandonment: causes, consequences and research challenges. *Curr Opin Environ Sustain* 1–6.
- Navarro, L.M., Pereira, H.M., 2012. Rewilding Abandoned Landscapes in Europe. *Ecosystems* 15, 900–912.
- Ojima, D.S., Galvin, K.A., Turner, B.L., 1994. The global impact of land-use change. *Bioscience* 44, 300–304.
- Perring, M.P., De Frenne, P., Baeten, L., Maes, S.L., Depauw, L., Blondeel, H., Carón, M.M., Verheyen, K., 2016. Global environmental change effects on ecosystems: the importance of land-use legacies. *Glob Chang Biol* 22, 1361–1371.
- Phalan, B., Onial, M., Balmford, A., Green, R.E., 2011. Reconciling food production and biodiversity conservation: land sharing and land sparing compared. *Science* 333, 1289–1291.
- Pidgeon, A.M., Mathews, N.E., Benoit, R., Nordheim, E. V, 2011. Response of in Avian Historic Habitat Change the Northern 15, 1772–1788.
- Radeloff, V.C., Hammer, R.B., Voss, P.R., Hagen, A.E., Field, D.R., Mladenoff, D.J., 2001. Human Demographic Trends and Landscape Level Forest Management in the Northwest Wisconsin Pine Barrens. *For Sci* 47, 229–241.
- Seto, K.C., Reenberg, A., Boone, C.G., Fragkias, M., Haase, D., Langanke, T., Marcotullio, P., Munroe, D.K., Olah, B., Simon, D., 2012. Urban land teleconnections and sustainability. *Proc Natl Acad Sci U S A* 109, 7687–92.
- Tilman, D., May, R.M., Lehman, C.L., Nowak, M.A., 1994. Habitat destruction and the extinction debt. *Nature* 371, 65–66.
- Turner, B.L., Janetos, A.C., Verburg, P.H., Murray, A.T., 2013. Land system architecture: Using land systems to adapt and mitigate global environmental change. *Glob Environ Chang* 23, 395–397.
- Verburg, P.H., Erb, K.-H., Mertz, O., Espindola, G., 2013. Land System Science: between global challenges and local realities. *Curr Opin Environ Sustain* 5, 1–5.
- Wallin, D.O., Swanson, F.J., Marks, B., 1994. Landscape Pattern Response to Changes in Pattern Generation Rules: Land-Use Legacies in Forestry. *Ecol Appl* 4, 569–580.

Chapter 1. Forest and agricultural land change in the Carpathian region – a meta-analysis of long-term patterns and drivers of change

Introduction

Land-cover change is a main component of global environmental change (Foley et al., 2005), affecting climate, biodiversity and ecosystem services, which in turn, affect land-use decisions (Ojima et al., 1994). Humans have altered land cover for centuries, but recent rates of change are higher than ever (Foley et al., 2005; Goldewijk, 2001; Hansen et al., 2010). The temporal dimension of change is particularly interesting because land-use legacies may persist for centuries (Foster et al., 2003). Over long time periods though, land-use transition theories predict gradual changes, primarily as a function of demographic and economic factors (DeFries et al., 2004; Foley et al., 2005). For example, forest transition theory postulates that gradual economic and demographic change leads to agricultural specialization and reforestation of marginal lands, and defines the transition point as the time of the lowest forest cover in a given country or region (Mather, 1992; Meyfroidt and Lambin, 2011). Different regions may experience these transitions at different points in time, depending on economic, political or institutional condition (Meyfroidt and Lambin, 2011) or go through multiple transition phases (Yeo and Huang, 2013), as land systems respond to institutional and economic changes (Lambin et al., 2001). Shifts in political systems, and the related socio-economic adaptations, such as those following the collapse of the Soviet Union, greatly affect land trends (Hostert et al., 2011). The question is though first, how land cover changes over long time periods, and how these changes vary depending on economic, institutional and social factors. Regional land change patterns are the combined result of changes at much finer scale, that are driven by complex economic, policy and institutional, demographic and market forces (Lambin and Meyfroidt, 2010; Verburg et al., 2009). These localized changes,

in turn, are constrained by interacting broad- and local-scale driving forces, especially in crisis situations (Cioroianu, 2007). While, the local-scale drivers of land-use change can be understood from case-studies (Foley et al., 2005), the variation of these drivers across regions can only be understood from a broader perspective.

Capturing land change under successive distinct economic periods and documenting change processes over large areas and long time periods (e.g., centuries) is often impossible due to the lack of consistent, broad-scale and long-term data. When that is the case, a meta-analysis can be a valuable tool for synthesizing knowledge and extracting broader scale patterns and drivers of change (Poteete and Ostrom, 2008; Rudel, 2008). Meta-analyses have been applied to assess, for example, long-term urban growth across the globe (Seto et al., 2011), desertification (Geist and Lambin, 2004), deforestation (Geist and Lambin, 2002), and tropical agriculture (Keys and McConnell, 2005). In regard to forest change, such a meta-analysis showed that tropical deforestation is a result of interacting proximate causes and underlying driving forces, which vary geographically and with historical context (Geist and Lambin 2002). Another meta-analysis focusing on forest cover in Mexico showed that cattle ranching and outmigration cause deforestation in lowland areas, while highland regions with outmigration experience forest cover increase (Rudel, 2008). Dryland degradation globally has been attributed to the combined effects of climate, economies and institutions which drive cropland expansion, overgrazing and infrastructure development (Geist and Lambin, 2004). In Central Eastern Europe, Kozak (2010) analyzed land change across a number of local case studies to describe forest transition in the Polish Carpathian Mountains as occurring between the two World Wars (WW). However, while most meta-analyses examined broad spatial extents and explain spatial variation, their temporal scale has been limited to decades, which limits the ability to isolate effects and legacies of major

socio-economic shifts across time and space. Furthermore, most meta-analyses of land change processes included only case studies that were published in English (Geist and Lambin, 2004; McConnell and Keys, 2005; Seto et al., 2011), thus not including local research and knowledge.

Broad scale, long term comparative studies across countries of Eastern Europe are still lacking (Bjørnsen-Gurung et al., 2009), despite the availability of a high number of local, regionally published studies. Given its long land-use history and multiple social, political and economic shocks, the Carpathian region represents a “natural experiment” (Gehlbach and Malesky, n.d.) to examine long-term land-use change and to develop a broader synthesis of land-use histories.

Our overall goal was to identify and quantify broad-scale and long-term land change patterns and processes during times of shocks, and the main driving forces of these changes. To do so, we conducted a meta-analysis of historical land change studies for the Carpathian region, reaching as far back as 1790s.

Specifically, our objectives were to:

1. Assess and quantify the main forest and agricultural changes in the Carpathian region for politically and economically distinct time periods over the past 250 years;
2. Assess the heterogeneity of the local-scale studies across the region;
3. Identify the main drivers of long-term land-use change and the impact of major socio-economic shocks on forest and agricultural change.

Methods

Study area

We studied the 350,000 km² Carpathian region in Eastern Europe, which comprises two major

eco-regions: the Carpathian Mountains and the Pannonian Plains. The study area includes parts of the Czech Republic, Poland, Ukraine and Romania, and all of Hungary and Slovakia (Figure 1.1), has a temperate climate, and landscapes consisting mostly of a mosaic of forests, pastures, and agricultural fields. The region harbors some of the largest contiguous temperate forests in Europe (Knorn et al., 2009; Kuemmerle et al., 2007) alongside high nature conservation value farmland (Paracchini et al., 2008). The Pannonian plains also represent one of the most fertile regions in Europe (Schiller et al., 2010). The Carpathian eco-region is a global biodiversity hotspot, particularly regarding plant diversity, and harbors rare old-growth and alpine meadow ecosystems and many wildlife species of conservation concern (e.g., brown bear, wolf, lynx, European bison, (Salvatori et al., 2002).

The region has a long land-use history, with centuries of agricultural and forest land use being influenced by changes in political, economic and demographic dynamics (Verburg et al., 2009). Land-cover changes during recent decades (since 1980s), have been captured by remote sensing analyses of the entire region, and showed overall increases in forest cover and agricultural abandonment (Griffiths et al., 2014; Kuemmerle et al., 2008). However, our understanding of long-term land-use trends remains scattered across numerous local-scale case-studies dispersed across the region (e.g., Feranec and Otáhel, 2009; Kaim, 2009; Ostafin, 2009) and a synthesis of these studies is lacking.

Theoretical land change predictions

In order to understand land-use trends in the region, we examined agricultural and forest change during five historical periods with distinctive socio-economic, political, technological and cultural characteristics, that were demarcated by several large-scale shocks: (1) the Habsburg and

Austro-Hungarian Empires (K.u.K. Monarchy), a time of agricultural modernization and the beginning of the industrial revolution, which ended with World War I (WW I), (2) the Interwar period, characterized by the emergence of several nation-states, industrialization and intensification up to World War II (WW II), (3) the Socialist period, defined by intensification, centrally planned economies and land reforms leading to nationalization and collectivization, which ended around 1990 in the Carpathian countries, (4) the Transition when countries established market economies and land reforms took place, which lasted roughly until 2000, and lastly (5) the Accession of most countries to the EU in either 2004 and 2007 (except Ukraine), a period influenced by EU's trade, agricultural, and economic policies. We considered this last time period to start in 2000 because that is when most countries already adjusted their regulations and legislation according to European standards.

We formulated a set of expected land change trends for each period, based on overall socio-economic and technological trends during the respective time periods. Specifically, we predicted that the expansion of the Austro-Hungarian Empire, and the industrial revolution, led to homogenization and specialization of land use practices (Bičík et al., 2001), manifested as an increase in agricultural lands and a decrease in forest cover up to WW II. Under socialism, natural resource use intensified (Cioroianu, 2007), and we hypothesized that agricultural expansion, intensification and in some countries collectivization, led to an increase of agricultural land, especially in lowland areas, while abandonment of marginal lands allowed for forest recovery. We expected that the collapse of socialism, followed by the changes in ownership structure and the establishment of market economies led to a decrease in agricultural lands due to abandonment and forest re-growth during the Transition period (Baumann et al., 2011; Kuemmerle et al., 2009a, 2009d; Prishchepov et al., 2013). We also expected that the

effects of the EU macro-economic policies would result in the increase of agricultural lands due to subsidies (Björnsen-Gurung et al., 2009) and continued forest transition (Lambin and Meyfroidt, 2010). We hypothesized thus increases in both agricultural lands (over previously abandoned fields) and forest cover after 2000 (Csaki and Jambor, 2009). For a summary of these expected trends, see Table 1.1.

Data

We collected case study information on forest and agricultural change both from peer-reviewed articles and grey literature. We used Google Scholar and regional scientific databases using combinations of “historic”, “land-use/ land-cover change”, and “maps” in English and the regional languages (Romanian, Slovakian, and Ukrainian) and complemented this information with traditional library research in the respective countries plus references from local experts in the Czech Republic, Hungary, Ukraine, Poland, Romania and Slovakia. For 85 publications, we extracted information about the study area, land cover at different time periods, and the main drivers of change. In approximately half of the cases, data was provided directly by authors of the paper. For the remaining publications we extracted the data using a structured form. From the total of 85 publications, we selected and analyzed those 66 papers (listed in the Appendix 1) that (1) were based on spatially-explicit data (historic maps, aerial photographs, and satellite imagery) and not only statistics; (2) examined land cover at least two points in time, and (3) included spatial data regarding the study location or coordinates of the study region. We excluded papers that did not meet at least one of the three criteria (Appendix 2). We hereafter refer to a case study as being a single geographical location at which either forest or agricultural (or both) land cover was reported during a given time period. Some papers contained several case studies, reporting land-cover change in multiple locations. We did not include in our meta-

analysis recent broad-scale remote sensing land change studies based on Landsat data (Alcantara et al., 2012; Baumann et al., 2011; Griffiths et al., 2014; Knorn et al., 2012b, 2009; Kuemmerle et al., 2011, 2009a, 2008, 2007; Prishchepov et al., 2012), due to their significantly different spatial extent, frequent overlap among studies, and generally short duration, but we considered them in the discussion of change drivers. In sum, the 66 papers contained a total of 102 case study locations, for which change rates were calculated for one or more time-periods (Figure 1.1 and Table 1.1). The spatial extent of the individual studies spans from 240 ha (Wolski, 2001) to over 2 million hectares (Grekov, 2002), with the median at 12,500 ha (Figure 1.1b). On a temporal scale, the shortest time period covered by case studies at a single location was 10 years (Grekov, 2002) , and the longest 238 years (Mojsey and Boltiziar, 2011). The mean time covered was 105 years.

Analysis

We developed a common land-cover class catalogue, which was applied to all studies. In most instances, this necessitated the aggregation of classes (e.g., ‘permanent’ and ‘seasonal crops’ were combined into ‘agriculture’). The final product was land-cover data for ‘forest’, ‘agriculture’ and ‘other’ land covers. We calculated the annual rate of change for each land-cover class following the model of FAO forest change assessments (Pandey, 1995) which uses a formula based on the compound interest law in order to compare among sites ((Puyravaud, 2003):

$$Ann_{change} = \left(\frac{A_2}{A_1} \right)^{1/(t_2-t_1)} - 1$$

(Equation 1)

A_1 and A_2 represent the area of land cover of interest (forest or agriculture land) at the times t_1 and t_2 . When a case study reported multiple rates within one of the five analyzed time-periods, we calculated weighted averages. Studies that reported a single rate of change across multiple time periods were mapped using a different symbol, as these depict change only between the beginning and end of the first and last period, missing variation within the selected time window. We defined change rates between $\pm 0.1\%$ change/year as ‘stable’ land use. Centroids were digitized to represent the location of each study and rates of change were calculated for each study and time period under investigation (Figure 1.1).

To identify the main drivers, we conducted a qualitative review, categorizing the major types of driving forces as suggested by Geist and Lambin (2004) and Bürgi et al. (2005): institutional, economic, social-demographic, cultural, and climatic. Because our analysis only captured changes in land cover and not in land-use intensity, technological drivers, such as the introduction of fertilizers, or mechanization, which would mostly lead to increased yields or crop rotation, were considered jointly with the economic factors. For each case study, we identified the two most important drivers of change as described by paper authors and regional experts. We counted the number studies that mentioned each driver and qualitatively reviewed each driver across case-studies and the four land change processes of interest (deforestation, reforestation, agricultural expansion, and agricultural abandonment).

Case study representativeness and robustness check

The case studies ranged widely in extent (240 ha to 2.1 million ha) and duration (from 2 to 180 years). We tested for correlation between the absolute values of the annual rate of change and (a) the size of study area, (b) the temporal extent of the studies and (c) the percentage cover at the

beginning of the study, but found only weak associations (adjusted R-squares of 0.036, 0.018, and 0.033 respectively). Spatially, land change research was concentrated in the Carpathian Mountains, while lowland areas were underrepresented, except in Ukraine. The highest density of studies was in Poland and Slovakia (Figure 1.1). Since 2000, case-studies on agricultural and forest change were relatively sparse due to the short time period under consideration (12 years).

In order to check if case studies represented the general conditions of the respective country's share of the study area, we examined three physical variables, mean elevation, mean slope, and dominant soil type, for each case study and compared mean values of the case-studies with the mean of country's share of the study area. We found that the dominant soil across all countries was Cambisol, as was the case for most of the case-studies, except in Hungary where Luvisols and Fluvisols were overrepresented (Figure 1.2). In terms of slope and elevation, case studies in the Czech Republic, Slovakia and Ukraine studies represented their country's physical conditions well. In Poland and Romania, many studies were carried out at higher-than-average elevations and slopes, but the means for the country's share of the study area fall close to the 1st quartile of the case studies distribution in all cases (Figure 1.2).

Results

Forest cover increase was the most common land-cover change over the past 250 years in the majority of studies. Among the time periods, we found the highest proportion of case studies reporting decreases in forest cover during the K.u.K. Monarchy (over 22% of studies). However, even this period, stable forest cover was the most common pattern (mean annual change +0.08%). Forest cover increased during all other periods, especially during the Transition and EU period (mean annual change +1.07% and +0.89%). In the Interwar period, 92% of studies reported stable or increasing forest cover (mean annual change 0.35%, Figure 1.3). A high

proportion of studies reported forest cover increase (65%) for the Socialist period, in particular in the northern part of the Carpathians (annual mean 0.33%), followed by continuing increasing forest cover during Transition and EU accession periods (73 % and 72% respectively). After 2000, forest cover increased (annual mean 0.89%), but in Romania we found high rates of forest cover loss (Figure 1.3 and Figure 1.4).

Agricultural change was generally complementary to forest change, where forests increased, agriculture decreased, and vice-versa. However, during the K.u.K Monarchy period, agriculture increased (70% of studies, mean annual increase of 0.12%), while forest cover was mostly stable (45% of studies), indicating agricultural expansion into other land covers (Figure 1.3, Figure 1.5). The mean annual change of agricultural land change during the Interwar period was -1.28%, despite relatively stable agricultural cover (55% of studies, $\pm 0.1\%$ annual change) reported in most studies. After 1945, most studies ($> 75\%$) reported a decrease of agricultural land-cover. During the Transition and EU accession periods, there were substantial decreases in agricultural cover (mean annual change of -1.61% and -1.20% respectively). Across time periods, the proportion of studies documenting loss of agricultural land increased constantly until 2000, but dropped slightly after the EU accession (Figure 1.3).

There were interesting regional patterns of change though: forest decreased during the K.u.K. Monarchy in the Romanian, Ukrainian, and Slovakian Carpathians, while it increased in the Polish Carpathians, and was stable in the Czech Republic ($\pm 0.1\%$ annual change). During the Interwar period the majority of the forest change case-studies (48%) reported stable or increasing (44%) forest cover ($>0.1\%$ annual change) but most of them were in Slovakia, and Poland, while cases of forest loss occurred in Hungary, Romania and Slovakia. Thus, across the region, forest transition occurred during the Interwar period, though we caution that patterns at elevations over

1000m in Ukraine and Romania were different (Shandra et al., 2013). The most rapid forest increase during the Socialist period occurred in the border region between Poland, Ukraine, and Slovakia (Figure 1.4), while deforestation occurred in lowland areas (e.g., Hungary) as well as in the mining district of central Slovakia. After 1990, forest cover increased across Poland, Slovakia, Czech Republic and Hungary, but there were still cases of forest loss in the Eastern Romanian Carpathians and southwestern Slovakia.

Agricultural change varied regionally: during the K.u.K. Monarchy, agriculture expanded mostly in the lowlands of Hungary, Czech Republic, and Ukraine, concurrent with forest loss, while agriculture decreased in the mountains of Slovakia and Poland. In the Interwar period agricultural land use peaked in parts of Hungary and southwest Slovakia, while agriculture declined in parts of the Polish Carpathians and northern Slovakia. During the Socialist time period, low but positive annual rates of agricultural expansion occurred in Romania and southeast of Hungary. In Slovakia and the Czech Republic, agriculture decreased slowly, whereas in Poland, agricultural land decreased by up to 5% per year (Woś, 2005). Since 1990s, agricultural decrease was least pronounced in the lowlands of Hungary, Ukraine, and the Czech Republic. In mountain areas, lower abandonment rates were reported in Ukraine, contrasting with higher rates for Romania and Slovakia (Figure 1.5). Since 2000 agriculture declined in 69% of the studies, but we caution that there are only few studies for this period.

Our analysis of the main drivers of land change examined the number of times at least one of the selected drivers of change (institutional, economic, socio-demographic, cultural, and climatic) was deemed important by the case-study authors and collaborators for each of the change processes. We found that institutional and economic factors were the most important drivers of agricultural expansion and deforestation, jointly accounting for more than 75% and 65%

respectively of the case studies. This class of drivers also included the technological developments that led to agricultural intensification and support forest transition, but our focus on land-cover areas did not allow to examine technological drivers in detail. In contrast, socio-demographic factors like migration or sector employment were more important for agricultural abandonment (42% of cases) and forest succession (36% of cases, Figure 1.6). Physical factors were also mentioned as drivers of change, for example climate supported forest succession on abandoned mountain pastures, where the timberline shifted to higher altitudes (Mihai et al., 2006; Shandra et al., 2013). Overall, abandonment of agriculture was largely driven by socio-demographic (42%) and institutional (31%) factors, with the economy playing a less important role (24%) (Figure 1.6).

Discussion

We identified temporal and spatial patterns of land-cover change and their driving forces over the last 250 years across the Carpathian Basin. Our results showed that forest change was closely related to agricultural dynamics and that rates and patterns of change were heterogeneous among politically distinct time periods, and varied regionally. Deforestation was less widespread than we had expected, and the observed changes differed from our expectations in particular during the K.u.K. Monarchy and Interwar periods. Between WW I and WW II, forest cover decrease stopped across the region. Our findings are concurrent with other studies (Kozak, 2003; Kuemmerle et al., 2011), indicating that the region as a whole experienced a forest transition during the Interwar period, despite regional differences (Shandra et al., 2013). After WW II, the observed forest cover increase was in line with our expectations (Table 1.2). While agricultural abandonment was widespread throughout the 20th century, increase in agricultural cover occurred only during the K.u.K. Monarchy. Contrary to our expectations, agricultural

abandonment started early, being a prominent process across the region already during the Interwar and Socialist periods. However, abandonment rates increased after the collapse of the Socialism. In general, forest and agricultural dynamics were complementary, but there were exceptions to this rule due to rapid urban or grassland-related land-cover changes. Agricultural expansion and deforestation were mostly driven by economic and political events, while land abandonment and reforestation were mostly driven by socio-demographic factors.

Our analysis highlighted regional variation in land change patterns, and in the major drivers of change across the study area. We primarily focused on patterns of two broad scale processes: deforestation followed by agricultural expansion and forest cover increase, related to agricultural abandonment. The rise of the Habsburg Empire and Austro-Hungarian Monarchy, which brought German settlers to the Carpathian region, and the industrial revolution of the 19th century, caused significant population growth, increasing demands for agricultural products (Vepryk, 2002). Deforestation for agricultural development was both an economic and a cultural process (Boltižiar and Chrastina, 2006; Mojses and Boltižiar, 2011; Skokanová et al., 2012), and as such, patterns of deforestation varied by land ownership. While Ukrainian smallholders cleared forest patches for agricultural use in lowland areas, large landowners did not deforest, but replaced mixed forest stands with spruce plantations for pulp production at high elevations (Vepryk, 2001). While forest clearing for agriculture was common (Chrastina and Boltižiar, 2010; Konkoly-Gyuró et al., 2011; Vepryk, 2002), deforestation was also related to expanding grassland and urban cover. For example, on Ukrainian mountain meadows, livestock farming increased partly due to Hungarian and Czech investment up to WWII, lowering the timberline (Sitko and Troll, 2008). In the Northern Romanian Carpathians, net forest cover decreased at

timberline since 1880s, but generally net forest cover increased at timberline due to decline of transhumance (Shandra et al., 2013).

Similarly, economic growth led to the drainage of wetlands for agriculture in Hungary (Biró et al., 2012; Konkoly-Gyuró et al., 2011; Nagy, 2008), the Czech Republic, and Slovakia (Demek et al., 2008; Drgona, 2004; Gerard et al., 2010, 2006b; Mojses and Bezák, 2010) and to the conversion of grasslands to row crops in Hungary (Chrastina and Boltižiar, 2008), Romania (Schreiber, 2003), and the Czech Republic (Chrastina and Boltižiar, 2008; Havlíček et al., 2011). During the Socialist time, annual forest cover loss was high due to the clearing of forested area of no economic value (small isolated patches and shrubby vegetation) in Slovakia and the Czech Republic (Demek et al., 2008; Špulerová, 2008; Stránská, 2008). Political goals of increasing agricultural production caused agricultural expansion in the Czech Republic (Demek et al., 2008; Skokanová et al., 2009; Štych, 2007). There was also considerable regional variation related to agricultural expansion: in some mountain areas (e.g., parts of the Polish and Slovak Carpathians) agricultural land remained privately owned and agriculture did not expand (Kozak, 2010; Mojses and Petrovič, 2013), while some agricultural expansion occurred in Romania (where 80% of the population was already employed in agriculture at the time of collectivization), and in the Great Plain of Hungary (about 50% of the population) (Kligman and Verdery, 2011). Deforestation between 1945 and 1990 was, however, not always related to agricultural expansion. For example, tourism and industrial development led to forest cover loss in the Southern Romanian Carpathians (Huzui et al., 2012) and the Tatra Mountains (Gerard et al., 2010, 2006a, 2006b). Similarly, since 1990, selective logging for household needs, illegal harvesting, and large scale clear-cuts due to loopholes in the forest laws of some countries (Irland and Kremenetska, 2009; Kuemmerle et al., 2009a) caused forest losses (Grozavu et al., 2012; Mihai et al., 2007, 2006)

with particularly heavy illegal logging reported in Romania (Knorn et al., 2012b; Shandra et al., 2013), and Ukraine (Kuemmerle et al., 2009a).

On the other hand, agricultural abandonment and reforestation occurred mostly after WWII, with few local exceptions during earlier times (Patru-Stupariu, 2011). Since 1880, forest cover increased along the timberline throughout the study area (Shandra et al., 2013). During the 19th and early 20th century, marginal agricultural sites in the Polish mountains exhibited the most abandonment due to harsh environmental conditions (Ostafin, 2009), while agriculture expanded in more favorable areas with little terrain, in line with the forest transition theory (Lambin and Meyfroidt, 2010; Lambin et al., 2001). The agricultural decrease was related to a shift of agricultural activities to more productive lands, as well as to industrialization (Gerard et al., 2010, 2006a, 2006b). During the Socialist time period, the forced industrialization of the 1970s led to migration from rural areas to cities, causing farmland abandonment, for example, in Romania (Schreiber, 2003). In the same period, forests increased along inaccessible areas of the Iron Curtain in the Czech Republic (Skokanová and Eremiášová, 2012), and Slovakia (Kaliivoda et al., 2010).

After the collapse of socialism, the lack of agricultural subsidies, decreased profitability (Müller et al., 2013; Prishchepov et al., 2012), and the bankruptcy of most large agricultural enterprises (Petrovič and Hreško, 2010; Turnock, 2002; Zaušková et al., 2011) caused widespread abandonment followed by reforestation (Boltiziar and Chrastina, 2008; Havlíček et al., 2009; Zaušková et al., 2011). Increasing emigration to western Europe (Munteanu et al., 2008; Petrovič, 2006) resulted in decreasing employment in the agricultural sector, reducing pressure on land and allowing forest succession to take place (Kozak, 2003; Kozak et al., 2007; Smaliychuk, 2010). In the Ukraine, after 1990, abandonment occurred mostly on large

agricultural fields, while subsistence agriculture reemerged on marginal lands in the mountains (Baumann et al., 2011). Last but not least, nature conservation policies contributed to stabilize or increase forest cover after 1945, and especially since 1990, in parts of Slovakia, Hungary and Poland (Gerard et al., 2006b; Konkoly-Gyuró et al., 2011; Olah and Boltižiar, 2009), even though the effectiveness of protected areas in Romania is uncertain (Knorn et al., 2012b). In mountain areas, forest increase was also triggered by decreasing grazing pressure (Mihai et al., 2007; Tirla et al., 2012; Zaušková et al., 2011) and changing climate (Mihai et al., 2006; Shandra et al., 2013; Tirla et al., 2012). On the other hand, after the EU accession, nature conservation and agricultural policies alongside with awareness of the loss of valuable mountain grasslands, resulted in a shift from arable land to high-nature value meadows and from forest to pastures (Bezák and Halada, 2010; Cebecaurová and Cebecauer, 2008; Zaušková et al., 2011).

Most of our case studies reported interactions among the drivers of land change, with broader political decisions being often the underlying factors constraining economic and social conditions (Cebecaurová and Cebecauer, 2008; Janicki, 2004; Sitko and Troll, 2008). The same driver also often caused different land change patterns in different parts of the region: for example during the Socialist time period, national policies led to agricultural expansion on fertile soils in Hungary (Chrastina and Boltižiar, 2008), while forced industrialization as a national policy caused migration and abandonment of agriculture in areas of Romania (Schreiber, 2003). Relative to other drivers of change, the effects of culture on land use may only become apparent at long temporal scales. This means that culture may not have been an important driver at the temporal scale of some of our case studies, which reported change on the order of decades rather than centuries, and culture thus being less prevalent in our summaries than its overall importance

would suggest. Furthermore individual effects of drivers were difficult to isolate because of the interplay between social, economic and political elements that lead to local land-use decisions.

It was beyond the scope of our analysis to assess changes in land-use intensity, since most case studies did not map these explicitly. However, across the region, notable changes include agricultural intensification and shifts in forest management. Intensification was driven mostly by economic and technological development throughout the 19th century (Demek et al., 2008; Havlíček et al., 2011; Skokanová et al., 2009), when both crop rotation and industrial fertilizers were introduced. Similarly, Soviet agricultural policies led to intensification (Cebecaurová and Cebecauer, 2008; Mojses and Bezák, 2010; Skokanová et al., 2009) while nationalization of land caused increase in property sizes and the shift from small-scale farms to large state-owned agricultural operations (Boltižiar and Chrastina, 2006; Krivosudsky, 2011; Štych, 2007; Štych et al., 2012). These changes did not necessarily affect the land cover, but led to landscape homogenization (Krivosudsky, 2011; Mojses and Boltižiar, 2011; Špulerová, 2008). Conversely, changes in forest use affected forest patterns and fragmentation: non-native species were planted for timber production (Chrastina and Boltižiar, 2010; Nagy, 2008) and heavy logging and clearcuts occurred during Soviet times in Romania and Slovakia (Boltižiar and Chrastina, 2008; Grozavu et al., 2012; Niculita et al., 2008) due to increased demand for wood. Despite the documented overall forest cover increase after 2000 (0.89% mean annual change), extensive forest disturbances - which do not necessarily alter the land-cover type - occurred in Romania, Poland, Ukraine and the Czech Republic (Griffiths et al., 2014; Kuemmerle et al., 2009a).

Overall, our analysis provided a synthesis of land change patterns and processes during time periods with very different and rapidly changing political and economic conditions. The strength of our analyses lied in the multi-language data sources as well as in the fact that we

complemented this information with traditional library research, accessing a wide base of local knowledge. Furthermore, by synthesizing land changes in 102 case-studies, our analysis improves the understanding of overall trends, and acknowledges regional and temporal differences, rather than extrapolating findings of single micro-scale studies. We showed that rates of change differed markedly over the past 250 years: after the collapse of the Austro-Hungarian Empire agricultural land declined, while the collapse of the socialism accelerated agricultural abandonment and forest cover increase. We also showed that recent land change trends do follow long term land changes in terms of direction of changes but the magnitude of these processes differs substantially across periods, with high rates of change being captured since the collapse of socialism. We acknowledge that some case-studies were focused on capturing change based on unique conditions, such as depopulated areas of Poland (e.g., Maciejowski, 2001; Warcholik, 2005; Wolski, 2001) or flooded villages in Slovakia (Petrovič and Bezák, 2010) so that our analysis might describe the very peaks of observed processes. However, despite the abrupt changes in political and economic systems, which might disrupt gradual land transitions, the forest transition theory holds true in this region with the shift from decreasing to increasing forest cover occurring between the two World Wars for the most case studies.

Our results show that despite repeated shocks, which can alter the intensity of long-term, gradual changes, forest transition theory holds true in the Carpathian region. The change point from forest decrease to forest increase occurred between the two World Wars in most of the areas. Around the world, the forest transition occurred at different points in time, highly dependent on the socio-economic, institutional conditions and global marked dynamics. For example, western European countries like France and Britain experienced forest transition already in the second

half of the 18th century (Mather, 1998, 1992) as a function of their land use specialization and developing economies (Barbier et al., 2010; Lambin and Meyfroidt, 2010). Countries, such as India, China, Vietnam, Chile, El Salvador, on the other hand, experienced forest transition recently (second half of the 20th century), driven by governance, global market dynamics and displacement of land use abroad (Mather, 2007; Meyfroidt et al., 2010).

In the Carpathians, the agricultural change was mostly mirrored by forest cover and also involved other land-cover classes, for which data availability was limited. Agricultural expansion up to the early part of the 20th century, coincides broadly with the trends observed during the 19th century in other developed regions such as the US, Canada or USSR, where agriculture expanded mostly over grasslands and forests (Goldewijk, 2001). In developing tropical countries, agricultural expansion and intensification is a more recent process (Keys and McConnell, 2005). Countries like Bhutan, Brazil, Costa Rica or China experience large increases in agricultural covers only in the second half of the 20th century (Meyfroidt et al., 2010). In contrast, during the same time, many post-soviet countries face land abandonment since the collapse of socialism (Alcantara et al., 2012; Prishchepov et al., 2013), a similar trend to the one we observed in the Carpathians. Regional differences were notable, especially due to physical factors and several interacting driving forces, but institutional, policy and economic drivers were most influential in shaping both deforestation and agricultural expansion. Socio- demographic factors like rural population decline were the key drivers for land abandonment. Overall, we highlighted the value of longitudinal studies of land change to reveal the strong effects that repeated socio-economic and institutional changes have on land-use and land-cover.

References

- Alcantara, C., Kuemmerle, T., Prishchepov, A. V., Radeloff, V.C., 2012. Mapping abandoned agriculture with multi-temporal MODIS satellite data. *Remote Sens Environ* 124, 334–347.
- Barbier, E.B., Burgess, J.C., Grainger, A., 2010. The forest transition: Towards a more comprehensive theoretical framework. *Land use policy* 27, 98–107.
- Baumann, M., Kuemmerle, T., Elbakidze, M., Ozdogan, M., Radeloff, V.C., Keuler, N.S., Prishchepov, A. V., Kruhlov, I., Hostert, P., 2011. Patterns and drivers of post-socialist farmland abandonment in Western Ukraine. *Land use policy* 28, 552–562.
- Bezák, P., Halada, L., 2010. Sustainable Management Recommendations to Reduce the Loss of Agricultural Biodiversity in the Mountain Regions of NE Slovakia. *Mt Res Dev* 30, 194–204.
- Bičík, I., Jeleček, L., Štěpánek, V., 2001. Land-use changes and their social driving forces in Czechia in the 19th and 20th centuries. *Land use policy* 18, 65–73.
- Biró, M., Czúcz, B., Horváth, F., Révész, A., Csatári, B., Molnár, Z., 2012. Drivers of grassland loss in Hungary during the post-socialist transformation (1987–1999). *Landsc Ecol* 28, 789–803.
- Björnsen-Gurung, A., Bokwa, A., Chelmicki, W., Elbakidze, M., Hirschmugl, M., Hostert, P., Ibisch, P., Kozak, J., Kuemmerle, T., Matei, E., Ostapowicz, K., Pociask-Karteczka, J., Schmidt, L., van der Linden, S., Zebisch, M., 2009. Global Change Research in the Carpathian Mountain Region. *Mt Res Dev* 29, 282–288.
- Bolťižiar, M., Chrastina, P., 2006. Využitie krajiny SV okraja Bakoňského lesa v Maďarsku (in Slovak). *Geogr Rev* 2, 49–61.
- Bolťižiar, M., Chrastina, P., 2008. Zmeny využitia zeme nížinnej poľnohospodárskej krajiny na príklade obce Nové Sady (1782–2002), (in Slovak). *Geoinformation* 4, 16–35.
- Bürgi, M., Hersperger, A.M., Schneeberger, N., 2005. Driving forces of landscape change - current and new directions. *Landsc Ecol* 19, 857–868.
- Cebecaurová, M., Cebecauer, T., 2008. Spatio-temporal trends of landscape development in southwest part of Slovakia: analysis of major landscape change types. *Ekológia Int J Ecol Probl Biosph* 27, 212–228.
- Chrastina, P., Bolťižiar, M., 2008. Zmeny krajiny Slovenskej enklávy Veľký Bánhedeš, in: *Slovenčina v Menšinovom Prostredí. Štúdie Z Druhej Medzinárodnej Konferencie Výskumného Ústavu Slovákov v Maďarsku Békešská Čaba 17.-18.10.2007* (In Slovak). *Výskumný ústav Slovákov v Maďarsku, Békešská Čaba*, pp. 397–408.
- Chrastina, P., Bolťižiar, M., 2010. Senváclav: Krajina – človek – kultúra slovenskej enklávy vo Vyšegrádskejších vrchoch, in: *Viedermann, E. (Ed.), Studia Historica Nitriensia* (in Slovak). *Filozofická fakulta Univerzity Konštantína Filozofa v Nitre, Nitra*, pp. 53–86.

- Cioroianu, A., 2007. Pe umerii lui Marx. O introducere în istoria comunismului românesc (in Romanian). Curtea Veche. Bucuresti.
- Csaki, C., Jambor, S., 2009. The Diversity of Effects of EU Membership on Agriculture in New Member States. Policy Stud Rural Transition FAO 4, 44p.
- DeFries, R.S., Foley, J.A., Asner, G.P., 2004. Land-use choices : balancing human needs and ecosystem function In a nutshell : *Frontiers Ecol* 2, 249–257.
- Demek, J., Havlíček, M., Chrudina, Z., Mackovčín, P., 2008. Changes in land-use and the river network of the Graben Dyjsko-svratecký úval (Czech Republic) in the last 242 years. *J Landsc Ecol* 1, 22–51.
- Drgona, V., 2004. Assessment of the landscape use changes in the city of Nitra. *Ekológia Int J Ecol Probl Biosph* 23, 385–392.
- Feranec, J., O'ahel', J., 2009. Land cover/land use change research and mapping in Slovakia. *Geogr Slovaca* 26, 169–190.
- Foley, J.A., DeFries, R.S., Asner, G.P., Barford, C., Bonan, G., Carpenter, S.R., Chapin, F.S., Coe, M.T., Daily, G.C., Gibbs, H.K., Helkowski, J.H., Holloway, T., Howard, E.A., Kucharik, C.J., Monfreda, C., Patz, J.A., Prentice, I.C., Ramankutty, N., Snyder, P.K., 2005. Global consequences of land use. *Science* (80-) 309, 570–574.
- Foster, D.R., Swanson, F.J., Aber, J., Burke, I., Brokaw, N., Tilman, D., Knapp, A., 2003. The importance of land-use legacies to ecology and conservation. *Bioscience* 53, 77–88.
- Gehlbach, S., Malesky, E.J., n.d. The Grand Experiment That Wasn't? New Institutional Economics and the Postcommunist Experience, in: Galiani, S., Itai, S. (Eds.), *Institutions, Property Rights and Economic Growth: The Legacy of Douglass North*. Forthcoming.
- Geist, H.J., Lambin, E.F., 2002. Proximate Causes and Underlying Driving Forces of Tropical Deforestation. *Bioscience* 52, 143.
- Geist, H.J., Lambin, E.F., 2004. Dynamic causal patterns of desertification. *Bioscience* 54, 817–829.
- Gerard, F., Bugár, G., Gregor, M., Halada, L., Hazeu, G., Huiti, H., Köhler, R., Kolář, J., Luque, S., Mál'cher, C., Olschofsky, K., Petit, S., Pino, J., Smith, G., Thomson, A., Wachowicz, M., Bezák, P., Boltižiar, M., De Badts, E., Halabuk, A., Manchester, S., Mojses, M., Petrovič, F., Roda, F., Roscher, M., Tuominen, S., Ziese, H., 2006a. BIOPRESS Final report, sections 5 and 6: Jan 2003- Dec 2005, EC-FPV Contract.
- Gerard, F., Petit, S., Smith, G., Thomson, A., Brown, N., Manchester, S., Wadsworth, R., Bugár, G., Halada, L., Bezak, P., Boltižiar, M., De badts, E., Halabuk, A., Mojses, M., Petrovič, F., Gregor, M., Hazeu, G., Mucher, C. a., Wachowicz, M., Huitu, H., Tuominen, S., Kohler, R., Olschofsky, K., Ziese, H., Kolar, J., Sustera, J., Luque, S., Pino, J., Pons, X., Roda, F., Roscher, M., Feranec, J., 2010. Land cover change in Europe between 1950 and 2000 determined employing aerial photography. *Prog Phys Geogr* 34, 183–205.

- Gerard, F., Thomson, A., Wadsworth, R., Gregor, M., Luque, S., Sandra, L., Huitu, H., Köhler, R., Olschofsky, K., Hazeu, G., Mucher, S., Halada, L., Bugár, G., Pino, J., 2006b. Land cover change in Europe from the 1950ies to 2000. Aerial photo interpretation and derived statistics from 59 samples distributed across Europe. Institute for World Forestry, University of Hamburg, Hamburg.
- Goldewijk, K.K., 2001. Estimating global land use change over the past 300 years: The HYDE Database. *Global Biogeochem Cycles* 15, 417.
- Grekov, S., 2002. Do suchasnoi dynamiki silskogospodarskogo zemlekoristuvannya v Karpato-Podilskomu regioni (in Ukrainian). *Nauk zapysky Ternoilskogo derjavnogo Pedagog Univ Ser Geogr* 138, 154–161.
- Griffiths, P., Kuemmerle, T., Baumann, M., Radeloff, V.C., Abrudan, I. V., Lieskovský, J., Munteanu, C., Ostapowicz, K., Hostert, P., 2014. Forest disturbances, forest recover, and changes in forest types across the Carpathian ecoregion from 1985 to 2010 based on Landsat image composites. *Remote Sens Environ* 151, 72–88.
- Grozavu, A., Marginit, M., Niculita, C., 2012. The dynamics of land use in the middle sector of the Moldova river drainage basin (Eastern Carpathians, Romania), in: Boltižiar, M. (Ed.), *Conference Abstracts of the 2nd Forum Carpathicum. From Data to Knowledge, from Knowledge to Action*. Slovak Academy of Sciences, Bratislava, p. 214.
- Hansen, M.C., Stehman, S. V, Potapov, P. V, 2010. Quantification of global gross forest cover loss. *Proc Natl Acad Sci USA* 107, 8650–5.
- Havlíček, M., Borovec, R., Svoboda, J., 2009. Long-term changes in land use in the Litava River basin. *Acta Pruhoniana* 31–37.
- Havlíček, M., Krejčíková, B., Chrudina, Z., Borovec, R., Svoboda, J., 2011. Land use and streams changes in the Velička river basin, the Kyjovka upper river basin and the Svratka upper river basin. *Acta Pruhoniana* 99 5–17.
- Hostert, P., Kuemmerle, T., Prishchepov, A. V., Sieber, A., Lambin, E.F., Radeloff, V.C., 2011. Rapid land use change after socio-economic disturbances: the collapse of the Soviet Union versus Chernobyl. *Environ Res Lett* 6, 8.
- Huzui, A.E., Călin, I., Pătru-Stupariu, I., 2012. Spatial pattern analyses of landscape using multi-temporal data sources. *Procedia Environ Sci* 2011 Int Conf Environ Identity 14, 98–110.
- Irland, L.C., Kremenetska, E., 2009. Practical Economics of Forest Ecosystem Management: The Case of the Ukrainian Carpathians, in: Soloviy, I.P., Keeton, W.S. (Eds.), *Ecological Economics and Sustainable Forest Management: Developing a Transdisciplinary Approach for the Carpathian Mountains*. Ukrainian National Forestry University Press, Lviv, p. 432.
- Janicki, R., 2004. Specific features of the landscape succession processes in Przemyskie Foothills, in: Kirchner, K., Wojtanowicz, J. (Eds.), *Cultural Landscapes. Regiograph*, Brno, p. 377.

- Kaim, D., 2009. Zmiany pokrycia terenu na pograniczu polsko-słowackim na przykładzie Małych Pienin (in Polish). *Przegląd Geogr* 81, 93–106.
- Kalivoda, H., Petrovič, F., Kalivodová, E., Kürthy, A., 2010. Influence of the landscape structure on the butterfly (Lepidoptera, Hesperioidea and Papilionoidea) and bird (Aves) taxocoenoses in Veľké Leváre (SW Slovakia). *Ekológia Int J Ecol Probl Biosph* 29, 337–359.
- Keys, E., McConnell, W.J., 2005. Global change and the intensification of agriculture in the tropics. *Glob Environ Chang* 15, 320–337.
- Kligman, G., Verdery, K., 2011. Peasants under siege: the collectivization of Romanian agriculture, 1949-1962. Princeton University Press.
- Knorn, J., Kuemmerle, T., Radeloff, V.C., Szabo, A., Mindrescu, M., Keeton, W.S., Abrudan, I. V., Griffiths, P., Gancz, V., Hostert, P., 2012. Forest restitution and protected area effectiveness in post-socialist Romania. *Biol Conserv* 146, 204–212.
- Knorn, J., Rabe, A., Radeloff, V.C., Kuemmerle, T., Kozak, J., Hostert, P., 2009. Land cover mapping of large areas using chain classification of neighboring Landsat satellite images. *Remote Sens Environ* 113, 957–964.
- Konkoly-Gyuró, E., Nagy, D., Balázs, P., Király, G., 2011. Assessment of land cover change in western Hungarian landscapes, in: Balázs, P., Konkoly-Gyuró, É. (Eds.), *TransEcoNet Workshop on Landscape History Proceedings*. University of West Hungary Press, pp. 75–89.
- Kozak, J., 2003. Forest Cover Change in the Western Carpathians in the Past 180 Years. *Mt Res Dev* 23, 369–375.
- Kozak, J., 2010. Forest Cover Changes and Their Drivers in the Polish Carpathian Mountains Since 1800, in: Nagendra, H., Southworth, J. (Eds.), *Reforestation Landscapes Linking Pattern and Process*. Landscape Series 10. Springer, pp. 253–273.
- Kozak, J., Estreguil, C., Troll, M., 2007. Forest cover changes in the northern Carpathians in the 20th century: a slow transition. *J Land Use Sci* 127–146.
- Krivosudsky, R., 2011. Land use development of south-east slopes of Malé Karpaty mts, in: Dobrovodská, M., Špulerová, J., Štefunková, D. (Eds.), *Research and Management of Historical Agricultural Landscape*. Institute of Landscape Ecology, Slovak Academy of Sciences, pp. 39–49.
- Kuemmerle, T., Chaskovskyy, O., Knorn, J., Radeloff, V.C., Kruhlov, I., Keeton, W.S., Hostert, P., 2009a. Forest cover change and illegal logging in the Ukrainian Carpathians in the transition period from 1988 to 2007. *Remote Sens Environ* 113, 1194–1207.
- Kuemmerle, T., Hostert, P., Radeloff, V.C., Perzanowski, K., Kruhlov, I., 2007. Post-socialist forest disturbance in the Carpathian border region of Poland, Slovakia, and Ukraine. *Ecol Appl* 17, 1279–95.

- Kuemmerle, T., Hostert, P., Radeloff, V.C., van der Linden, S., Perzanowski, K., Kruhlov, I., 2008. Cross-border Comparison of Post-socialist Farmland Abandonment in the Carpathians. *Ecosystems* 11, 614–628.
- Kuemmerle, T., Müller, D., Griffiths, P., Rusu, M., 2009b. Land use change in Southern Romania after the collapse of socialism. *Reg Environ Chang* 9, 1–12.
- Kuemmerle, T., Oloffson, P., Chaskovskyy, O., Baumann, M., Ostapowicz, K., Woodcock, C.E., Houghton, R.A., Hostert, P., Keeton, W.S., Radeloff, V.C., 2011. Post-Soviet farmland abandonment, forest recovery, and carbon sequestration in western Ukraine. *Glob Chang Biol* 17, 1335–1349.
- Lambin, E.F., Meyfroidt, P., 2010. Land use transitions: Socio-ecological feedback versus socio-economic change. *Land use policy* 27, 108–118.
- Lambin, E.F., Turner, B.L., Geist, H.J., Agbola, S.B., Angelsen, A., Bruce, J.W., Coomes, O.T., Dirzo, R., Fischer, G., Folke, C., George, P.S., Homewood, K., Imbernon, J., Leemans, R., Li, X., Moran, E.F., Mortimore, M., Ramakrishnan, P.S., Richards, J.F., Skånes, H., Steffen, W., Stone, G.D., Svedin, U., Veldkamp, T.A., Vogel, C., Xu, J., 2001. The causes of land-use and land-cover change: moving beyond the myths. *Glob Environ Chang* 11, 261–269.
- Maciejowski, W., 2001. Zmiany użytkowania ziemi i ich wpływ na funkcjonowanie środowiska przyrodniczego w zlewni górnej Wilszni (Beskid Niski) w okresie 1920-2000 (in Polish), in: German, K., Balon, J. (Eds.), *Przemiany Środowiska Przyrodniczego Polski a Jego Funkcjonowanie*. Instytut Geografii i Gospodarki Przestrzennej Uniwersytetu Jagiellońskiego, Kraków.
- Mather, A.S., 1992. The forest transition. *Area* 24, 367–379.
- Mather, A.S., 1998. The forest transition : a theoretical basis. *Area* 30, 117–124.
- Mather, A.S., 2007. Recent Asian forest transitions in relation to foresttransition theory. *Int For Rev* 9, 491–502.
- McConnell, W.J., Keys, E., 2005. Meta-analysis of agricultural change, in: Moran, E.F., Ostrom, E. (Eds.), *Seeing the Forest and the Trees. Human-Environment Interactions in Forest Ecosystems*. MIT Press, Cambridge, pp. 325–354.
- Meyfroidt, P., Lambin, E.F., 2011. Global Forest Transition: Prospects for an End to Deforestation. *Annu Rev Environ Resour* 36, 343–371.
- Meyfroidt, P., Rudel, T.K., Lambin, E.F., 2010. Forest transitions, trade, and the global displacement of land use. *Proc Natl Acad Sci USA* 107, 20917–22.
- Mihai, B., Savulescu, I., Sandric, I., 2007. Change Detection Analysis (1986-2002) of Vegetation Cover in Romania. *Mt Res Dev* 27, 250–258.

- Mihai, B., Savulescu, I., Sandric, I., Oprea, R., 2006. Application de la detection des changements a l'etude de la dynamique de la vegetation des Monts de Bucegi Bucegi (Carpathes Méridionales, Roumanie)(in French). *Télédetection* 6, 215–231.
- Mojšes, M., Bezák, P., 2010. Parížske Močiare, in: Bezák, P., Izakovičová, Z., Miklós, L. (Eds.), *Reprezentatívne Typy Krajiny Slovenska* (in Slovak). Ústav krajiny ekológie Slovenskej Akadémie Vied, Bratislava, p. 180.
- Mojšes, M., Boltížiar, M., 2011. Using spatial Metrics for assessment of the landscape structure changes of the Beša dry polder. *Tájökológiai Lapok* 9, 415–428.
- Mojšes, M., Petrovič, F., 2013. Land use changes of historical structures in the agricultural landscape at the local level - Hriňova case study. *Ekológia Int J Ecol Probl Biosph* 32, 1–12.
- Müller, D., Leitão, P.J., Sikor, T., 2013. Comparing the determinants of cropland abandonment in Albania and Romania using boosted regression trees. *Agric Syst* 117, 66–77.
- Munteanu, C., Geitner, C., Scharr, K., 2008. Consequences of historical and modern land use on cultural landscapes and biodiversity of the Maramureş Mountains, in: Gaceu, L. (Ed.), *Journal of EcoAgriTourism, Proceeding of the International Conference BIOATLAS 2008*. Transilvania University of Brasov, Romania, Brasov, pp. 91–98.
- Nagy, D., 2008. A történeti felszínborítás térképezése a Tisza-völgyben., in: Flachner, Z., Kovács, A., Kelemen, É. (Eds.), *A Történeti Felszínborítás Térképezése a Tiszavölgyben* (in Hungarian). Nagykorú-Eger-Budapest, Budapest, pp. 40–58.
- Niculita, C., Lupascu, A., Minea, I., 2008. Evolutia suprafetelor forestiere in ocoalele silvice Ciurea si Podu Iloaiei intre anii 1894-2000 (in Romanian). *Lucr Stiint* 51, 195–201.
- Ojima, D.S., Galvin, K.A., Turner, B.L., 1994. The global impact of land-use change. *Bioscience* 44, 300–304.
- Olah, B., Boltížiar, M., 2009. Land use changes within the Slovak biosphere reserves's zones. *Ekológia Int J Ecol Probl Biosph* 28, 127–142.
- Ostafin, K., 2009. Zmiany granicy rolno-leśnej w środkowej część Beskidu Średniego od połowy XIX wieku do 2005 roku (in Polish). roku, Wydaw UJ. Wydawnictwo Uniwersytetu Jagiellońskiego Kraków.
- Pandey, D., 1995. Forest resources assessment 1990: tropical forest plantation resources. FAO For Pap FAO Rome 128, 90p.
- Paracchini, M., Petersen, J., Hoogeveen, Y., 2008. High Nature Value Farmland in Europe: An estimate of the distribution patterns on the basis of land cover and biodiversity data., agrienv.jrc.ec.europa.eu/ Luxemburg.
- Patru-Stupariu, I., 2011. Peisaj si gestiunea durabila a teritoriului. Aplicatii la culoarul transcarpatic Bran-Rucar-Dragoslavele (in Romanian), 2nd ed. University Press Bucharest, Bucharest.

- Petrovič, F., 2006. The changes of the landscape with dispersed settlement. *Ekológia Int J Ecol Probl Biosph* 25, 201–211.
- Petrovič, F., Bezák, P., 2010. Novobanská štálová oblasť, in: Bezák, P., Izakovičová, Z., Miklós, L., Al., E. (Eds.), *Reprezentatívne Typy Krajiny Slovenska* (in Slovak). Ústav krajinnéj ekológie Slovenskej Akadémie Vied, Bratislava, p. 180.
- Petrovič, F., Hreško, J., 2010. Vplyv výstavby vodnej nádrže Domaša na súčasnú krajinnú štruktúru (in Slovak). *Životné Prostr* 44, 182–186.
- Poteete, A.R., Ostrom, E., 2008. Fifteen Years of Empirical Research on Collective Action in Natural Resource Management: Struggling to Build Large-N Databases Based on Qualitative Research. *World Dev* 36, 176–195.
- Prishchepov, A. V., Müller, D., Dubinin, M., Baumann, M., Radeloff, V.C., 2013. Determinants of agricultural land abandonment in post-Soviet European Russia. *Land use policy* 30, 873–884.
- Prishchepov, A. V., Radeloff, V.C., Baumann, M., Kuemmerle, T., Müller, D., 2012. Effects of institutional changes on land use: agricultural land abandonment during the transition from state-command to market-driven economies in post-Soviet Eastern Europe. *Environ Res Lett* 7, 024021.
- Puyravaud, J.P., 2003. Standardizing the calculation of the annual rate of deforestation. *For Ecol Manage* 177, 593–596.
- Rudel, T.K., 2008. Meta-analyses of case studies: A method for studying regional and global environmental change. *Glob Environ Chang* 18, 18–25.
- Salvatori, V., Okarma, H., Ionescu, O., Dovhanych, Y., Find, S., 2002. Hunting legislation in the Carpathian Mountains : implications for the conservation and management of large carnivores. *Wildlife Biol* 8, 3–10.
- Schiller, H., Miklós, D., Sass, J., 2010. The Danube River and its Basin Physical Characteristics, Water Regime and Water Balance, in: Brilly, M. (Ed.), *Hydrological Processes of the Danube River Basin: Perspectives from the Danubian Countries*. Springer, pp. 25–77.
- Schreiber, W., 2003. Analiza peisajelor geografice din partea de vest a Câmpiei Transilvaniei (in Romanian). *Presa Univ. Clujeană, Cluj Napoca*.
- Seto, K.C., Fragkias, M., Güneralp, B., Reilly, M.K., 2011. A meta-analysis of global urban land expansion. *PLoS One* 6, e23777.
- Shandra, O., Weisberg, P., Martazinova, V., 2013. Influences of Climate and Land Use History on Forest and Timberline Dynamics in the Carpathian Mountains During the Twentieth Century, in: Kozak, J., Ostapowicz, K., Bytnerowicz, A., Wyżga, B. (Eds.), *The Carpathians: Integrating Nature and Society Towards Sustainability*. Springer, pp. 209–223.

- Sitko, I., Troll, M., 2008. Timberline Changes in Relation to Summer Farming in the Western Chornohora (Ukrainian Carpathians). *Mt Res Dev* 28, 263–271.
- Skokanová, H., Eremiášová, R., 2012. Changes in the secondary landscape structure and the connection to ecological stability: the cases of two model areas in the Czech Republic. *Ekológia Int J Ecol Probl Biosph* 31, 33–45.
- Skokanová, H., Havlíček, M., Borovec, R., Demek, J., Eremiášová, R., Chrudina, Z., Mackovčín, P., Rysková, R., Slavík, P., Stránská, T., Svoboda, J., 2012. Development of land use and main land use change processes in the period 1836–2006. *J Maps* 8, 88–96.
- Skokanová, H., Stránská, T., Havlíček, M., Borovec, R., Eremiášová, R., Rysková, R., Svoboda, J., 2009. Land use dynamics of the South Moravian region during last 170 years. *Geoscape* 4, 58–65.
- Smaliychuk, A., 2010. Doslidjennya suchasnoi dynamyky geoekosistem nyzkogir'ya Karpat u mejah Lvivskoi oblasti z vykoristannyam GIS tehnologiy (in Ukrainian). *Fiz Geogr ta Geomorfol* 3, 83–91.
- Špulerová, J., 2008. Land use changes in the Veselovianka river catchment in the Horná Orava region. *Ekológia Int J Ecol Probl Biosph* 27, 326–337.
- Stránská, T., 2008. Dynamika vývoje krajinné struktury okolí Dolních Kounic (in Czech), in: Herber, V. (Ed.), *Fyzickogeografický Sborník 6*. Masarykova univerzita, Brno, pp. 263–268.
- Štych, P., 2007. Regional differentiation of long-term land-use changes in the Czech Republic. Charles University in Prague. PhD Thesis.
- Štych, P., Bičík, I., Stonáček, V., Bláha, J., 2012. Change of Land Use Patterns 1827-2005, in: Bičík, I., Himiyama, Y., Feranec, J., Štych, P. (Eds.), *Land Use/Cover Changes in Selected Regions in the World. Volume VII. IGU-LUCC*, Prague, pp. 37–41.
- Tirla, L., Matei, E., Vdulie, I., Cuculici, R., 2012. Analysis of the vegetation cover in the eastern Căpățâni Mountains (Southern Carpathians, Romania) between 1973 – 2005 using remote sensing and GIS data, in: Boltižiar, M. (Ed.), *Conference Abstracts of the 2nd Forum Carpaticum. From Data to Knowledge, from Knowledge to Action*. Slovak Academy of Sciences, Bratislava, p. 214.
- Turnock, D., 2002. Ecoregion-based conservation in the Carpathians and the land-use implications. *Land use policy* 19, 47–63.
- Vepryk, N., 2001. Istoryko-landshaftoznavchyy analiz lisogospodarskogo pryrodokorystuvannya (na prykladi Pivnichnoi Bukoviny u kinci 18 – na pochatku 20 stolittya) (in Ukrainian). *Nauk Visnyk Chernivetskogo Univ Ser Geogr* 120, 156–171.
- Vepryk, N., 2002. Agrolandshafty i zemlerobstvo Pivnichnoi Bukoviny u kinci 18-na pochatky 20 stolittya (in Ukrainian). *Nauk Visnyk Chernivetskogo Univ Ser Geogr* 138, 135–149.

- Verburg, P.H., Berkel, D.B., Doorn, A.M., Eupen, M., Heiligenberg, H. a. R.M., 2009. Trajectories of land use change in Europe: a model-based exploration of rural futures. *Landsc Ecol* 25, 217–232.
- Warcholik, W., 2005. Rejestracja różnic w przebiegu granicy rolno-leśnej w Beskidzie Niskim na obszarze Polski i Słowacji (1933-1975) z wykorzystaniem GIS (in Polish). *Probl Zagospodarowania Ziemi Górskich* 51, 59–69.
- Wolski, J., 2001. Kierunki zmian krajobrazu okolic bieszczadzkiej wsi Caryńskie (in Polish). *Pr Geogr IGiPZ PAN* 179, 149–167.
- Woś, B., 2005. Zmiany pokrycia terenu w wybranych gminach Beskidów w drugiej połowie XX w. na podstawie analizy zdjęć lotniczych (in Polish). *Teledetekcja Środowiska* 35, 1–114.
- Yeo, I.-Y., Huang, C., 2013. Revisiting the forest transition theory with historical records and geospatial data: A case study from Mississippi (USA). *Land use policy* 32, 1–13.
- Zaušková, Ľ., Feješ, J., Kysucká, K., 2011. Zmeny a pustnutie poľnohospodárskej krajiny v k.ú. Podhorie (in Slovak). *Univerzita Mateja Bela, Banská Bystrica*.

Tables and figures



Table 1.1 Time periods, their duration, the expected land changes and the number of studies that report land change for the specific period. The first number (*) indicates that the annual rate of change has been calculated for only one period. The second number (**) indicates that the case-study spans at least two time periods, and the annual rate of change is calculated based only on land cover at the beginning and end of the considered time span.

Time period	Duration	Expected land change		Number of case studies	
		Forest	Agricultural	Forest (n)	Agriculture (n)
K.u.K Monarchy	1750-1914	-	+	31* / 51**	24* / 43**
Interwar	1914-1945	-	+	29* / 72**	28* / 46**
Socialist	1945-1990	+	+	46* / 96**	37* / 63**
Transition	1990-2000	+	-	46* / 84**	42* / 68**
EU accession	2000-2012	+	+	37* / 60**	26* / 47**

Table 1.2 Comparison of expected and observed land changes for each time period and the mean annual rates of change, calculated for all the case studies for which change rates were not spanning more than one period (marked * in Table 1.1). For these calculations, only studies that report annual change for single periods were considered.

Time period	Expected land changes		Mean annual rate of change		Observed land changes	
	Forest	Agriculture	Forest	Agriculture	Forest	Agriculture
K.u.K Monarchy	-	+	0.08%	0.12%	0	+
Interwar	-	+	0.35%	-1.28%	+	-
Socialist	+	+	0.33%	-0.54%	+	-
Transition	+	-	1.07%	-1.61%	+	-
EU accession	+	+	0.89%	-1.20%	+	-

Figure 1.1 a) Study area, including spatial extent of case studies (grey) and centroids for 102 case study locations (triangles). Country codes: AT: Austria, HU: Hungary, PL: Poland, CZ: Czech Republic, SK: Slovakia, UA: Ukraine, RO: Romania, MD: Moldova, HR: Serbia, SI: Slovenia. b) Spatio-temporal distribution of 102 study locations (one location may include multiple case-studies).








Figure 1.3 Proportion of studies reporting decreasing ($<-0.1\%$ annually), stable (-0.1% to 0.1% annually) and increasing ($>0.1\%$ annually) cover for each time period for a) forest and c) agriculture and distribution of annual rates of change per time period for b) forest and d) agricultural cover. Country codes: CZ: Czech Republic, HU: Hungary, PL: Poland, RO: Romania, SK: Slovakia, UA: Ukraine.

Forest change



Agriculture change





Figure 1.4 Spatial and temporal distribution of forest change case studies. Annual rates of change are mapped for each case study and time period. Studies are represented by centroids. The size of the symbols indicate the amount of change, the colors indicate the direction of change (increase/stability/decrease). Shaded colors indicate that annual rates are calculated for more than one time period. Country codes: CZ: Czech Republic, HU: Hungary, PL: Poland, RO: Romania, SK: Slovakia, UA: Ukraine.




Figure 1.5 Spatial and temporal distribution of agricultural change case studies. Annual rates of change are mapped for each case study and time period. Studies are represented by centroids. The size of the symbols indicate the amount of change, the colors indicate the direction of change (increase/stability/decrease). Shaded colors indicate that annual rates are calculated for more than one time period. Country codes: CZ: Czech Republic, HU: Hungary, PL: Poland, RO: Romania, SK: Slovakia, UA: Ukraine.





Figure 1.6 Main classes of land change drivers and the relative importance of drives for each land change process in the study area. The proportions are calculated based on the number of times a driver was deemed as important in influencing change.



Appendix

Appendix 1.1 List of publications included in the meta-analysis

- | | |
|----------------------------------|---------------------------------|
| Boltižiar and Chrastina (2006) | Mojšes and Boltižiar (2011) |
| Boltižiar and Chrastina (2008) | Mojšes and Petrovič (2013) |
| Bugár et al. (2010) | Monastyrskiy (2010) |
| Cebecaurová and Cebecauer (2008) | Moyzeová and Izakovičová (2010) |
| Chrastina and Boltižiar (2008) | Muchová and Petrovič (2010) |
| Chrastina and Boltižiar (2010) | Nagy (2008) |
| Dec et al. (2009) | Niculita et al. (2008) |
| Demek et al. (2008) | Ostafin (2009) |
| Drgona (2004) | Ostapowicz & Ostafin (in prep) |
| Gerard et al. (2006a) | Ostapowicz and Kozak (2011) |
| Gerard et al. (2006b) | Patru-Stupariu (2011) |
| Grekov (2002) | Patru-Stupariu et al. (2011) |
| Grozavu et al. (2012) | Petrovič (2006) |
| Havlíček and Borovec (2008) | Petrovič and Bezák (2010) |
| Havlíček et al. (2009) | Petrovič and Hreško (2010) |
| Havlíček et al. (2011) | Petrovič and Muchová (2008) |
| Hurbánek and Pazúr (2007) | Pietrzak (2002) |
| Huzui et al. (2012) | Reiser (2006) |
| Jančovič et al. (2010) | Schreiber (2003) |
| Janicki (2004) | Shandra et al. (2013) |
| Kaim (2009) | Skokanová et al. (2009) |
| Kalivoda et al. (2010) | Špulerová (2008) |
| Konkoly-Gyuró et al. (2011) | Štefunková and Petrovič (2011) |
| Kozak (2003) | Stránská (2008) |
| Kozak et al. (2004) | Štych (2007) |
| Kozak et al. (2007) | Štych et al. (2012) |
| Kozak (unpublished) | Vepryk (2000) |
| Krivosudsky (2011) | Vepryk (2001) |
| Labuda and Pavličková (2006) | Vepryk (2002) |
| Maciejowski (2001) | Warcholik (2005) |
| Mihai et al. (2006) | Wolski (2001) |
| Mihai et al. (2007) | Woś (2005) |
| Mojšes and Bezák (2010) | Zaušková et al. (2011) |

Appendix 1.2 List of publications included in the meta-analysis

Paper not included	Criteria not met
Badea, (2011)	1
Bičík et al., (2001)	1
Bochko, (2010)	1
Boucníková and Kučera, (2005)	1,2
Feranec and Oťahel', (2008)	1,3
Feranec et al., (2000)	1,3
Feurdean, (2010)	1,3
Kilianová et al., (2008)	1
Konkoly-Gyuró et al., (2007)	3
Korjyk, (2001)	1
Kozak, (1994)	3
Lavruk, (2011)	1
Malahova, (2009)	1
Miklovoda and Gazuda, (2010)	1
Mulková et al., (2012)	1
Perovych et al., (2011)	1
Sitko and Troll, (2008)	1
Slivka and Savjuk, (2011)	1,3
Tirla et al., (2012)	3

References

- Badea, R., 2011. The Dynamics of Forest Areas in the Vâlsan Basin. *Forum Geogr.* X, 372–378.
- Bičík, I., Jeleček, L., Štěpánek, V., 2001. Land-use changes and their social driving forces in Czechia in the 19th and 20th centuries. *L. Use Policy* 18, 65–73.
- Bochko, O., 2010. Analiz zemelnoi ploschi Zakarpatskoi oblasti v suchasnyh umovah in Ukrainian). *Nauk. Visnyk Mukachivskogo Derjavnogo Univ.* 9, 146–152.
- Boucníková, E., Kučera, T., 2005. How natural and cultural aspects influence land cover changes in the Czech Republic? *Ekológia. Int. J. Ecol. Probl. Biosph.* 24, 69–82.
- Feranec, J., Oťahel', J., 2008. Land cover changes in Slovakia in the period 1970–2000. *Geogr. časopis* 60, 113–128.
- Feranec, J., Šúri, M., Oťahel', J., Cebecauer, T., Kolář, J., Soukup, T., Zdeňková, D., Waszmuth, J., Vâjdea, V., Vîjdea, A.M., Nitica, C., 2000. Inventory of major landscape changes in the Czech Republic, Hungary, Romania and Slovak Republic 1970s – 1990s. *Int. J. Appl. Earth Obs. Geoinf.* 2, 129–139.
- Feurdean, A.N., 2010. Forest conservation in a changing world: natural or cultural? Example from the Western Carpathians forests, Romania. *Stud. Univ. Babes-Bolyai, Geol.* 55, 45–48.
- Kilianová, H., Pechanec, V., Zapletalová, Z., 2008. Land use changes of agriculture landscape in Trkmanka catchment, in: Špulerová, J., Hrnčiarová, T. Eds.), *Biodiversity of Agricultural Landscape*. Institute of Landscape Ecology SAS, Bratislava, pp. 199–205.
- Konkoly-Gyuró, E., Jombach, S., Tatai, Z., 2007. Indicators for landscape identity : a novel social sustainability impact issue Basic source : Corine Land Cover Database. *Sens. Newsl.* 5 8–10.
- Korjyk, V., 2001. Lisystist Bukoviny yak funktsia antropogenizacii dovkilliya in Ukrainian). *Nauk. Visnyk Chernivetskogo Univ. Ser. Geogr.* 104, 36–60.
- Kozak, I., 1994. Ekologichni osoblyvosti typovyh biogeocenotichnyh kompleksiv (in Ukrainian), in: *Antropogenni Zminy Biogeocenotichnogo Pokryvu v Karpatskomu Regioni*. Golubets, Kyiv, p. 166.
- Lavruk, M., 2011. Geografia poloninskigi gospodarstva Gutsulshchiny na pochatky XXI stolittya in Ukrainian). *Visnyk Lviv. Univ. ser. Geogr.* 39, 218–230.

- Malahova, C., 2009. Analiz I otsinka suchasnogo stanu vedennya derjavnogo zeelnogo kadastru naselenyh punktiv na prykladi Lvivskoi oblasti in Ukrainian). Geod. Kartogr. I Aerofotozнимannya 72, 128–134.
- Miklovoda, V., Gazuda, M., 2010. Agrarian land use in regions of the Ukrainian Carpathians. Econ. Ind. 49, 163–171.
- Mulková, M., Popelka, P., Popelková, R., 2012. Landscape changes in the central part of the Karviná region from the first half of the 19 th century to the beginning of the 21st century. Ekológia. Int. J. Ecol. Probl. Biosph. 31, 75–91.
- Perovych, D., Perovych, L., Ludchak, O., 2011. Analiz vykoristannya zemelnyh resursiv Ivano-Frankivschyny – in Ukrainian. Suchasni dosyagnennya Geod. Nauk. ta vyrobnitstva 1, 214–217.
- Sitko, I., Troll, M., 2008. Timberline Changes in Relation to Summer Farming in the Western Chornohora Ukrainian Carpathians). Mt. Res. Dev. 28, 263–271.
- Slivka, R., Savjuk, M., 2011. Zminy u strukturi zemlekoristuvannya na teritorii Boikivschyny u XX stolitti in Ukrainian). Visnyk Priкарпатського Національного Університету. Серія Біол. 25, 31–34.
- Tirla, L., Matei, E., Vdulie, I., Cuculici, R., 2012. Analysis of the vegetation cover in the eastern Căpățâni Mountains Southern Carpathians, Romania) between 1973 – 2005 using remote sensing and GIS data, in: Boltižiar, M. Ed.), Conference Abstracts of the 2nd Forum Carpaticum. From Data to Knowledge, from Knowledge to Action. Slovak Academy of Sciences, Bratislava, p. 214.

Chapter 2. Legacies of 19th century land use shape contemporary forest cover

Introduction

Land use and land cover change are major components of global change, causing daunting sustainability challenges (Foley et al., 2005; Lambin and Geist, 2006; Sarukhán and Whyte, 2005). The effects of past land use (hereafter ‘land use legacies’) on the structure and functioning of current land system can be long-lasting. Legacies manifest themselves in all parts of ecosystems (Foster et al., 2003; Wallin et al., 1994) and can persist for decades (Wallin et al., 1994) or even centuries (Boucher et al., 2013; Thompson et al., 2013). The ecological effects of past land uses on current ecosystem structure are fairly well understood (Boucher et al., 2013; Foster et al., 2003; Rhemtulla and Mladenoff, 2007; Thompson et al., 2013) and path dependency has been conceptually acknowledged in land change science as an uncertainty factor (Brown et al., 2005; Lambin and Geist, 2006; National Research Council, 1998; Verburg et al., 2004), but empirical evidence on how much land use legacies affect contemporary land use change and land management is still scarce.

Past land uses can affect all parts of ecosystems (Foster et al., 2003; Wallin et al., 1994). For example, soil composition and nutrient content that were altered in the Eastern US during European settlement, are affecting plant abundances today (Thompson et al., 2013). The vegetation composition of historically ploughed areas has fewer shrubs and a distinct understory vegetation compared to continuously forested areas (Eberhardt et al., 2011; Motzkin and Foster, 2002). Similarly, prior farming in sagebrush ecosystems causes lower shrubs and forb cover today (Foster et al., 2003; Morris et al., 2011) while the high proportion of shrubby vegetation in dry areas, such as Chaco, New Mexico is due to overharvesting by the Anasazi (800 BC) as well as overgrazing and high stocking densities in the 1800s (Brown and Archer, 1989; Foster et al.,

2003; Fredrickson et al., 1998; Gibbens et al., 2005; Swetnam et al., 1999). Past land use decisions affecting possibilities of future change are probably best exemplified by urban area expansion, where path dependence constrains the possibility to revert an urban area to agricultural land (Lambin and Geist, 2006). Although land use legacies are widely acknowledged, the magnitude of their effect on contemporary land use dynamics at broad spatial and temporal scales is rarely quantified. The increasing number of studies and datasets capturing long term land use and land cover change (Başnou et al., 2013; Gerard et al., 2010) offers exciting new opportunities for the quantification of the legacies that past land uses exert on contemporary land change processes.

Forests are particularly likely to exhibit land use legacies, because they are persistent elements in landscapes due to the long lifespan of trees. Land use legacies can affect both forest structure and management decisions. For example, forests that were farmed during Roman times in Western Europe have a different seed bank than those that were always forested, including higher abundance of species that colonize abandoned land, and fewer seeds of poor dispersers (Dupouey et al., 2002a; Jan Plue et al., 2009). Historic land use leads to the occurrence of fruit tree species in oak forest systems (Plieninger et al., 2010a) and affects both forest structure and composition including basal area, tree density, and woody plant richness (Plieninger et al., 2010a; Rhemtulla et al., 2009). Even in cases in which forest composition is similar to that of historic forests, for example after agricultural abandonment in the Northeastern US, the relative importance of tree species is different and depends on the historic use (Thompson et al., 2013). Similarly, Mayan overexploitation of forests affects forest structure until today, due to changes in micro-topography, soil moisture, nutrient content and the location of ancient settlements (Foster et al.,

2003). However, while past land uses clearly affect current forest patterns, their impact is difficult to predict, especially if land management is constrained by such legacies.

Contemporary forest management may be severely constrained by historic uses and prior management practices, and land use legacies can play a defining role of the pathways of future forest change. Harvesting regimes can exert substantial legacy-effects because they establish an age-structure that can persist for several rotation cycles, even when management changes subsequently (Wallin et al., 1994). Similarly, rates of forest disturbance from either harvest or fires influence forest types in following decades, leading to less coniferous cover in the Russian Far East (Cushman and Wallin, 2000). Historic housing density, reforestation and fire suppression since the early 20th century affect forest management at landscape level at the end of the 20th century in the Midwestern US (Radeloff et al., 2001). In sum, both forest composition and structure are closely related to historic land uses, reaching back from decades to centuries.

However, while the long-term persistence of legacies in ecosystem structure and composition is relatively well understood in ecology, and path dependency has been an established concept in land change science (Lambin and Geist, 2006; National Research Council, 1998) the role of past land uses in modulating contemporary forest disturbance patterns has not been well quantified. Anecdotal evidence suggests that recent vegetation changes, such as shrub encroachment on overexploited agricultural land (Cramer et al., 2008; Foster et al., 2003) or reforestation on historically cleared pastures (Bezák and Mitchley, 2014; Sitko and Troll, 2008) are a consequence of past land uses. Moreover, past management affects ecosystem health and the susceptibility to change (Main-Knorn et al., 2009). Land use legacies may affect the pace and the timing of forest disturbance, making the consideration of land use history important when predicting future forest changes. However, the extent to which contemporary forest disturbance

is determined by land use legacies remains unclear, especially in comparison to other major drivers of land change such as environmental or socio-economic factors. In other words, it remains unknown how much forest disturbance is modulated by historic land use, and how much by other factors (Amacher et al., 2003; Beach et al., 2005; Geist and Lambin, 2001).

Our goal was to analyze the effects of land use legacies on contemporary forest disturbance in the Carpathian region, by assessing (1) the magnitude of contemporary forest disturbance, (2) the relation to spatial determinants of forest disturbance, and (3) changes in main forest types. Here, we define forest disturbance as full loss of forest cover due to forest management (e.g., clear-cutting), natural disturbances such as pests and storms (often followed by salvage logging), and deforestation (conversion to other land uses). Our first hypothesis was that there is more contemporary forest disturbance in areas that were not forested in the mid-19th century (hereafter ‘new forests’) compared to areas forested at that time (hereafter ‘old forests’) because new forest, mostly plantations, are more likely to be intensively managed than old forests and because the age and species composition of old forests makes them more resilient to disturbance. Second, we expected to find that contemporary disturbance is higher in new forests irrespective of environmental, socio-political and accessibility variation. Our third hypothesis was that ‘new forests’ have a higher proportion of coniferous forest than ‘old forests’, due to forest management practices of the late 19th and early 20th century, including widespread plantations.

Methods

Study area

We studied the Carpathian region in Eastern Europe (~265,000 km²), because the region experienced multiple socio-economic, political, and land management shifts over the past two

centuries, providing an ideal ‘natural experiment’ for the study of land-use legacies (Munteanu et al., 2014). The study area includes parts of two major eco-regions, the Carpathian Mountains and the Pannonian plains, and parts of Romania, Slovakia, Ukraine, Poland, Czech Republic and Hungary (Figure 2.1). Our study period from 1860 to 2010 captured a century and a half of land-use history, starting with the peak of the Habsburg Empire in the mid-19th century.

The land cover in the Carpathian mountains consists of a mosaic of forests, small agricultural fields, grassland areas, and scattered settlements (Kozak et al., 2013b; Kuemmerle et al., 2008). The Carpathian mountains harbor some of the largest contiguous forests of Europe, a high proportion of which are ecologically valuable (Knorn et al., 2012a). The Pannonian Plains consist mostly of large agricultural fields (Kuemmerle et al., 2009b; Schiller et al., 2010), intermixed with forest plantations and urban areas. The study area has a temperate climate with elevations up to 2500 m above sea level and varying microclimates (Kozak et al., 2013b). At low elevations deciduous forests (*Quercus* sp, *Fagus sylvatica*, *Carpinus betulus*, *Populus* sp, and *Robinia pseudoaccaia*) are common, while at high elevations coniferous forests are dominant (*Pinus* sp, *Picea abies*, *Abies alba*). Pine plantations for pulp production occur in the lowlands of Hungary and Romania (Bartha and Oroszi, 1995). The average tree line in the Carpathian mountains is 1600 m (Kozak et al., 2013a). Historically, the land cover of the Pannonian plains was grasslands and wetlands, but due to the high fertility of soils and population growth, many natural ecosystems were converted to agriculture (Bellon, 2004; Frisnyák S., 1990; Jordan et al., 2005; Szilassi et al., 2006).

The current land-cover patterns reflect centuries of land management. Overall, forest cover increased in the Carpathian region since the turn of the 20th century. Most of the study region

experienced a forest transition, i.e., a shift from net deforestation to net forest expansion, between the two World Wars (Kozak et al., 2007; Kuemmerle et al., 2011; Munteanu et al., 2014) and forest area increased especially after the breakdown of socialism in 1989, albeit at varying rates (Baumann et al., 2011; Griffiths et al., 2014). In the Carpathian mountains alone, forest area increased from 39.4% to 40.3% between 1985 and 2010 (Griffiths et al., 2014).

Large scale forest disturbances have occurred in the Carpathian region since the 19th century, partly because the forest management policies of the Habsburg Empire focused on timber production. After WWII, large areas of forest in Romania and Ukraine were harvested to pay war debts to the Soviet Union (Kligman and Verdery, 2011). Forest management for timber and pulp led to increased harvesting of hardwoods (Chirita, 1981) and to the establishment of spruce monocultures both before and during the socialist time period (Irland and Kremenetska, 2009; Keeton et al., 2013). After the collapse of socialism in 1991, disturbance rates were also high: from 1985 to 1995 disturbance peaks occurred in Poland, Czech Republic, Ukraine and northern Romania, and from 1995 to 2000 in the Romanian Carpathians (Griffiths et al., 2014). Overall, since 1985, as much as 20% of the Carpathian forests experienced stand-replacing disturbances (Griffiths et al., 2014). Following the collapse of the Soviet Union, most countries adopted restitution laws that reverted publicly-owned land to pre WWII owners (Bemmann and Grosse, 2001; Hartvigsen, 2014; Irimie and Essmann, 2009; Swinnen, 1999) who often harvested their forest for financial gains. However, differences in the timing of restitution laws, the strength of governance, and in economic and socio-demographic factors among countries caused differences in harvesting patterns (Griffiths et al., 2014, 2012; Kuemmerle et al., 2009d). In other words, while the institutional and socio-economic shifts associated with the transition to market-oriented economics certainly affected the rate of recent forest harvest in the Carpathian mountains

(Griffiths et al., 2014; Knorn et al., 2012b), the drivers of forest harvest and management have only been studied at broad scales (Levers et al., 2014) and the role of past land use for contemporary forest disturbance remains unclear.

Historic and contemporary land use and land cover data

We reconstructed historic forest area and historic forest types for the region in 1860s and 1960s from several collections of historical maps (Table 2.1), most of which were available in digital, georeferenced format (Arcanum Adatbázis Kft, 2015). We verified point location accuracy by a back-dating approach that associates the location of the digitized point with nearby landmarks in all available maps. From the historic maps we extracted forest cover information at two points, first during the Habsburg Empire (1805-1918) and second, during Socialism (1945-1990). We labeled 92,000 points arranged in a regular 2 x 2 km grid as either forest or non-forest for each time point (roughly 21% of the points being forested in each time slice). Where possible, we also mapped forest types as coniferous, mixed, or deciduous. Depending on the time period, forest type information was available for 62%-96% of the data points. Our point grid matched that of the 2007 INSPIRE directive (Infrastructure for Spatial Information in the European Community) and LUCAS (Land Use and Cover Area frame Survey, (Gallego and Delince', 2010).

To estimate contemporary forest disturbance, we mapped forest disturbance at 5-year time intervals from 1985 to 2010, and forest types for 1985s and 2010s, based on 30-m resolution Landsat TM/ETM+ image composites with an overall accuracy of 85.8% for the forest disturbance map (Griffiths et al., 2014, 2013a). We assigned the disturbance information at the specific point location to each grid point in our historic dataset. Our dependent variable, forest disturbance, captured loss of closed-canopy forest cover either due to harvesting, which was

predominantly clear-cutting, or natural disturbances, which were often followed by salvage logging. Selective logging was generally not captured, and we did not consider forest recovery or reforestation, which were beyond the scope of this paper. We analyzed forest disturbances between 1985 and 2010 (hereafter recent disturbance) because the interval captures two events that affected land management in the Carpathian region: the countries' transition to market economies (after 1989), and the accession to the European Union (in 2004 or 2007) of all countries in the study area, except Ukraine.

We defined a point as disturbed if it experienced forest loss in any 5-year time interval between 1985-2010 (Griffiths et al., 2014). We further restricted the definition of forest disturbance to only those areas that were forest in the 1960s maps in order to reconcile the remote sensing data (representing forest land cover), with the historic maps (representing forest land use). Via this step, we excluded cases of reforestation and spontaneous afforestation and of abandoned agricultural lands that were re-cultivated after 2000 (Griffiths et al., 2013b). Our data selection also minimized classifications errors in the 1985 classification due to limited Landsat image availability. We defined a point as not disturbed if it was continuously forested in the 1960s, 1985, 1990, 1995, 2000, 2005 and 2010. We eliminated data points above 1600 m, the average timberline in the Carpathians. In sum, for modelling purposes, we restricted our analysis to only those points that were forested in 1960s, a total of 19,947 points. For the forest type analysis we used a minimum of 12,497 points for the year 1860 and a maximum of 19,360 for the year 2010. Of all disturbed points, 43% experienced disturbance between 1985-1995, and approximately 4.6% of those were disturbed after 1985 and not reforested by 2010.

Land use legacy models

We selected 16 covariates that we expected to correlate to 1985-2010 forest disturbance. One covariate represented the historic forest cover in the 1860s and the rest captured environment (6 variables), socio- demographics (2 variables), and accessibility (7 variables, Table 2.2). We used the presence or absence of forest cover in 1860s as the indicator of land use legacy. We extracted all raster values for the 2-km point grid and used the binary response variables for forest disturbance (0/1) as the depended variable.

We fitted multiple logistic regression models (Hosmer and Lemeshow, 1980) to explain contemporary forest disturbance and to estimate the role of historic forest extent for recent disturbances. We fitted an overall model using the full dataset (19,947 data points), and country-specific models (Müller et al., 2009) to capture socio-economical and institutional diversity (Appendix 2.1). We performed variable selection using an exhaustive search (Hosmer et al., 2013) based on the Akaike Information Criterion (AIC) and retained the best performing model. Because we were interested in estimating the effect of historic land uses on recent change, we refitted the best model including the legacy variable for those countries where the best performing model did not include land use legacies (Ukraine, Hungary and Czech Republic). We found no changes in the signs of the model coefficients and also no major changes in coefficient values. The changes in AIC values were always less than 3 for the model including legacies. We tested for interactions between historic forest cover and two environmental variables (slope and elevation) to assess whether contemporary forest disturbance occurred in topographically marginal areas with historic forest cover (Müller and Zeller, 2002), but found that interaction terms were not significant and coefficients were close to zero (results not shown). We checked

the degree of spatial autocorrelation of the dependent variable using semivariograms of model residuals (Curran, 1988; Griffith, 2003) and did not find significant spatial autocorrelation.

To address our first objective regarding the importance of legacy effects for forest disturbance, we calculated the odds ratio of our logistic models, which compares the relative rates of forest disturbance in old and new forests depending on historic land cover. Values higher than 1 indicated higher odds of disturbance in new forests. We did not report significance levels or confidence intervals in our analysis because our data grid represents effectively a full census of historic and recent land cover and because the estimate of the effect that we observed is independent of sample size (Lohr, 2010). Finally, we checked model performance using receiver operating curves (ROC, (Freeman and Moisen, 2008)) and evaluated model utility by calculating the area under the ROC curve (AUC).

To address our second objective regarding the relationship between land-use legacies and other spatial determinants of forest disturbance, we compared the proportion of forest disturbance in old and new forests along a gradient of environmental and accessibility values (variables described in Table 2.2). We split our data into two groups (old and new forests), computed the proportion of disturbance in each group along gradients of other continuous variables in our data, and plotted partial dependence of the disturbance proportion.

To address our third objective of the effect of legacies on the forest types, we estimated the proportion of forest types at four time points. These estimates were based on historic land use maps for the 1860s and 1960s (Table 2.1) and forest-type classifications for 1985s and 2010s (Griffiths et al., 2014). For each time slice (Table 2.1) where forest type information was available, we calculated the percentage of each forest type in the overall forest. Forest type data

was complete for the time layers 1960, 1985 and 2010, but was absent for approximately 15% of the whole area in 1860, and 49% of contemporary Poland. We assumed that forest types in areas with no forest type information followed the same pattern as in areas where forest type information was available (Appendix 2.5). Here, we considered the legacy effects of past management by assessing changes in forest types and shifts in the proportion of coniferous, mixed and deciduous forest over time. We also analyzed the historic forest types of recently disturbed forests, the 2010 forest types of new forests for each country as well as the 1860 and 2010 forest types of old forest. When we calculated the percentage of each forest type in 2010 for areas not forested in 1860, we compared our results with the 1960s forest types in new forests to check for consistency between the two time periods.

Results

Land use legacies were strongly related to recent forest disturbance in the Carpathian region. Forest disturbance occurred more often in new forests, established after 1860, than in areas that were already forested in 1860. Legacy effects remained important when controlling for other determinants of forest disturbance. Together, land-use history, topography, climate, and accessibility explained patterns of forest disturbance in the Carpathian region well in our overall model, but there were differences among countries (see below). However, probabilities of disturbance were always higher in new forests, irrespective of environment, accessibility and socio-politics. Legacies of past land use also affected the proportions of forest types. Areas with a historically high coniferous cover (e.g., Czech Republic and Ukraine) decreased the percentage of coniferous forests by 2010. In Romania and Hungary, on the other hand, recent disturbance occurred mostly in historically deciduous and mixed forests, and the percentage of coniferous forests increased.

Forest disturbance legacies

Across all Carpathian countries, forest disturbance was more likely in areas that were not forested in 1860s (new forests), compared to forested areas in 1860s (old forest) (Figure 2.2). From the total set of 19,947 points, 73% were forested in 1860s and still forested in 1960s, 1985s and 2010s. Observed forest disturbance between 1985-2010 in old forests was 13% (of forested area), but 18% in new forest (Figure 2.2, Appendix 2.1). The ratios of forest disturbances in new relative to old forests, were consistently higher in all countries, with the maximum percentage of observed disturbance in old forests in the Czech Republic and Ukraine (22% and 16%, respectively) and the maximum percentage of observed disturbance in new forest in Czech Republic (26%), Slovakia (21%) and Ukraine (21%) (Figure 2.2).

The logistic regression results suggested that even when controlling for environmental, accessibility and socio-political factors, the odds of forest disturbance were 49% higher for new forests than for old forests (Figure 2.3). However, legacies varied by country: they were strongest in Poland (odds 88% higher), weakest in Ukraine (odds 34% higher), and not important in the Czech Republic and Hungary (Figure 2.3, Appendix 2.2). The AUC for the seven models varied from 0.62 (Czech Republic) to 0.78 (Poland). The overall model had an AUC value of 0.66. Our findings suggest that even after controlling for the environmental, accessibility and socio-political differences in the study region, the historic land uses played an important role in determining the location of recent forest disturbance.

Spatial determinants of disturbance

Our models confirmed the importance of environment (topography, temperature, length of growing season) and accessibility (distance to cities and settlements) for forest disturbance in all

Carpathian countries. In the overall model, slope, annual mean temperature, and the length of the growing season were important predictors, alongside our country dummy variable, which captured at national level, processes not captured by other variables, such as land reforms, strength of institutions, or accessibility differences (Levers et al., 2014; Müller and Sikor, 2006). In all countries, areas with steep slopes were less likely to be disturbed, and areas afar from major cities, and close to human settlement were more likely to be disturbed (Appendix 2.3). Our models did not show strong quantitative evidence for a relationship of forest disturbance with population density in the 1990s or distance to railroads (Appendix 2.3).

Our comparison of the proportion of disturbance in old and new forests across the full range of slopes, temperatures, precipitations, and accessibilities, indicated that the proportion of disturbance was consistently higher in new forests both when we summarized our data (Figure 2.4) and when modelling disturbance while controlling for other spatial determinants of change (Appendix 2.4). Disturbance decreased with increasing slope, temperature, crop suitability, and length of the growing season in both old and new forests, but the consistently higher proportion of disturbance in new forests remained. The shorter the growing season, the more new forests were disturbed. Within 15 km of roads, new forests were more likely to be disturbed, and although at distances higher than 20 km, older forests were more likely to be disturbed, we had only few observations in this data range. Forests closer to settlements were also more likely to be disturbed.

Changes in main forest types

We analyzed forest type abundance in the Carpathian region for four time points (1860s, 1960s, 1985s, and 2010s). We found that coniferous forest cover declined in all countries except

Romania and Hungary. Slovakia and Czech Republic reached their peak coniferous forest cover in the early socialist period (1960s) while Hungary, Romania and Ukraine reached a peak in the late 1980s. The forests of most countries were mainly coniferous and mixed in 2010 (65% to 98% in Ukraine and Romania respectively). Only Slovakia and Hungary had over 50% deciduous forests in 2010. Overall, coniferous cover in 2010 was 24%, roughly 5% lower than in 1860.

When we analyzed the 1860s forest types of areas that were disturbed after 1985, we found that over 50% of the disturbance occurred in 1860s coniferous stands, except for Romania and Hungary where the proportion of coniferous forest was low in the 1860s (Figure 2.5a). Overall, the 2010s forest types in new forests were 28% coniferous, 27% mixed, and 40% deciduous, but there were marked differences among countries: in Romania, Hungary and Slovakia, over 50% of the forests were deciduous and mixed, while Ukraine and Czech Republic had an approximately equal proportion of deciduous, coniferous, and mixed forests (Figure 2.5d). The proportion of coniferous and deciduous new forests was very similar in 1960 and 2010. Most recent disturbances (1985-2010) occurred in 1860s coniferous stands, less Hungary, where there were few coniferous to start out with (Figure 2.5b). The proportion of forest types in old forests shifted towards less coniferous in all countries except Romania and Hungary (Figure 2.5c and e). Generally, there was a higher proportion of coniferous in new forests than in old forests across countries (Figure 2.5d and e). Forest disturbances between 1985 and 1995 also affected forest types in 2010. Of all disturbed points that were initially coniferous, only 37% remained coniferous after disturbance. Conversely, 90% of the disturbed deciduous and mixed forests retained their composition.

Discussion

Our results showed that land-use legacies were an important spatial determinant of forest disturbance in the Carpathian region. These results are important because land-use legacies are rarely included in models analyzing drivers of forest change (Levers et al., 2014; Pazúr et al., 2014; Verburg et al., 2009). Our results are unique among land-use legacy studies (Bellemare et al., 2002; Dupouey et al., 2002a; Foster et al., 2003), because we provided evidence for legacies affecting forest management decisions and subsequent rates of forest disturbance. Land-use legacies from 150 years ago greatly affected recent patterns of forest disturbance even when controlling for environment, accessibility and socio-political factors. Indeed, areas not forested in the 1860s (new forests) had 49% higher odds of recent disturbance, than areas forested in 1860s (old forests). The probability of disturbance was consistently higher in new forests, across the full range of covariates, underpinning the importance of considering land-use legacies when assessing and modeling forest change. Forest management caused a decrease of coniferous in old forests between 1860 and 2010 and a higher percentage of coniferous in new forests than in old forests.

Forest disturbance legacies

The Carpathian region is a hotspot of cultural, political and socio-economic diversity with a rich land management history (Munteanu et al., 2014), thus providing an ideal ‘natural experiment’ to assess the effects of land-use legacies on rates of disturbance and the abundance of forest types. In the 19th century, a large proportion of the study region was part of the Habsburg Monarchy, later the Austro-Hungarian Empire, which managed forests intensively for wood production. In the mountainous regions of contemporary northern Romania, Ukraine, northern Slovakia and southern Poland (historical regions of Bessarabia, Bukovina, Galicia, and Maramures), forest

harvest intensified during the Habsburg rule (Bohateret, 2012). In addition, fast growing, productive tree species such as Norway spruce (*Picea abies*) were widely planted for pulp production and erosion control. In the lowlands, following a period of agricultural expansion and timber scarcity, forests were planted outside their prior ranges during Hungarian and Austrian rule, and later again during Socialism (Konkoly-Gyuró et al., 2011, 2012). Especially Hungary, Slovakia and Romania planted large areas of poplar (*Populus* sp.), black locust (*Robinia pseudoacacia*) and pine (*Pinus* sp., (Bartha and Oroszi, 1995; Chirita, 1981; Konkoly-Gyuró et al., 2011). With hardwood rotation ages of about 70 years for pulp and 120 years for saw-timber (Chirita, 1981; Disescu, 1954), it is likely that early 20th century plantations have recently reached a harvestable age, which may be one reason for the high rate of forest disturbance in new forests. Furthermore, natural disturbance events, such as wind throws or insect outbreaks, preferentially affect forests vulnerable through previous management practices, such as plantations of even-aged monocultures (Klopcic et al., 2009; Schelhaas et al., 2003; Svoboda et al., 2012). When natural disturbances occur, salvage logging is common, which may be another explanation for the strong legacy effect that we observed. Spruce plantations in the Carpathians are susceptible to pests such as bark beetle (Keeton et al., 2010), pollution (Carrier and Krippel, 2009; Main-Knorn et al., 2009; Modrzyński, 2003), floods (Glenz et al., 2006), wind and snowstorms (Falt'an et al., 2009), and fluctuations in climate (Bouriaud and Popa, 2008a), all potentially causing higher disturbance rates in new forests. We caution though, that plantations are also common in old forests, especially in mountain regions of Romania where historic mixed stands have been recently replaced by spruce. Other past management practices that may still affect rates of disturbance are forest grazing and litter raking, common in the Habsburg Empire since the 19th century (Erb et al., 2013). Likewise, historic forest ownership structures may

affect current disturbance rates when land owners decide to preserve or to manage forests for timber production (Ostafin, 2009). Irrespective of the mechanisms, which likely vary in space, our results showed that recent forest management is greatly restricted by historic land uses and forest management decisions.

Spatial determinants of disturbance

We analyzed the proportion of forest disturbance (logging, and natural disturbances typically followed by salvage logging) in old and new forests in relation to environmental and accessibility variables and found that disturbance was consistently higher in new forests, across the entire range of spatial determinants of change. The relationship of disturbance to the different determinants of change is interesting in its own right as well though.

We included variables that captured known spatial determinants of land-use change and forest management (Geist and Lambin, 2001; Müller et al., 2013; Pazúr et al., 2014). We did not include data on policies, markets and economic factors that may underlie forest disturbance patterns (Amacher et al., 2003; Beach et al., 2005; Geist and Lambin, 2001; Lambin et al., 2001) because these were only available at country level, and our sample size did not allow us to examine country effects. However, we captured these differences partly via the country dummy variable and explored them in country-specific models.

Our results indicated that areas with less rough terrain were more likely to be disturbed (probably due to difficult access). Our results are thus consistent with other analyses of drivers of forest cover change, which found higher disturbance probability in areas with less rough terrains and mild slopes (Levers et al., 2014; Nagendra et al., 2003; Wendland et al., 2011), and that such areas often represent deciduous and mixed forests at lower elevations. We found slightly more

disturbance in areas farther away from cities and roads, but most of our data was concentrated at distances less than 10km away from roads.

Areas with low crop suitability experienced higher rates of disturbance. High rates of forest disturbance on low quality soils are common in areas that have been spruce plantations for long times (Chirita, 1981) because of soil acidification. Furthermore, new forests in Hungary, Romania and Slovakia were often planted in areas prone to erosion and on poor soils that were depleted of nutrients. Areas with low soil quality generally have more forest health problems (Schulze et al., 1989). We note that deforestation is often higher on better soils (Pfaff, 1999; Veldkamp et al., 1992), but this is due to their suitability for agriculture (Etter et al., 2006; Grau et al., 2005; Veldkamp et al., 1992) rather than natural causes due to forest management. In the recent land use history of the Carpathian region, agricultural clearing is not common, and only some of the recently abandoned agricultural land has been brought back into production (Griffiths et al., 2013b).

We found interesting differences among countries in terms of the importance of land use legacies in relation to other spatial determinants. Areas afar from cities were less likely to be disturbed in Poland and Czech Republic than in Romania and Ukraine, most likely because the former countries have reliable forest protection system and a high percentage of state-owned land (Kuemmerle et al., 2009c), and have experienced fewer institutional changes and shifts in environmental policies in recent decades (“Polityka ekologiczna panstwa w latach 2009-2012,” 2008). On the other hand, Romania and Ukraine had higher occurrence of disturbance in remote areas, most likely because institutions there are weaker, forest restitution caused widespread harvesting in private forests (Giurgiu, 2010a), and protection is not always effective (Irland and Kremenetska, 2009; Knorn et al., 2012b). Despite repeated suggestions that a high portion of the

logging might be illegal due to poor regulatory framework (Knorn et al., 2012b; Kuemmerle et al., 2009a), it remains hard to quantify to what extent this influenced our results. In Hungary, disturbance was higher near settlements and rivers and in accessible areas that could be easily harvested. We speculate that this could be due to policies for erosion control, soil quality enhancement, and pulp plantations along river ways (Bartha and Oroszi, 1995; Konkoly-Gyuró et al., 2012).

Individual disturbance events can affect the observed relationship between old and new forests, if these disturbances are very large. In Slovakia, a large windthrow occurred in the Tatra Mountains in 2004 (Falt'an et al., 2009; Griffiths et al., 2014). A large part of the windblown area was not forested in the 1860s and was subsequently planted with spruce thus becoming susceptible to natural disturbances (Falt'an et al., 2009). A large part of the windblown area was already affected by a historic disturbances around 1915s and 1940s (Zielonka et al., 2010) and planted with spruce. The high likelihood of disturbance in new forests that our analysis uncovered in Slovakia may thus be at least partly influenced by this singular natural disturbance event.

Change in forest type

Land use legacies also affected forest types. Overall, new forest had a higher percentage of conifers than old forests, and the percentage of coniferous trees in old forests was higher in 2010 than in 1860 in all countries except Romania and Hungary, where the coniferous cover was low in the 1860s. Our results supported prior findings about the importance of historic management on recent forest composition, species abundance, and ecosystem health (Bellemare et al., 2002; Dupouey et al., 2002a; Foster et al., 2003; Wallin et al., 1994). In most Carpathian countries,

historic extensive harvest for wood production and the susceptibility of spruce plantations to natural disturbances resulted in a decline in coniferous forests. Where natural regeneration occurred following clear-cuts in the late 19th and early 20th century, the forest shifted towards a larger proportion of mixed and deciduous tree types, and this was in particular the case in Ukraine, Slovakia and Czech Republic. Romania and Hungary had a high proportion of deciduous and mixed forests in the 1860s and an increasing proportion of coniferous over time. Here, contemporary forest disturbance occurred mostly in historically mixed and deciduous forests and we explain this diverging legacy by economically-driven plantations during Habsburg and Socialist times (Chirita, 1981; Dincă, 1955; Konkoly-Gyuró et al., 2011).

We argue that the relative abundance of forest types, similar to disturbance patterns, is the result of legacies related to forest management practices in each country. Norway spruce was the predominant production tree species across the Carpathian region since the mid-19th century (Irland and Kremenetska, 2009) due to its fast growth rate and because it provided both pulp and timber. Romania and Hungary increased their percentage of coniferous forests over historic deciduous forests due to Soviet pressure during the 1950s (Banu, 2004; Kligman and Verdery, 2011) to repay war debts in timber. Furthermore, our data indicated a strong increase of coniferous forests in Romania and Ukraine between 1960 and 1985. Here, after WW II, forests transferred to state ownership and intensive forest management for wood and pulp production led to widespread spruce plantations (Chirita, 1981; Irland and Kremenetska, 2009). However, while planting clearcuts with spruce was common in many regions, others relied on natural succession (Dincă, 1955; Gosciniński, 2014).

We caution that the interpretation of forest type changes relies on the assumption that the composition of all forests followed the same distribution as the forests for which we had forest

type information in the 1860s maps. We did not report country results for Poland here, because we were missing 1860s forest type data for half of its area in the study region. We caution that the legacy effects that we revealed could vary slightly depending on data preprocessing, the definition of legacy effects and the methodology used to map forest disturbance. Here, we relied on forest-non forest data and on forest type classifications as indicators of past land management and legacies. We tested the consistency of the observed differences between old and new forests using nine different definitions of disturbance, and found a consistent pattern of more disturbances in new forest in all cases (results not shown). Furthermore, the historical data used in our analysis depicts land use, whereas the recent remote sensing analysis captures land cover, and in order to make the two datasets comparable, we restricted our analysis to forest disturbances within 1960s forests. Including reforestation and disturbance on abandoned fields in the analysis dataset could also potentially alter the observed legacy effect, but analyzing this was beyond the scope of this paper. We also caution that our data does not capture historic forest harvests – but including historic clear-cuts in our models, would likely increase the observed disturbance difference between old and new forests. Given the nature of our dataset our results should be interpreted at broad, national and regional scales. Furthermore, in the Carpathian region, our analysis may reflect particularly strong legacy effects, due to political shifts and land management changes (Munteanu et al., 2014).

Conclusions

Our study showed that contemporary forest disturbance patterns were heavily influenced by 150-year old land-use legacies, even when controlling for environmental, accessibility, socio-political spatial determinants. Specifically, forest disturbance was more likely in areas where forests occurred for a shorter time period. This is good news for the conservation of Carpathian forests

that have been in place for a longer time and highlights that past land uses are important when deciding which areas to protect or harvest. In a scientific context, the consideration of past land uses as spatial determinants of change could enhance the performance of forest change assessments and of predictions of future land change trajectories. Our results suggest strong land use legacy effects can be present over centuries. This is of concern because we are currently in a phase of rapid land use change globally (Baumann et al., 2014; Grau et al., 2008; Meyfroidt and Lambin, 2009), and land use legacies may restrict management possibilities, in the Carpathians and worldwide, while creating legacies for future generations. Forests are lost at rapid rates in many areas of the globe (Hansen et al., 2013). Knowledge of legacy effects of these trends could help land managers when making decisions which areas to use for timber production, agriculture, or conservation in the future.

References

- Amacher, G.S., Conway, M.C., Sullivan, J., 2003. Econometric analyses of nonindustrial forest landowners: Is there anything left to study? *J For Econ* 9, 137–164.
- Arcanum Adatbázis Kft, 2015. Historical Maps of the Habsburg Empire [WWW Document]. Mapire.eu. URL <http://mapire.eu/en/>
- Banu, F., 2004. Asalt asupra economiei Romaniei de la Solagra la SOVROM (1936-1956) (in Romanian). Nemira, Bucuresti.
- Bartha, D., Oroszi, S., 1995. Magyar erdők. (Hungarian Froests), in: M, D. (Ed.), Pannon Enciklopédia. Főszerk. Járayné Komlódy. pp. 220–230.
- Başnou, C., Álvarez, E., Bagaria, G., Guardiola, M., Isern, R., Vicente, P., Pino, J., 2013. Spatial patterns of land use changes across a Mediterranean metropolitan landscape: implications for biodiversity management. *Environ Manage* 52, 971–80.
- Baumann, M., Kuemmerle, T., Elbakidze, M., Ozdogan, M., Radeloff, V.C., Keuler, N.S., Prishchepov, A. V., Kruhlov, I., Hostert, P., 2011. Patterns and drivers of post-socialist farmland abandonment in Western Ukraine. *Land use policy* 28, 552–562.
- Baumann, M., Radeloff, V.C., Avedian, V., Kuemmer, 2014. Land-use change in the Caucasus during and after the Nagorno-Karabakh conflict. *Reg Environ Chang*.
- Beach, R.H., Pattanayak, S.K., Yang, J.-C., Murray, B.C., Abt, R.C., 2005. Econometric studies of non-industrial private forest management: a review and synthesis. *For Policy Econ* 7, 261–281.
- Bellemare, J., Motzkin, G., Foster, D.R., Forest, H., 2002. Legacies of the agricultural past in the forested present : an assessment of historical land-use effects on rich mesic forests. *J Biogeography* 29, 1401–1420.
- Bellon, T., 2004. Living together with nature. Farming on the river flats in the valley of the Tisza. *Acta Ethnogr Hungarica* 49, 243–256.
- Bemmann, A., Grosse, W., 2001. Changes of forest ownership in the countries of transition in Middle and East Europe. *Allg Forst Und Jagdzeitung* 172, 173–183.
- Bezák, P., Mitchley, J., 2014. Drivers of change in mountain farming in Slovakia: from socialist collectivisation to the Common Agricultural Policy. *Reg Environ Chang* 14, 1343–1356.
- Bohateret, V.M., 2012. Readjusting Romania ' s Forestry Policy with a View to the Year 2050. *J Settlements Spat Plan* 1, 27–42.
- Boucher, Y., Grondin, P., Auger, I., 2013. Land use history (1840–2005) and physiography as determinants of southern boreal forests. *Landsc Ecol* 29, 437–450.
- Bouriaud, O., Popa, I., 2008. Comparative dendroclimatic study of Scots pine, Norway spruce, and silver fir in the Vrancea Range, Eastern Carpathian Mountains. *Trees* 23, 95–106.

- Brown, D.G., Page, S., Riolo, R., Zellner, M., Rand, W., 2005. Path dependence and the validation of agent based spatial models of land use. *Int J Geogr Inf Sci* 19, 153–174.
- Brown, J.R., Archer, S., 1989. Woody plant invasion of grasslands : establishment of honey mesquite (*Prosopis glandulosa* var . *glandulosa*) on sites differing in herbaceous biomass and grazing history 80, 19–26.
- Carrier, J.G., Krippel, E., 2009. Comprehensive Study of European Forests Assesses Damage and Economic Losses from Air Pollution. *Environ Conserv* 17, 365.
- Chirita, C., 1981. Padurile Romaniei. Studiu Monografic. Ed Academiei RSR, Bucharest.
- Cramer, V.A., Hobbs, R.J., Standish, R.J., 2008. What's new about old fields? Land abandonment and ecosystem assembly. *Trends Ecol Evol* 23, 104–12.
- Curran, P.J., 1988. The semivariogram in remote sensing: An introduction. *Remote Sens Environ* 24, 493–507.
- Cushman, S.A., Wallin, D.O., 2000. Rates and patterns of landscape change in the Central Sikhote-alin Mountains, Russian Far East. *Landsc Ecol* 15, 643–659.
- Dincă, I., 1955. Realizarea primului cincinal la împăduriri, important succes al politicii de refacere a fondului forestier al R.P.R. *Rev Padur* 9.
- Disescu, R., 1954. Exploatabilitatea, virsta exploatabilitatii si ciclul de productie la arboretele de molid (in Romanian). *Analele ICAS XV*, 529–246.
- Dupouey, J.L., Dambrine, E., Laffite, J.D., Moares, C., 2002. Irreversible Impact of Past Land Use on Forest Soils and Biodiversity. *Ecology* 83, 2978–2984.
- Eberhardt, R.W., Foster, D.R., Motzkin, G., Hall, B., 2011. Conservation of Changing Landscapes : Vegetation and Land-Use History of Cape Cod National Seashore. *Ecol Appl* 13, 68–84.
- Erb, K.-H., Kastner, T., Luyssaert, S., Houghton, R.A., Kuemmerle, T., Olofsson, P., Haberl, H., 2013. Bias in the attribution of forest carbon sinks. *Nat Clim Chang* 3, 854–856.
- Etter, A., McAlpine, C., Pullar, D., Possingham, H., 2006. Modelling the conversion of Colombian lowland ecosystems since 1940: drivers, patterns and rates. *J Environ Manage* 79, 74–87.
- Falt'an, V., Katina, S., Bánovský, M., Pazúrová, Z., 2009. The influence of site conditions on the impact of windstorms on forests: The case of the High Tatras Foothills (Slovakia) in 2004. *Morav Geogr Reports* 17, 44–52.
- Foley, J.A., DeFries, R.S., Asner, G.P., Barford, C., Bonan, G., Carpenter, S.R., Chapin, F.S., Coe, M.T., Daily, G.C., Gibbs, H.K., Helkowski, J.H., Holloway, T., Howard, E.A., Kucharik, C.J., Monfreda, C., Patz, J.A., Prentice, I.C., Ramankutty, N., Snyder, P.K., 2005. Global consequences of land use. *Science* (80-) 309, 570–574.
- Foster, D.R., Swanson, F.J., Aber, J., Burke, I., Brokaw, N., Tilman, D., Knapp, A., 2003. The

- importance of land-use legacies to ecology and conservation. *Bioscience* 53, 77–88.
- Fredrickson, E., Havstad, K.M., Estell, R., Hyder, P., 1998. Perspectives on desertification: south-western United States. *J Arid Environ* 39, 191–207.
- Freeman, E.A., Moisen, G.G., 2008. A comparison of the performance of threshold criteria for binary classification in terms of predicted prevalence and kappa. *Ecol Modell* 217, 48–58.
- Frisnyák S., 1990. Magyarország történeti földrajza. (Historical geography of Hungary). Tankönyvkiadó, Budapest.
- Gallego, J., Delincé', J., 2010. The European land use and cover area-frame statistical survey, in: Benedetti, R., Bee, M., Espa, G., Piersimoni, F. (Eds.), *Agricultural Survey Methods*. John Wiley & Sons, pp. 151–168.
- Geist, H.J., Lambin, E.F., 2001. What Drives Tropical Deforestation? A meta-analysis of proximate and underlying causes of deforestation based on subnational case study evidence., *LUCC Report Series*. Louvain-la-Neuve.
- Gerard, F., Petit, S., Smith, G., Thomson, A., Brown, N., Manchester, S., Wadsworth, R., Bugár, G., Halada, L., Bezak, P., Boltiziar, M., De badts, E., Halabuk, A., Mojses, M., Petrovič, F., Gregor, M., Hazeu, G., Mucher, C. a., Wachowicz, M., Huitu, H., Tuominen, S., Kohler, R., Olschofsky, K., Ziese, H., Kolar, J., Sustera, J., Luque, S., Pino, J., Pons, X., Roda, F., Roscher, M., Feranec, J., 2010. Land cover change in Europe between 1950 and 2000 determined employing aerial photography. *Prog Phys Geogr* 34, 183–205.
- Gibbens, R.P., McNeely, R.P., Havstad, K.M., Beck, R.F., Nolen, B., 2005. Vegetation changes in the Jornada Basin from 1858 to 1998. *J Arid Environ* 61, 651–668.
- Giurgiu, V., 2010. Declinul suprafeței pădurilor și marginalizarea împăduririlor. *Rev Padur* 2.
- Glenz, C., Schlaepfer, R., Iorgulescu, I., Kienast, F., 2006. Flooding tolerance of Central European tree and shrub species. *For Ecol Manage* 235, 1–13.
- Gosciniński, W., 2014. Raport o stanie lasow w Polsce. Warszawa, Poland.
- Grau, H.R., Gasparri, N.I., Aide, T.M., 2005. Agriculture expansion and deforestation in seasonally dry forests of north-west Argentina. *Environ Conserv* 32, 140.
- Grau, H.R., Gasparri, N.I., Aide, T.M., 2008. Balancing food production and nature conservation in the Neotropical dry forests of northern Argentina. *Glob Chang Biol* 14, 985–997.
- Griffith, D.A., 2003. *Spatial Autocorrelation and Spatial Filtering: Gaining Understanding Through Theory and Scientific Visualization*. Springer.
- Griffiths, P., Kuemmerle, T., Baumann, M., Radeloff, V.C., Abrudan, I. V., Lieskovský, J., Munteanu, C., Ostapowicz, K., Hostert, P., 2014. Forest disturbances, forest recover, and changes in forest types across the Carpathian ecoregion from 1985 to 2010 based on Landsat image composites. *Remote Sens Environ* 151, 72–88.
- Griffiths, P., Kuemmerle, T., Kennedy, R.E., Abrudan, I. V., Knorn, J., Hostert, P., 2012. Using

- annual time-series of Landsat images to assess the effects of forest restitution in post-socialist Romania. *Remote Sens Environ* 118, 199–214.
- Griffiths, P., Linden, S. Van Der, Kuemmerle, T., Hostert, P., 2013a. A Pixel-Based Landsat Compositing Algorithm for Large Area Land Cover Mapping 6, 2088–2101.
- Griffiths, P., Müller, D., Kuemmerle, T., Hostert, P., 2013b. Agricultural land change in the Carpathian ecoregion after the breakdown of socialism and expansion of the European Union. *Environ Res Lett* 8, 045024.
- Hansen, M.C., Potapov, P. V, Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A., Thau, D., Stehman, S. V, Goetz, S.J., Loveland, T.R., Kommareddy, A., Egorov, A., Chini, L., Justice, C.O., Townshend, J.R.G., 2013. High-resolution global maps of 21st-century forest cover change. *Science* 342, 850–3.
- Hartvigsen, M., 2014. Land reform and land fragmentation in Central and Eastern Europe. *Land use policy* 36, 330–341.
- Hosmer, D.W., Jr., Lemeshow, S., Sturdivant, R.X., 2013. *Applied Logistic Regression*. John Wiley & Sons.
- Hosmer, D.W., Lemeshow, S., 1980. Goodness of fit tests for the multiple logistic regression model. *Commun Stat - Theory Methods* 9, 1043–1069.
- Irimie, D.L., Essmann, H.F., 2009. Forest property rights in the frame of public policies and societal change. *For Policy Econ* 11, 95–101.
- Ireland, L.C., Kremenetska, E., 2009. Practical Economics of Forest Ecosystem Management: The Case of the Ukrainian Carpathians, in: Soloviy, I.P., Keeton, W.S. (Eds.), *Ecological Economics and Sustainable Forest Management: Developing a Transdisciplinary Approach for the Carpathian Mountains*. Ukrainian National Forestry University Press, Lviv, p. 432.
- Jordan, G., Rompaey, A. van, Szilassi, P., Csillag, G., Mannaerts, C., Woldai, T., 2005. Historical land use changes and their impact on sediment fluxes in the Balaton basin (Hungary). *Agric Ecosyst Environ* 108, 119–133.
- Keeton, W.S., Angelstam, P., Baumflek, M., Bihun, Y., Chernyavskyy, M., Crow, M.S., Deyneka, Y., Elbakidze, M., Farley, J., Kovalyshyn, V., Mahura, B., Myklush, S., Nunery, J.R., Solovity, I., Zahvoyska, L., 2013. Sustainable forest management alternatives for the Carpathian Mountain region, with a focus on Ukraine, in: Kozak, J., Ostapowicz, K., Bytnerowicz, A., Wyzga, B. (Eds.), *The Carpathians: Integrating Nature and Society Towards Sustainability*. Springer-Verlag, Berlin and Heidelberg, pp. 331–352.
- Keeton, W.S., Chernyavskyy, M., Gratzner, G., Main-Knorn, M., Shpylchak, M., Bihun, Y., 2010. Structural characteristics and aboveground biomass of old-growth spruce–fir stands in the eastern Carpathian mountains, Ukraine. *Plant Biosyst* 144, 148–159.
- Kligman, G., Verdery, K., 2011. *Peasants under siege: the collectivization of Romanian agriculture, 1949-1962*. Princeton University Press.

- Klopčič, M., Poljanec, A., Gartner, A., Boncina, A., 2009. Factors Related to Natural Disturbances in Mountain Norway Spruce (*Picea abies*) Forests in the Julian Alps. *Ecoscience* 16, 48–57.
- Knorn, J., Kuemmerle, T., Radeloff, V.C., Keeton, W.S., Gancz, V., Biriş, I.-A., Svoboda, M., Griffiths, P., Hagatis, A., Hostert, P., 2012a. Continued loss of temperate old-growth forests in the Romanian Carpathians despite an increasing protected area network. *Environ Conserv* 40, 182–193.
- Knorn, J., Kuemmerle, T., Radeloff, V.C., Szabo, A., Mindrescu, M., Keeton, W.S., Abrudan, I. V., Griffiths, P., Gancz, V., Hostert, P., 2012b. Forest restitution and protected area effectiveness in post-socialist Romania. *Biol Conserv* 146, 204–212.
- Konkoly-Gyuró, É., Bacsárdi, V., Tirászi, Á., 2012. Perception of landscape changes in three trans-boundary focus areas based on oral history surveys with local inhabitants, stakeholders and experts. *TransEcoNet WP6: Identities and Strategies. Action 6.1. Sopron*.
- Konkoly-Gyuró, E., Nagy, D., Balázs, P., Király, G., 2011. Assessment of land cover change in western Hungarian landscapes, in: Balázs, P., Konkoly-Gyuró, É. (Eds.), *TransEcoNet Workshop on Landscape History Proceedings*. University of West Hungary Press, pp. 75–89.
- Kozak, J., Estreguil, C., Troll, M., 2007. Forest cover changes in the northern Carpathians in the 20th century: a slow transition. *J Land Use Sci* 127–146.
- Kozak, J., Ostapowicz, K., Byterowicz, A., Wyzga, B., 2013a. The Carpathian Mountains: Challenges for the Central and Eastern European Landmark, in: *The Carpathians: Integrating Nature and Society Towards Sustainability*. Springer, Berlin, Heidelberg, pp. 1–11.
- Kozak, J., Ostapowicz, K., Bytnerowicz, A., Wyzga, B. (Eds.), 2013b. *The Carpathians: Integrating Nature and Society Towards Sustainability, Environmental Science and Engineering*. Springer Berlin Heidelberg, Berlin, Heidelberg.
- Kuemmerle, T., Chaskovskyy, O., Knorn, J., Radeloff, V.C., Kruhlov, I., Keeton, W.S., Hostert, P., 2009a. Forest cover change and illegal logging in the Ukrainian Carpathians in the transition period from 1988 to 2007. *Remote Sens Environ* 113, 1194–1207.
- Kuemmerle, T., Hostert, P., Radeloff, V.C., van der Linden, S., Perzanowski, K., Kruhlov, I., 2008. Cross-border Comparison of Post-socialist Farmland Abandonment in the Carpathians. *Ecosystems* 11, 614–628.
- Kuemmerle, T., Hostert, P., St-Louis, V., Radeloff, V.C., 2009b. Using image texture to map farmland field size: a case study in Eastern Europe. *J Land Use Sci* 4, 85–107.
- Kuemmerle, T., Kozak, J., Radeloff, V.C., Hostert, P., 2009c. Differences in forest disturbance among land ownership types in Poland during and after socialism. *J Land Use Sci* 4, 73–83.
- Kuemmerle, T., Müller, D., Griffiths, P., Rusu, M., 2009d. Land use change in Southern Romania after the collapse of socialism. *Reg Environ Chang* 9, 1–12.

- Kuemmerle, T., Oloffson, P., Chaskovskyy, O., Baumann, M., Ostapowicz, K., Woodcock, C.E., Houghton, R.A., Hostert, P., Keeton, W.S., Radeloff, V.C., 2011. Post-Soviet farmland abandonment, forest recovery, and carbon sequestration in western Ukraine. *Glob Chang Biol* 17, 1335–1349.
- Lambin, E.F., Geist, H.J. (Eds.), 2006. *Land-Use and Land-Cover Change - Local Processes and Global Impacts*, Global Cha. ed. Springer.
- Lambin, E.F., Turner, B.L., Geist, H.J., Agbola, S.B., Angelsen, A., Bruce, J.W., Coomes, O.T., Dirzo, R., Fischer, G., Folke, C., George, P.S., Homewood, K., Imbernon, J., Leemans, R., Li, X., Moran, E.F., Mortimore, M., Ramakrishnan, P.S., Richards, J.F., Skånes, H., Steffen, W., Stone, G.D., Svedin, U., Veldkamp, T.A., Vogel, C., Xu, J., 2001. The causes of land-use and land-cover change: moving beyond the myths. *Glob Environ Chang* 11, 261–269.
- Levers, C., Verkerk, P.J., Müller, D., Verburg, P.H., Butsic, V., Leitão, P.J., Lindner, M., Kuemmerle, T., 2014. Drivers of forest harvesting intensity patterns in Europe. *For Ecol Manage* 315, 160–172.
- Lohr, S., 2010. *Sampling: Design and Analysis*, 2nd ed. Brooks/Cole, Boston MA.
- Main-Knorn, M., Hostert, P., Kozak, J., Kuemmerle, T., 2009. How pollution legacies and land use histories shape post-communist forest cover trends in the Western Carpathians. *For Ecol Manage* 258, 60–70.
- Meyfroidt, P., Lambin, E.F., 2009. Forest transition in Vietnam and displacement of deforestation abroad. *Proc Natl Acad Sci USA* 106, 16139–44.
- Modrzyński, J., 2003. Defoliation of older Norway spruce (*Picea abies* /L./ Karst.) stands in the Polish Sudety and Carpathian mountains. *For Ecol Manage* 181, 289–299.
- Morris, L.R., Monaco, T. a., Sheley, R.L., 2011. Land-Use Legacies and Vegetation Recovery 90 Years After Cultivation in Great Basin Sagebrush Ecosystems. *Rangel Ecol Manag* 64, 488–497.
- Motzkin, G., Foster, D.R., 2002. Grasslands, heathlands and shrublands in coastal New England: historical interpretations and approaches to conservation. *J Biogeogr* 29, 1569–1590.
- Müller, D., Kuemmerle, T., Rusu, M., Griffiths, P., 2009. Lost in transition: determinants of post-socialist cropland abandonment in Romania. *J Land Use Sci* 4, 109–129.
- Müller, D., Leitão, P.J., Sikor, T., 2013. Comparing the determinants of cropland abandonment in Albania and Romania using boosted regression trees. *Agric Syst* 117, 66–77.
- Müller, D., Sikor, T., 2006. Effects of postsocialist reforms on land cover and land use in South-Eastern Albania. *Appl Geogr* 26, 175–191.
- Müller, D., Zeller, M., 2002. Land use dynamics in the central highlands of Vietnam: a spatial model combining village survey data with satellite imagery interpretation. *Agric Econ* 27, 333–354.

- Munteanu, C., Kuemmerle, T., Boltiziar, M., Butsic, V., Gimmi, U., Kaim, D., Király, G., Konkoly-Gyuró, É., Kozak, J., Lieskovský, J., Mojses, M., Müller, D., Ostafin, K., Ostapowicz, K., Shandra, O., Štych, P., Walker, S., Radeloff, V.C., 2014. Forest and agricultural land change in the Carpathian region—A meta-analysis of long-term patterns and drivers of change. *Land use policy* 38, 685–697.
- Nagendra, H., Southworth, J., Tucker, C., 2003. Accessibility as a determinant of landscape transformation in western Honduras: linking pattern and process. *Landsc Ecol* 18, 141–158.
- National Research Council, 1998. *People and Pixels: Linking Remote Sensing and Social Science*. The National Academies Press, Washington, DC.
- Ostafin, K., 2009. Zmiany granicy rolno-leśnej w środkowej części Beskidu Średniego od połowy XIX wieku do 2005 roku (in Polish). roku, Wydaw UJ. Wydawnictwo Uniwersytetu Jagiellońskiego Kraków.
- Pazúr, R., Lieskovský, J., Feranec, J., Oľahel', J., 2014. Spatial determinants of abandonment of large-scale arable lands and managed grasslands in Slovakia during the periods of post-socialist transition and European Union accession. *Appl Geogr* 54, 118–128.
- Pfaff, A.S.P., 1999. What Drives Deforestation in the Brazilian Amazon? *J Environ Econ Manage* 37, 26–43.
- Plieninger, T., Schaich, H., Kizos, T., 2010. Land-use legacies in the forest structure of silvopastoral oak woodlands in the Eastern Mediterranean. *Reg Environ Chang* 11, 603–615.
- Plue, J., Meuris, S., Verheyen, K., Hermy, M., 2009. The importance of artefacts of ancient land use on plant communities in Meerdaal forest, Belgium. *Belgian J Bot* 142, 16.
- Polityka ekologiczna państwa w latach 2009-2012 [WWW Document], 2008. URL http://www.mos.gov.pl/g2/big/2009_11/8183a2c86f4d7e2cdf8c3572bdba0bc6.pdf
- Radeloff, V.C., Hammer, R.B., Voss, P.R., Hagen, A.E., Field, D.R., Mladenoff, D.J., 2001. Human Demographic Trends and Landscape Level Forest Management in the Northwest Wisconsin Pine Barrens. *For Sci* 47, 229–241.
- Rhemtulla, J.M., Mladenoff, D.J., 2007. Why history matters in landscape ecology. *Landsc Ecol* 22, 1–3.
- Rhemtulla, J.M., Mladenoff, D.J., Clayton, M.K., 2009. Legacies of historical land use on regional forest composition and structure in Wisconsin, USA (mid-1800s-1930s-2000s). *Ecol Appl* 19, 1061–78.
- Sarukhán, J., Whyte, A., 2005. *Ecosystems and Human Well-Being*. Island Press, Washington DC.
- Schelhaas, M.-J., Nabuurs, G.-J., Schuck, A., 2003. Natural disturbances in the European forests in the 19th and 20th centuries. *Glob Chang Biol* 9, 1620–1633.

- Schiller, H., Miklós, D., Sass, J., 2010. The Danube River and its Basin Physical Characteristics, Water Regime and Water Balance, in: Brilly, M. (Ed.), *Hydrological Processes of the Danube River Basin: Perspectives from the Danubian Countries*. Springer, pp. 25–77.
- Schulze, E.-D., Lange, O.L., Oren, R. (Eds.), 1989. *Forest Decline and Air Pollution. A Study of Spruce (Picea abies) on Acid Soils*. Springer.
- Sitko, I., Troll, M., 2008. Timberline Changes in Relation to Summer Farming in the Western Chornohora (Ukrainian Carpathians). *Mt Res Dev* 28, 263–271.
- Svoboda, M., Janda, P., Nagel, T.A., Fraver, S., Rejzek, J., Bače, R., 2012. Disturbance history of an old-growth sub-alpine *Picea abies* stand in the Bohemian Forest, Czech Republic. *J Veg Sci* 23, 86–97.
- Swetnam, T.W., Allen, C.D., Betancourt, J.L., 1999. Applied Historical Ecology: Using the past to manage for the future. *Ecol Appl* 9, 1189–1206.
- Swinnen, J.F.M., 1999. The political economy of land reform choices in Central and Eastern Europe. *Econ Transit* 7, 637–664.
- Szilassi, P., Jordan, G., van Rompaey, A., Csillag, G., 2006. Impacts of historical land use changes on erosion and agricultural soil properties in the Kali Basin at Lake Balaton, Hungary. *CATENA* 68, 96–108.
- Thompson, J.R., Carpenter, D.N., Cogbill, C. V, Foster, D.R., 2013. Four centuries of change in northeastern United States forests. *PLoS One* 8, e72540.
- Veldkamp, E., Weitz, A.M., Staritsky, I.G., Huising, E.J., 1992. Deforestation trends in the Atlantic Zone of Costa Rica: A case study. *L Degrad Dev* 3, 71–84.
- Verburg, P.H., Berkel, D.B., Doorn, A.M., Eupen, M., Heiligenberg, H. a. R.M., 2009. Trajectories of land use change in Europe: a model-based exploration of rural futures. *Landsc Ecol* 25, 217–232.
- Verburg, P.H., Eck, J.R.R. Van, Nijs, T.C.M. De, Dijst, M.J., Schot, P., 2004. Determinants of land-use change patterns in the Netherlands. *Environ Plan B Plan Des* 31, 125–150.
- Wallin, D.O., Swanson, F.J., Marks, B., 1994. Landscape Pattern Response to Changes in Pattern Generation Rules: Land-Use Legacies in Forestry. *Ecol Appl* 4, 569–580.
- Wendland, K.J., Lewis, D.J., Alix-Garcia, J., Ozdogan, M., Baumann, M., Radeloff, V.C., 2011. Regional- and district-level drivers of timber harvesting in European Russia after the collapse of the Soviet Union. *Glob Environ Chang* 21, 1290–1300.
- Zielonka, T., Holeksa, J., Fleischer, P., Kapusta, P., 2010. A tree-ring reconstruction of wind disturbances in a forest of the Slovakian Tatra Mountains, Western Carpathians. *J Veg Sci* 21, 31–42.

Tables and figures

Table 2.1 Maps and satellite images used for forest cover mapping

Time layer	Data range of maps	Map scale/ resolution	Map source/ description
1860s	1819-1873	1:28.800	Second Austrian Military Survey
1960s	1949-1983	1:50.000 and 1:25.000	Soviet and National Topographic Maps from the Cold War period
1985s	1984-1987	30m	Landsat TM composite
1990s	1988-1992	30m	Landsat TM composite
1995s	1993-1997	30m	Landsat TM composite
2000s	1998-2002	30m	Landsat TM/ ETM+ composite
2005s	2003-2007	30m	Landsat TM/ ETM+ composite
2010s	2008-2012	30m	Landsat TM/ ETM+ composite

Table 2.2 List of predictors used in the forest disturbance models (n=19947), including data sources, measurements in units (Unit), their spatial resolution (SpRes), mean values (Mean), standard deviation (SD), range (Min, Max)

	Variable	Description	Source	Unit	SpRes	Mean	SD	Min	Max
Response	dist_8510	Forest disturbance 1985-2010	Griffiths et al, 2013	Yes/No	30m	factor	N/A	N/A	N/A
Historic land use	FNF1860	forest cover 1860	ref hist maps		vector	factor	N/A	N/A	N/A
Environmental	elev	elevation	Farr et al, 2007	m	90m	693.83	327.22	77.00	1598.00*
	slope	slope	Farr et al, 2007	°	90m	12.79	7.25	0.00	54.09
	temp	Annual Mean Temperature in C*10 from WORLDCLIM	Hijmans et al, 2005	C * 10	~1km	67.86	18.17	6.00	114.00
	precip	Annual Precipitation in mm from WORLDCLIM	Hijmans et al, 2005	mm	~1km	767.21	118.94	524.00	1481.00
	crop_si	Crop suitability index	FAO (GAEZ), 2014	%	~8km	3466.58	2221.02	0.00	10000.00
	grow_ss	Length of growing season	FAO (GAEZ), 2014	days	~8km	205.21	19.93	143.00	253.00
Accessibility	acc_50k	accessibility to nearest 50k inhabitants town, time in minutes	Nelson, 2008	min	~1km	175.24	131.42	1.00	869.00
	dist_city	Euclidean distance to nearest major city in km	ESRI, 2008	km	vector	44.36	19.55	1.10	96.77
	dist_settl	Euclidean distance to nearest settlement in km	EEA, 2013	km	vector	2.97	2.51	0.00	17.87
	dist_road	Euclidean distance to nearest road	CIESIN & ITOS, 2013	km	vector	7.97	6.64	0.00	50.10
	dist_border	Euclidean distance to nearest current border in km		km	vector	52.72	49.64	0.00	213.81
	dist_rail	Euclidean distance to nearest railroad in km	ESRI, 2008	km	vector	13.94	11.76	0.01	70.46
	dist_river	Euclidean distance to nearest main river in km	Vogt et al 2007	km	vector	5.28	3.43	0.00	19.97
Socio-political	cntry	country delineation	ESRI, 2008	N/A	vector	factor	N/A	N/A	N/A
	pop90	population count for year 1990	CIESIN, 2005	pers	~5km	1081.45	1927.00	0.00	38026.50

*The maximum elevation was truncated at 1600m, the average timberline for the Carpathian Mountains (see Methods)

Figure 2.1 Study area in Eastern Europe and forest cover maps for the 1860s, 1960s and 2010s. CZ: Czech Republic, HU: Hungary PL: Poland, RO: Romania, SK: Slovakia, UA: Ukraine




Figure 2.2 Percentage forest disturbance in the Carpathian Region (overall) and by country (RO: Romania, SK: Slovakia, UA: Ukraine, PL: Poland, CZ: Czech Republic, HU: Hungary) in old forests and new forests. Old forests refer to areas that were forested throughout 1860s-1985, new forests to areas not forested in 1860s, but forested in 1960s. The graph indicates consistently more forest disturbance in new forests. Number of observations per country in brackets.




Figure 2.3 Odds of forest disturbance (in %) in new forests (not forested in 1860s) compared to old forests (forested in 1860s), overall and in country models (RO: Romania, SK: Slovakia, UA: Ukraine, PL: Poland, CZ: Czech Republic, HU: Hungary)




Figure 2.4 Proportion of forest disturbance (left y axis) in old forests (black line) and new forests (grey line) in relation to elevation, slope, annual mean temperature, annual precipitation, crop suitability, length of growing season, accessibility, distance to cities, settlements, roads and rivers. The graph represents a data summary of all observed disturbance. The dotted line represents number of forest observations in 1985 (right y axis). Continuous variables were divided in 30 equal interval data bins and the proportion of disturbance as well as the mean number of forest observations was plotted for each bin. For model based results please see Appendix 2.4





Figure 2.5 (a) Forest types by year and country since 1860s, (b) 1860s forest types of disturbed forests between 1985-2010, (c) 1860s forest types of old forests (d) 2010s forest types in new forests, and (e) 2010s forest types of old forests . RO: Romania, SK: Slovakia, UA: Ukraine, CZ: Czech Republic, HU: Hungary. Poland has been excluded from this graph due to scarcity of forest type data. Note: Figures are not for entire country territory but for the study region depicted in Figure 2.1. Forest type distribution in (a) is not identical to Griffiths et al (2013) because the data presented here is restricted to 1960 forest cover.



Appendix

Appendix 2.1 Number of observations for the entire dataset and for each country for each old forests (F) and new forests (NF). Not disturbed = forested in 1960 and never disturbed between 1985-2010, Disturbed = forested in 1960 and disturbed at least once in a five year time step between 1985-2010.

	1860 NF	1860 F
Overall model		19947
not disturbed	2336	14703
disturbed	527	2381
ROMANIA		8218
not disturbed	847	6276
disturbed	184	911
SLOVAKIA		4011
not disturbed	302	3114
disturbed	81	514
UKRAINE		3079
not disturbed	281	2277
disturbed	77	444
POLAND		1955
not disturbed	560	1198
disturbed	81	116
CZECH REPUBLIC		1524
not disturbed	159	1008
disturbed	57	300
HUNGARY		1160
not disturbed	187	830
disturbed	47	96


Appendix 2.2 Model outputs for best selected models: regression coefficients, odds ratio (OR) of disturbance on historic non forest vs forest, and model performance as area under ROC curve (AUC). An odds ratio > 1 indicates higher rate of disturbance on 1860 non-forest areas.

Model	Coefficients	OR	AUC
Overall	-0.401	1.494	0.66
RO	-0.516	1.676	0.682
SK	-0.529	1.698	0.685
UA	-0.289	1.336	0.643
PL	-0.631	1.879	0.784
CZ	-0.186	1.204	0.622
HU	-0.253	1.288	0.698

Appendix 2.3 Number of best performing models ($\Delta\text{AIC} < 3$), of those number of models that include land use legacy as an explanatory variable, ΔAIC of best model including legacies. The second part of the table shows the odds ratios for each selected variable in the exhaustive search

	Overall	RO	SK	UA	PL	CZ	HU
Exhaustive model search							
Number of models $\Delta\text{AIC} < 3$	10	31	65	93	45	154	117
Number of legacy models $\Delta\text{AIC} < 3$	10	31	65	46	45	2	2
ΔAIC of legacy model	0	0	0	0.12	0	2.83	2.7
Odds ratios and coefficient signs							
(Intercept)	5.17	10.6	17.26	22.00	0.002	0.02	853
FNF1860	1.52	1.69	1.69	1.35	1.89	1.20	1.28
elev			1.00	0.99			0.99
slope	0.96	0.97	0.95	0.96	0.93	0.96	0.96
temp	0.98	0.97	0.98		1.07		0.95
precip	1.00				1.00		
crop_si	0.99	0.99	0.99				
grow_ss	0.99	0.99	0.98	0.97	0.97	1.01	0.98
acc_50k	0.99		0.98		0.99		
dist_city	1.00	1.00	1.00	1.01	0.98	0.99	
dist_settl	0.93	0.95	0.92	0.91	0.96		1.22
dist_road	1.01		1.02	1.01			
dist_border		0.99					1.00
dist_rail			0.97	1.01			
dist_river	0.97		0.97		0.92		
pop90					0.99		
centry							
HU	1.11						
SK	0.85						
UA	0.81						
CZ	1.44						
PL	0.55						

Appendix 2.4 Disturbance in old forests (black line) and new forests (grey line) in relation to elevation, slope, annual mean temperature, annual precipitation, crop suitability, length of growing season, accessibility, distance to cities, settlements, roads and rivers. The graph represents regression results once we controlled for all other spatial determinants of change. For a data-based summary please see Figure 2.4



Appendix 2.5. Numbers of observations for forest type data by year and country. Forest type (FT) column indicates the number of points for which forest type information was available. No forest type (NoFT) column indicates the number of data points that were identified as forest but for which forest type information could not be retrieved. (RO: Romania, SK: Slovakia, UA: Ukraine, PL: Poland, CZ: Czech Republic, HU: Hungary)

	1860		1960		1985		2010	
	FT	NoFT	FT	NoFT	FT	NoFT	FT	NoFT
RO	5227	1031	7390	1	7806	412	7995	223
SK	2881	383	4009	0	3948	62	3854	157
UA	1500	358	3024	8	3001	78	2987	92
PL	662	641	1003	0	1919	36	1934	21
CZ	1301	216	1524	0	1481	43	1478	46
HU	926	234	990	0	1091	69	1112	48

Chapter 3. Historical forest management in Romania is imposing strong legacies on contemporary forests and their management

Introduction

Land use dynamics have transformed the Earth's ecosystems to an unprecedented extent (Foley et al., 2005). Long-term forest changes, in particular, have major consequences for ecosystem functioning, carbon storage, climate regulation and biodiversity (DeFries et al., 2004; Newbold et al., 2015). Globally, forest cover loss increased from roughly 7% in 1700 to over 21% in 1990 (Ellis et al., 2013; Goldewijk, 2001) although several countries in Europe and Asia experienced forest transition (Mather, 1998) in late 19th and early 20th century (Meyfroidt and Lambin, 2011) and are currently increasing in forest cover, and carbon sequestration (Erb et al., 2013; Rautiainen et al., 2010). Even though deforestation is declining in some countries (Gold et al., 2006), forest loss due to harvesting and natural disturbances remains high in many areas of the globe (Hansen et al., 2013; Potapov et al., 2014). Forest change is clearly related to socio-economic, political, institutional and environmental drivers (Lambin et al., 2001) but uncertainty about the role of past land uses, also referred to as path dependency, remains a concern for land change assessments. Long term human influence on forests can create legacies that may affect ecosystem functioning, structure and management of ecosystems for centuries (Foster et al., 2003; Munteanu et al., 2015) but the link between past and contemporary land management practices is still poorly understood.

Historical land management decisions affect contemporary landscape patterns across the globe (Foster et al., 2003) and land use legacies can manifest themselves in many aspects of forest ecosystems such as occurrence of disturbance, composition or age patterns. In Eastern Europe, forest disturbance occurs more frequently in areas that were not forested a century ago,

indicating that disturbance patterns are affected by past land management (Munteanu et al., 2015). Similarly, past forest fires and harvests diminish the coniferous forests in the Russian Far East (Cushman and Wallin, 2000) and historically farmed forests in Western Europe show a higher abundance of species that colonize abandoned land, and fewer poor dispersers (Dupouey et al., 2002b; J. Plue et al., 2009). Furthermore, the intensity of historical farming affects forest species composition (Atkinson and Marín-Spiotta, 2015; Plieninger et al., 2010b), indicating that effects of past management may persist for a long time into the future. Finally, age structure can also be a reflection of past land management, because age-patterns established by harvesting can persist for multiple rotation cycles, even under different management practices (Wallin et al., 1994). In summary, this highlights the persistence of land use legacies even after changes in land use type (Munteanu et al., 2015; Thompson et al., 2013) indicating that past land management may constrain forest management for centuries thereafter.

Although forested areas have increased in Europe in the 20th century (Fuchs et al., 2014; Gold et al., 2006; Munteanu et al., 2014), forest disturbance in the past decades is high in Eastern Europe (Griffiths et al., 2014; Hansen et al., 2013) and the forest composition and age structure are altered (Munteanu et al., 2015; Vilén et al., 2012). Contemporary patterns of forest harvesting in Europe vary among countries and have been explained by a suite of factors including site conditions, forest resource availability (Levers et al., 2014), institutional and political context (Baumann et al., 2011; Kuemmerle et al., 2007), ownership structures (Kuemmerle et al., 2009c) and level of protection (Butsic et al., 2015; Knorn et al., 2012b). However, most of these factors can act at different spatial and temporal scales and their effects can change over time, so that the links between past drivers and contemporary change remain unclear.

Eastern Europe represents a particularly interesting natural experiment for studying the relationship between past and contemporary forest change in relation to land tenure, political systems and conservation efforts because the region has a long history of human use (Giosan et al., 2012), very good data records starting as early as the 18th century (Timár et al., 2010) and experienced multiple shifts in institutions, land tenure, and socio-economic pressures both in time and space (Munteanu et al., 2014). Furthermore, current rates of forest harvesting are high (Griffiths et al., 2014) and controversial (Knorn et al., 2012a; Kuemmerle et al., 2009a), but their relationship to past forest management is still largely unexplored.

Our goal here was to examine the connections between historical forest management (as depicted by historical forest cover, species composition, age structure and harvesting) versus contemporary forest patterns in Romania. Specifically, we investigated how past and contemporary forest disturbances (harvesting or natural disturbances which are often followed by salvage logging) are related to ownership structures, forest composition and forest age distribution. We explored possible cause-effect relationships based on forestry census data and remote sensing estimate and focused on lingering effects of historical management in contemporary forests, such as altered forest composition, age structure and shifting disturbance patterns related to forest ownership.

Methods

Study area

We studied forest legacies in Romania (238 381 km²) because the region represents an ideal natural experiment of changing forest management over time. Currently all forests in Romania are managed under the same legislation and consistent forest management plans (Ioras and

Abrudan, 2006), but the region has historically experienced very different forest management regimes because it was split between the Habsburg and Ottoman Empires during the 18th and the 19th century (Munteanu et al., 2015).

Romania is ecologically highly diverse, including parts of five major vegetation ecoregions: Carpathian Montane Coniferous Forests, Pannonian Mixed Forests, Central European Mixed Forests, East European Forest Steppe and Pontic Steppe (European Environment Agency, 2003). The climate is temperate, with continental influences in the northeast and Mediterranean influences in the south. The mean elevation is 330 m and 27% of the country is covered by forest (National Institute of Statistics, 2012). Romania has a total population of 22 million (National Institute of Statistics, 2012), mostly concentrated in urban regions and a per capita GDP of \$13200 (Central Intelligence Agency, 2013), among the lowest in the EU. Historically, land tenure in Romania was split between private owners, churches, institutions and state (Bouriaud, 2008). Historical forest management in Romania was mostly focused on natural regeneration. In the early 1900s, roughly 25% of the Romanian forests were coppice forests, and the remaining 75% were either selectively logged or high forests (i.e., even-aged). Of the high forests, about 10% would be usually clear cut, the rest being managed as shelterwood cuts. Even clearcuts had to retain 50 trees/ha for natural regeneration (Antonescu, 1909).

After the Second World War (WWII) all land was nationalized and managed by the state. Soviet policies heavily influenced forest management leading to widespread clear cuts and planting of fast-growing species. With the collapse of the Soviet Union in 1990, land was partially returned to former private owners following three restitution laws in 1991, 2000 and 2007 (Ioras and Abrudan, 2006). In 2007, Romania joined the European Union, which brought with it new regulations to increase nature conservation (Butsic et al., 2015) and new land management

regulations, such as a requirement for management plans for private forests (Ioras and Abrudan, 2006). However, forests experienced high levels of disturbance after 1990, and particularly after 2000 (Griffiths et al., 2014; Potapov et al., 2014), including the loss of valuable ecosystems and old-growth forests (Knorn et al., 2012a). Contemporary forest management in Romania is largely based on natural regeneration (Schulze et al., 2014). In 2014, only about 1% of the forests were clear-cut and about 12% were shelterwood. About a half of the forests are managed solely by sanitary harvests and about 30% were thinned (Institutul National de Statistica, 2015a).

In addition to the national-level analyses, we conducted three case studies situated in the Eastern Carpathian Mountains to compare historical and contemporary management at a finer spatial resolution. All studies were situated at elevations between 700 – 1100 m and had a total area of 14000 ha (Figure 3.1). The three case study areas are characterized by similar ecological conditions (temperate climate, average yearly temperature around 7 Celsius, average precipitation of 800 mm, dominant soil class of Cambisol, (Institutul de Cercetări și Amenajări Silvice București, 1951; Ministry of Agriculture and Forestry, 1945; Romanian Church Forest Administration, 1926) and hence similar historical forest composition (beech and mixed beech, fir, and spruce forest). Forest management practices and policies were homogeneous since the 1950s until the early 1990s in all three areas because they were under state management, but forests in Humor are currently mostly state managed and in Oituz and Madaras mostly privately managed (Institutul de Cercetări și Amenajări Silvice București, 1951; Ministry of Agriculture and Forestry, 1945; Romanian Church Forest Administration, 1926). Furthermore, our case study areas differed highly in their historical policy, management practices and ownership structures because they were situated on either side of the Ottoman-Austrian-Hungarian border (Table 3.1). This means that the case study areas captured a variety of historical forest management types and

hence provide a great opportunity to examine the role of forest management legacies on current forest composition, structure and disturbance patterns.

Overview of long term forest dynamics

We analyzed long-term forest dynamics in Romania in relation to major socio-economic shifts and ownership changes based on an extensive literature review and national-level statistics. We relied on forest cover statistics about major forest types (coniferous, beech, oak, others) for the years 1924, 1954, 1964, 1980, 1985, 1994, 2006 and 2010 (Directia Centrala de Statistica, 1985, 1980, 1964; Directiunea Statisticeii Generale, 1954; Institutul National de Statistica, 2010, 2006, 1994; Ministerul Agriculturii și Domeniilor, 1924). We used average disturbance data reported in the 1924 forestry statistic for the decade of 1912-1922, in combination with age structure data to reconstruct average disturbance for the decade of 1902-1912. We only extrapolated the age structure for young forest classes because this method will result in estimates with high uncertainties for mature forests. The disturbance value for 1870 is reported in the literature (Nicolau-Barlad, 1944). For the years 1960 to 2014 we calculated disturbed areas based on FAO harvest volume data (FAO (United Nations Food and Agriculture Programme), 2015), which we converted to area estimates (ha) using an average volume/ha value of 400 cubic meters. The conversion factor was chosen based on average dendrometric values for contemporary forests of harvestable age in Romania (i.e. forests 80 years of age or older) (Rusu and Cojinovschi, 2014) and is comparable to timber volumes for clear-cuts in other parts of the world (Masek et al., 2011). We cross validated these estimates with annual disturbance rates reported in remote sensing analysis (Griffiths et al., 2014; Potapov et al., 2014) and found differences of only 1-2% in disturbance of forest areas for the overlapping years. However, our estimation is rather conservative because we assumed that harvest volumes stayed constant over time for the period

1960-2014. Volume density may have increased in recent years (Rautiainen et al., 2010; Vliet et al., 2015) and if this was the case for Romania too, our estimates of disturbance may underestimate the amount of historical harvest.

We analyzed national ownership patterns based on 1924 statistical data at the county level (Ministerul Agriculturii și Domeniilor, 1924) and national statistics for 2010 and 2014 (Curtea de Conturi a României, 2013). We relied on bibliographical sources on ownership data for 1940 and for the socialist period (1948-1990) (Bouriaud and Popa, 2008b; Giurescu, 1981; Nicolau-Barlad, 1944). We compared the proportions of three ownership types in each time periods: public (state owned), institutional, and private.

Since the mid-19th century Romania experienced five major land privatization events concomitant to socio-economic and political shifts such as wars and revolutions. In 1872 serfs were liberated and received land for farming, and in 1921 WWI soldiers received land as war compensation. After WWII all land was nationalized and managed by the state. Following the collapse of the Soviet Union in 1990, three restitution laws, ensured that forest passed back into private ownership in 1991, 2000 and 2007 (Ioras and Abrudan, 2006) (Figure 3.2).

Historical and contemporary spatial data

Our spatial analysis was largely based on forest inventory data for two spatial scales (country level and forest management unit) and focused on two time periods: early 20th century when the study region was under influence of the Habsburg and the Ottoman Empire (hereafter historical) and following the collapse of the Soviet Union and EU accession (hereafter contemporary). In order to analyze forest extent, composition, age classes and disturbances we relied on county-level forest inventory statistics for the historical period (Ministerul Agriculturii și Domeniilor,

1924) and aggregated spatial and statistical data at the county level for the contemporary time period. We digitized forest statistics on age classes and forest composition for 1924 and yearly forest disturbance for the decade 1912-1922. Data was available for the 60 historical counties of Romania according to the 1930 administrative boundaries (Max Planck Institute for Demographic Research and Chair for Geodesy and Geoinformatics, 2015) (Figure 3.1). For the contemporary time period, we integrated four major data sources: two national statistics (Institutul de Cercetări și Amenajări Silvice București, 2015; Institutul National de Statistica, 2015b) and two spatial broad scale data sets, one on forest disturbance (Hansen et al., 2013) and one on forest composition (Brus et al., 2011). We aggregated these data at the county level using administrative boundaries of the 42 Romanian counties of 2014 (Figure 3.1). In order to limit effects of inconsistencies in our data sources and ensure comparability, we used the baseline of national statistics, to which we assigned disturbance rates and species composition from the spatial datasets (Table 3.2).

At the forest management unit level, we obtained forest extent, composition, age and disturbance from forest management plans dated from 1926 to 1945 (Table 3.2). Contemporary forest management plans for the years 2008 to 2014 were available in GIS format and we compared them with digitized historical records to assess shifts in composition, disturbance and age structure.

Forest disturbance

For our analysis, we define disturbance as loss of forest cover due to forest harvest and natural disturbances (which are in Romania most commonly followed by salvage logging). At the national level we relied on historical disturbance data from 1912 to 1922 from forestry statistics

(Ministerul Agriculturii și Domeniilor, 1924). Historical data on forest harvest was reported by foresters in the field and subsequently centralized for each county, and we expect that this data could underestimate the amount of historical harvest. For the contemporary period (2000-2013) we mapped disturbance at county level using remote sensing data (Hansen et al., 2013) complemented with county level statistics for selective and shelterwood logging, because remote sensing data does usually not capture fine-scale disturbances (Kittredge et al., 2003). At the forest management unit level we compared the historical and contemporary occurrence of disturbance based on the forest management plans.

Forest composition

For all of Romania, we compared historical and contemporary extent of four major tree species (beech, oak, fir, and spruce) at the county level using the 1924 and 2014 statistics and reported change as percentage of the total forested area. 1924 data was summarized by 1930 administrative regions. For the contemporary dataset we compiled two data-sources of species distribution: statistical data on the area covered by major forest type (coniferous, deciduous and mixed) at the county level (Institutul National de Statistica, 2015b) and spatial information on the distribution of tree species groups in Europe at 1×1 km (Brus et al., 2011). We calculated percentage of tree species per county and assigned them to major forest types. We finally summarized tree species areas by county in order to obtain more detailed statistics. At the forest management unit level, composition is reported as percentage species in a given stand. For the three case studies, we compared historical and contemporary extent and percentage of species for each forest management unit.

Forest age

Across Romania, statistical data on age class distribution was available to us only at regional level for 2014 (Institutul de Cercetări și Amenajări Silvice București, 2015), and at the country level for 1964 (Directia Centrala de Statistica, 1964) and 1924 (Ministerul Agriculturii și Domeniilor, 1924). We aggregated all data at the national scale and analyzed changes over time. At forest management unit level, we compared shifts in age distribution between the historic and contemporary time periods at the stand level.

Comparison of historical and contemporary data with alternative data sources

Because the reliability of historical forestry statistics is often questionable (Kuemmerle et al., 2011; Schelhaas et al., 2003), we compared the historical datasets used in our analysis with other values reported in the literature and with statistical surveys carried out in the same region by other actors. For the contemporary time period we compared our data with remote sensing estimates and alternative national statistics for Romania (Table 3.3). Overall, we found only small differences between datasets, indicating that datasets used in our analysis captured the status of Romanian forests well. We found the smallest difference between the French forestry statistics from 1900 (Ministere de L'Agriculture du Commerce de L'Industrie ed des Domaines, 1900)a and the Romanian forestry statistic dated 1924, with a 0.4% percentage difference (Ministere de L'Agriculture du Commerce de L'Industrie ed des Domaines, 1900; Ministerul Agriculturii și Domeniilor, 1924) (Table 3.3). The largest difference between datasets occurred in the case of forest disturbance between 1900 and 1924 by 12.5% percent, but this is very likely due to the difference in the reporting year (Table 3.3a). We also checked the correlation of datasets available on contemporary forest cover, disturbance and composition and observed a

maximum correlation of 0.98 for disturbance estimates and a minimum correlation of 0.81 for species distribution (Table 3.3b).

Results

Overview of long term forest dynamics

Historical forest management, in particular past, extensive forest harvest, is strongly reflected in contemporary age structure, composition and disturbance patterns across Romania. Overall, forest area increased in Romania by roughly 308000 ha since 1924, and the country experienced forest transition (i.e., the shift from decreasing to increasing forest area) in the first half of the 20th century. The lowest forest cover occurred sometime between 1920s, when disturbance was particularly high (93000 ha) and 1955 when forest inventory area was at its minimum (5735000 ha). Forest harvest reached its highest point in the late 19th century (with over 100,000 ha being harvested in one year, Figure 3.2 and Figure 3.3). The contemporary Romanian forest inventory reports 6.3 mil ha of forest, which does not include shrub encroachment and forest succession on abandoned lands (estimated at 2.2 mil ha, Hansen et al., 2013). Overall, annual forest disturbance decreased from 1.40 % of the total forest cover in 1924 to 0.71% in 2013.

Forest composition also changed substantially in Romania, with the proportion of deciduous forests decreasing strongly since 1924, when beech accounted for 39% and oak for 22% of the total forest cover. The maximum coniferous cover was reached in Romania in the mid-1980s (31%) (Figure 3.2).

Forest ownership changed drastically during several historic land reforms and post-socialist privatization. Our data indicated that in Romania in 1924, land ownership was divided between private land owners (3298000 ha), state (1556000 ha) and other institutions (1217000 ha), i.e.,

roughly 54, 26 and 20% respectively. Privately owned land decreased by 1940 to 48% of the total forest area. In 1948 all forest was passed into state ownership (Ioras and Abrudan, 2006). Total state ownership lasted until 1991 when following the collapse of the socialism land started being privatized. Post socialist statistics report a shift in ownership to 30% private, 53% state and 17% other institutions, with a higher percentage of publicly owned forests than before WWII (Figure 3.2). In 2014, private forests represented roughly one third of the private forest in 1924. The cross-tabulation of forest disturbance and ownership patterns showed that in 1924, 54% of the forests were privately owned, but as much as 66% of the disturbances occurred in privately owned forests and only 20% in state forests. Spatial information on disturbance by ownership type for 2010 was not available to us.

Forest disturbance

Forest area increased in Romania since 1924 (when it covered 6072000 ha) by 5% and the annual amount of forest harvested (clear cuts and final cuts) between 2000-2013 dropped by 50% (~ 42000 ha/ year) compared to 1912-1922 (~85000 ha/year). Historically, forest harvest was concentrated in the more accessible, lowland areas of Romania, especially in the south and east of the country, where individual counties had annual harvesting rates between 4-6% of their forest cover (Constanta, Ilfov, Vlasca, Olt and Covurlui). Contemporary forest harvesting is concentrated mostly in Northern Carpathians and the northern half of Transylvania (Suceava, Bistrita-Nasaud, Harghita, Covasna, Cluj, Mures, Neamt, Bacau), as well as in the south-east of the country (Calarasi, Ialomita), where forest cover was low to begin with (10% of the county territory). In contrast to overall lower harvesting rates across Romania, in some of the Eastern Carpathian counties, contemporary forest disturbance was higher than historic forest disturbance (Figure 3.3).

At the local level, disturbance decreased in all cases, but most prominently in the case of Humor, where there was almost no disturbance in the period 2000-2010. For the Madaras and Oituz cases, the difference in the amount of harvest was small, but the disturbances were historically clustered in space and more evenly distributed in the contemporary time period. (Figure 3.4)

Forest composition

We found that the total area, proportion and spatial distribution of main tree species changed drastically across Romania since 1924. Forest composition shifted towards higher proportion of coniferous (*Picea* sp, *Pinus* sp, *Larix decidua* and *Pseudotsuga menziesii*) and some deciduous species (*Tilia cordata*, *Populus* sp, *Betula* sp, and *Alnus* sp), which are now more homogeneously distributed in space. Norway spruce increased in area since 1924 (by 6.75%), currently covering an area of 1590000 ha in Romania. Spruce was historically concentrated at higher elevations and in the northern part of the Carpathians, but is now also found at lower elevations. Beech and fir decreased in area (by 14.66% and 1.05% respectively), losing a total of 861000 ha, mostly to spruce plantations. For both species, we found a more spatially homogeneous distribution amongst most counties of Romania: beech declined in the southern Carpathians and the west of Romania and increased slightly in the south of the country. Contemporary oak cover was roughly the same in Romania as in 1924 (ca. 1400000 ha, amounting 22% of the forest cover) but the abundance and spatial distribution shifted greatly from a center of their distribution in southern Transylvania and the western part of the country towards the southern and eastern regions of the country. We recorded highest loss of oak from the historic regions of Alba de Jos, Tarnava Mica and Tarnava Mare where oak comprised between 30-50% of all forests in 1924 to only 10-20% in 2010 (Figure 3.5).

At the local scale, our three case studies confirmed the trends observed for Romania as a whole: a drop in the percentage of fir, beech (and oak in Oituz, where it was present to start with) as well a strong increase in spruce. Overall, forest stands were historically larger and fairly homogeneous in their species composition but became patchier in the contemporary time period. Spruce was more widespread in early 21st century with the exception of Madaras, where forest cover decreased altogether due to contemporary natural disturbances. In the Humor case study, fir area decreased from 1440 ha to 970 ha, being largely replaced by spruce (350 ha) and beech (115ha). Beech experienced a slight increase from 955 ha to 1030 ha (Figure 3.6). In the Oituz case study, the decline of fir, beech and oak was mirrored by an increase in spruce and hornbeam, with generally smaller homogenous stands (Figure 3.6).

Forest age

In contemporary Romania, more forests are even-aged and the area of old forests decreased compared to the historical time period. Age structure data was available only at regional level for 2014, and at the county level for 1924. We complemented this dataset with national level statistics for 1964 and aggregated all data to the national level. Old forests (over 80 years) had a higher percentage in 1924 (25 % of all forests) compared to 2014 (21% of all forests). In 1924 as much as 49% of all forests were in age classes below 40 years old, with a total of 1887000 ha being younger than 20 years old. Overall, we observed an equalization of age structure over time, with roughly 10-17% forest in each age class. Between 1924 and 2014, forests over 100 years declined from 14% to 9% and forests between 80-100 increased by 1% (Figure 3.7).

When cross-tabulating ownership and age structure, we found that historically the largest proportion of forests under 20 years old (61%) was privately owned, whereas old forest (>100

years) where roughly evenly distributed between state, institutional, and private land owners. In 2014, only a small proportion of old forests (17%) was privately owned, and the state owned most of the old forests in Romania, as much as 191000 ha more than in 1924.

At the local scale, our results indicated that forests were historically older compared to the contemporary period, with the exception Madaras, where a long history of spruce plantations led to successive wind disturbances and very young forest. In both Humor and Oituz, we found a high loss of forests in age classes older than 100 years, and an overall tendency of even distribution among age classes. In the case of Humor, contemporary stands were mostly 20-60 years old, whereas in Oituz most stands were 100 years or older. (Figure 3.7).

Discussion

Overview of long term forest dynamics

Our results showed that age structure, composition and disturbance patterns have changed greatly since the early 20th century in Romania and we argue that legacy effects of forest management from nearly a century ago are still greatly reflected in contemporary forests. Forest cover increased in Romania and disturbance is much lower than in early 20th century; but due to intensive management in the past, contemporary forests have a higher percentage of spruce and less beech and oak. We suggest that major shifts in the amount of disturbance and in species composition may be related to changes in governance and land tenure because disturbance peaked around the time of agrarian reforms in the 1920s and post-socialist privatization in the 1990s and 2000s .

Our study captured several major changes in land tenure systems, and we suggest that forest disturbance was closely related to changes in forest ownership. Specifically, our data captured

three land-ownership trends: a) decrease in private land from 1926 to 1948, following the agrarian reform of 1921 (Ioras and Abrudan, 2006; Ministerul Agriculturii și Domeniilor, 1924; Nicolau-Barlad, 1944), b) entirely state-owned land from 1948 to 1989 (Bouriaud and Popa, 2008b; Giurescu, 1981; Nicolau-Barlad, 1944), and c) increase of private land as the result of three privatization laws (Law 18/1991, Law 1/2000 and Law 247/2005) from 1990 to 2014 (Ioras and Abrudan, 2006). These three periods roughly coincide with a decrease, stagnation, and increase in the amount of forest disturbance, suggesting that forest disturbance may be related to changes in land tenure, and specifically to the share of privately owned land. Our results are in line with global forestry literature which indicates that forests with stable ownership have significantly lower rates of harvest (Jin and Sader, 2006), and that harvest rates are higher in private forests (Kittredge et al., 2003). In Romania, missing or unclear regulations and the widespread lack of management plans for privately owned forests may provide a potential explanation for the high rates of contemporary harvesting in private forests that we observed. With 50-75% (roughly 700,000ha) of its private forests lacking forest management plans, Romania is one of only five European countries in which management requirements are not fully consolidated across land ownership forms (Schmithuesen and Hirsch, 2010). Our results suggest that despite of 50 years of Socialism, when all forests were managed by the state (Ioras and Abrudan, 2006) legacies of historical shifts in governance can affect forest ecosystems far into the future, and may be related to the loss of old growth forests and changes in species composition. Similarly, our results highlighted the importance of stable governance and land tenure in maintaining forest area, age structure, composition and harvest rates.

Forest disturbance

Based on our results, forest cover increased in Romania since the 1920s and forest transition, i.e., the shift from a decrease to an increase in forest cover (Mather, 1998; Rudel et al., 2005), occurred approximately in the interwar period, consistent with case-studies in the region (Munteanu et al., 2014). Forest area in the current Romanian territory was as high as 10 million ha until 1860s (Nicolau-Barlad, 1944). By 1900 forest cover decreased in Romania by 3 mil ha – (Giurgiu, 2010a, 2010b) due to agrarian reforms at the end of the 19th century, which granted forested land to serfs for farming (Giurgiu, 2010b; Hitchins, 1994). Harvest rates were very high between 1912 and 1922, lowering the total tree cover to a minimum of 5023000 ha in the mid-1920s due to high timber needs for war purposes. Another agrarian reform in 1921, caused around 1 mil ha of clearings (Florescu, 1937; Giurescu, 1981; Giurgiu, 2010b; Sabau, 1957). Overall our study suggested that changes in regulations and high demand for agricultural products led to a rapid decrease in forest cover until WWII.

Following WWII, and especially after 1975, Soviet policies increased forest cover (Marea Adunare Nationala, 1976) by establishing forest plantations outside the historical range of forests (Munteanu et al., 2015). All forest were managed centrally and harvests were planned for 10-year time intervals, making reported forest harvest relatively constant (Marea Adunare Nationala, 1976). We observed a peak in harvest around 1965, partly due to war reparations paid to Russia in oil and timber (Banu, 2004). Disturbance peaked again in 1982-1985 when Romania was paying off loans to the International Monetary Fund (Ban, 2012). Following the collapse of the Soviet Union, disturbance rates were also high in Romania (Griffiths et al., 2014; Hansen et al., 2013; Knorn et al., 2012b; Potapov et al., 2011, 2014) especially following major privatization laws in 1991, 2000, and 2005 (Ioras and Abrudan, 2006). This finding provided

further evidence on how institutional instability may increase harvesting patterns (Baumann et al., 2011; Dragoi et al., 2011; Prishchepov et al., 2012). However, we highlight that rates of forest harvesting after 1990 were lower than pre-1990, a fact that is missed by most post-socialist studies.

We found higher historical harvest in the Ottoman and Romanian regions than in the Austrian ones, and attributed this to the increase in exports following the Adrianople Peace Treaty and the removal of the Ottoman timber monopoly in 1829 (Cojocaru-Țuiac, 2010). Furthermore, the impact of agrarian reforms was higher in fertile areas than in mountain regions (Giurgiu, 2010a). In Transylvania, counties located closer to Vienna and with less mountains experienced more deforestation. However, the Northeastern Carpathian region was heavily prized for its timber, both by Austrians and Ottomans (Cojocaru-Țuiac, 2010), and this is where we observed widespread forest harvesting. During the post-socialist period, we found a shift in disturbance patterns, where the mountain regions experienced higher disturbance rates, likely due to more abundant forest resources and increased accessibility.

Forest composition

Our results indicated an overall homogenization of the spatial distribution of tree species, with an increase in spruce (especially in Transylvania) and a shift in the spatial distribution of oaks (especially to Moldova and Wallachia). Our results also suggested that historical forest management - different across empire borders - may have increased the abundance of conifers on the Austrian and of oak on the Romanian side of the border. During the Austrian forest management of the 19th century, conifers such as spruce and pine were widely planted for pulp, timber and for erosion control in Transylvania (Popa, 2001). In contrast, historical Ottoman and

later Romanian forest management was centered on the cut and leave method. This meant that entire watersheds would be clear-cut, but at least 50 trees / ha were left standing as seed sources to ensure natural regeneration (Ministry of Agriculture and Forestry, 1945), leading to dominance of beech and oak. In addition, several oak species (*Quercus rubra*, *Q. frainetto*) were planted historically for erosion control and land reclamation in southern and eastern Romania. However, the increasing percentage of oak and beech in lowland regions coincided with the reduction of the species' ranges in central Transylvania. Although Transylvania still hosts some of the most biodiverse oak wooded pastures (Hartel et al., 2013; Öllerer, 2014), their extent very likely declined severely both during Austrian and Socialist rule (Rus, 2014) due to high value of the timber and because they were cleared for grazing or agriculture (Giurescu, 1981; Rus, 2014). Socialist forest management also affected the current ecosystem composition. Between 1948 and 1989 large clearcuts were prescribed to pay off war debts (Banu, 2004) and international loans, followed by extensive spruce plantations both within (Cojocaru-Țuiac, 2010) and outside forest ranges (2 mil ha between 1948 and 1975, (Marea Adunare Nationala, 1976). The area of spruce increased, while that of fir and deciduous species decreased.

Our local case studies provided additional evidence for legacy effects in forest composition. Madaras was mostly deciduous in Austrian military maps of the mid-19th century (Timár et al., 2010) and our data from the early 20th century indicated that spruce plantation occurred in the early 20th century. The area was clear-cut and restocked with spruce several times which may explain the wide-scale wind throws followed by salvage logging which we observed in contemporary management plans. In Humor, the shift from fir to spruce and beech could be a result of spruce plantation which encouraged natural regeneration of beech instead of fir (Damian, 1978; Isciuc, 2010). Finally, in Oituz we found a relatively high proportion of

successional species like hornbeam or pines which are a good indication of the effect of the cut and leave management. Beech and oak decreased here, likely because their regenerative power was smaller than that of successional species.

Forest age

In terms of forest age, Romania has less very young forests (0-20 years) since 1924 but also less forests older than 100 years. A relative equal distribution of age classes is desirable from a management perspective, because it ensures a sustained wood production for timber and pulp (Halbritter and Deegen, 2015). However, old-growth forest have a high natural and conservation value as they provide habitat for a wide range of species, provide ecosystem services and store carbon (Keeton et al., 2010; Wirth et al., 2009) and their loss is unfortunate from a conservation perspective.

Forest management in post WWII Romania aimed to maximize timber production (Banu, 2004; Giurescu, 1981) to pay war debts and economic loans (Ban, 2012; Banu, 2004), and this led to a decrease in old forests, including some of the last old-growth forests of Eastern Europe (Knorn et al., 2012a; Veen et al., 2010). Although consistent with remote-sensing studies indicating the loss of old-growth forests in Romania (Knorn et al., 2012a), our results also highlight that a large proportion of Romanian forests were already managed in the early 20th century despite their old-growth like structure. However, mature secondary vegetation can provide important ecosystem services, have high biodiversity and conservation value (Newbold et al., 2015).

Local scale case studies confirmed the overall loss of old forests, especially in Oituz and Humor, where stands over 120 years old disappeared since 1924, likely as a result of socialist management to maintain equal age classes (Giurescu, 1981). In Madaras, the young forests of

1945 reflect harvests and spruce plantations of the early 20th century, while the contemporary proportion of young stands is likely due to the wind-throws in 1995-1997 (Popa, 2000)

We caution that uncertainty may be introduced in our datasets by elements such as different methods in assembling forest statistics, clear definition of forest disturbance across data sources and ability for clear forest species identification. Overall, we expect that our historic estimates are more accurate in Transylvania and Bucovina, where historic management plans were available (Nicolau-Barlad, 1938; Stinghe, 1939). We expect historic ownership data to be reliable because detailed inventories were required in the course of the agrarian reforms. The historic tree species compositions may include errors depending on surveyors' ability to differentiate between spruce and fir or various oak species.

Conclusions

Our results suggested that contemporary forests were heavily affected by historical forest management and that changes in institutions and ownership patterns may drastically and rapidly affect disturbance patterns and forest composition. We interpret these results to mean that effects of past management and institutional shifts can linger for centuries, and this is important because many regions of the world are currently experiencing drastic changes in their governance and ownership patterns. Such changes may have snowballing effects on forest systems, their functioning and the services they provide for a long time into the future, making a sound understanding of forest legacies important for both conservation and management. In a regional context, Romania harbors some of the last old-growth forests in Europe, which are in decline across the continent. Romania represents also a major source of timber internationally, although harvest rates have decreased. Our results highlight the need to protect the remaining old forests,

which are declining, as well as the need to balance conservation and management goals in the future in order to ensure sustainable forests in Eastern Europe.

References

- Antonescu, P., 1909. Amenajarea Padurilor. *Rev Padur* 1, 305–317.
- Atkinson, E.E., Marín-Spiotta, E., 2015. Land use legacy effects on structure and composition of subtropical dry forests in St. Croix, U.S. Virgin Islands. *For Ecol Manage* 335, 270–280.
- Ban, C., 2012. Sovereign Debt, Austerity, and Regime Change: The Case of Nicolae Ceausescu's Romania. *East Eur Polit Soc* 26, 743–776.
- Banu, F., 2004. Asalt asupra economiei Romaniei de la Solagra la SOVROM (1936-1956) (in Romanian). Nemira, Bucuresti.
- Baumann, M., Kuemmerle, T., Elbakidze, M., Ozdogan, M., Radeloff, V.C., Keuler, N.S., Prishchepov, A. V., Kruhlov, I., Hostert, P., 2011. Patterns and drivers of post-socialist farmland abandonment in Western Ukraine. *Land use policy* 28, 552–562.
- Bouriaud, L., 2008. Proprietatea și dreptul de proprietate asupra pădurilor între reconstituire și re-compunere.
- Bouriaud, O., Popa, I., 2008. Comparative dendroclimatic study of Scots pine, Norway spruce, and silver fir in the Vrancea Range, Eastern Carpathian Mountains. *Trees* 23, 95–106.
- Brus, D.J., Hengeveld, G.M., Walvoort, D.J.J., Goedhart, P.W., Heidema, A.H., Nabuurs, G.J., Gunia, K., 2011. Statistical mapping of tree species over Europe. *Eur J For Res* 131, 145–157.
- Butsic, V., Munteanu, C., Griffiths, P., Knorn, J., Radeloff, V.C., Kuemmerle, T., 2015. The impact of protected areas on forest harvesting in the Carpathian Mountains 1985-2010 in review.
- Central Intelligence Agency, 2013. *The World Factbook 2013-14*. Washington, DC.
- Cojocaru-Țuiac, C., 2010. *Plutăritul pe Bistrița : Incursiune in istorie*. Universitas XXI, Iasi, Romania.
- Curtea de Conturi a Romaniei, 2013. *Sinteza Raportului de audit privind ” Situația patrimonială a fondului forestier din România , în perioada 1990- 2012 ”*.
- Cushman, S.A., Wallin, D.O., 2000. Rates and patterns of landscape change in the Central Sikhote-alin Mountains, Russian Far East. *Landsc Ecol* 15, 643–659.
- Damian, I., 1978. *Împăduriri* . Editura Didactică și Pedagogică, București,.
- DeFries, R.S., Foley, J.A., Asner, G.P., 2004. Land-use choices : balancing human needs and ecosystem function In a nutshell : *Frontiers Ecol* 2, 249–257.
- Directia Centrala de Statistica, 1964. *Anuarul statistic al Republicii Socialiste România*. Editura Științifică, Bucharest.

- Directia Centrala de Statistica, 1980. Anuarul statistic al Republicii Socialiste România. Editura Stiintifica, Bucharest.
- Directia Centrala de Statistica, 1985. Anuarul statistic al Republicii Socialiste Romania. Editura Stiintifica, Bucharest.
- Directiunea Statisticeii Generale, 1954. Anuarul statistic al R.P.R. Editura Stiintifica, Bucharest.
- Dragoi, M., Popa, B., Blujdea, V., 2011. Improving communication among stakeholders through ex-post transactional analysis - case study on Romanian forestry. *For Policy Econ* 13, 16–23.
- Dupouey, J.L., Dambrine, E., Laffite, J.D., Moares, C., 2002. Irreversible Impact of Past Land Use on Forest Soils and Biodiversity. *Ecology* 83, 2978–2984.
- Ellis, E.C., Kaplan, J.O., Fuller, D.Q., Vavrus, S., Klein Goldewijk, K., Verburg, P.H., 2013. Used planet: a global history. *Proc Natl Acad Sci USA* 110, 7978–85.
- Erb, K.-H., Kastner, T., Luyssaert, S., Houghton, R.A., Kuemmerle, T., Olofsson, P., Haberl, H., 2013. Bias in the attribution of forest carbon sinks. *Nat Clim Chang* 3, 854–856.
- European Environment Agency, 2003. DMEER: Digital Map of European Ecological Regions [WWW Document]. URL <http://www.eea.europa.eu/data-and-maps/figures/dmeer-digital-map-of-european-ecological-regions>
- FAO (United Nations Food and Agriculture Programme), 2015. The Global Forest Resources Assessments (FRA) [WWW Document]. URL <http://www.fao.org/forestry/fra/en/> (accessed 1.1.15).
- Florescu, M., 1937. Pericolul pulverizării pădurilor României. *Rev pădurilor* 5.
- Foley, J.A., DeFries, R.S., Asner, G.P., Barford, C., Bonan, G., Carpenter, S.R., Chapin, F.S., Coe, M.T., Daily, G.C., Gibbs, H.K., Helkowski, J.H., Holloway, T., Howard, E.A., Kucharik, C.J., Monfreda, C., Patz, J.A., Prentice, I.C., Ramankutty, N., Snyder, P.K., 2005. Global consequences of land use. *Science* (80-) 309, 570–574.
- Forest Design, 2010. Oituz Forest Management.
- Foster, D.R., Swanson, F.J., Aber, J., Burke, I., Brokaw, N., Tilman, D., Knapp, A., 2003. The importance of land-use legacies to ecology and conservation. *Bioscience* 53, 77–88.
- Fuchs, R., Herold, M., Verburg, P.H., Clevers, J.G.P.W., Eberle, J., 2014. Gross changes in reconstructions of historic land cover/use for Europe between 1900 and 2010. *Glob Chang Biol* 21, 299–313.
- Giosan, L., Coolen, M.J.L., Kaplan, J.O., Constantinescu, S., Filip, F., Filipova-Marinova, M., Kettner, A.J., Thom, N., 2012. Early anthropogenic transformation of the Danube-Black Sea system. *Sci Rep* 2, 582.
- Giurescu, C., 1981. Pădurea în viața și istoria poporului român. RSR Publishing House, Bucuresti.

- Giurgiu, V., 2010a. Declinul suprafeței pădurilor și marginalizarea împăduririlor. *Rev Padur* 2.
- Giurgiu, V., 2010b. Considerații asupra stării pădurilor României - partea I : Declinul suprafeței pădurilor și marginalizarea împăduririlor. *Rev Padur* 2.
- Gold, S., Korotkov, A., Sasse, V., 2006. The development of European forest resources, 1950 to 2000. *For Policy Econ* 8, 183–192.
- Goldewijk, K.K., 2001. Estimating global land use change over the past 300 years: The HYDE Database. *Global Biogeochem Cycles* 15, 417.
- Griffiths, P., Kuemmerle, T., Baumann, M., Radeloff, V.C., Abrudan, I. V., Lieskovský, J., Munteanu, C., Ostapowicz, K., Hostert, P., 2014. Forest disturbances, forest recover, and changes in forest types across the Carpathian ecoregion from 1985 to 2010 based on Landsat image composites. *Remote Sens Environ* 151, 72–88.
- Halbritter, A., Deegen, P., 2015. A combined economic analysis of optimal planting density, thinning and rotation for an even-aged forest stand. *For Policy Econ* 51, 38–46.
- Hansen, M.C., Potapov, P. V, Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A., Thau, D., Stehman, S. V, Goetz, S.J., Loveland, T.R., Kommareddy, A., Egorov, A., Chini, L., Justice, C.O., Townshend, J.R.G., 2013. High-resolution global maps of 21st-century forest cover change. *Science* 342, 850–3.
- Hartel, T., Dorresteyn, I., Klein, C., Máthé, O., Moga, C.I., Öllerer, K., Roellig, M., von Wehrden, H., Fischer, J., 2013. Wood-pastures in a traditional rural region of Eastern Europe: Characteristics, management and status. *Biol Conserv* 166, 267–275.
- Hitchins, K., 1994. *Rumania 1866-1947*. Oxford University Press, Oxford.
- INS, 2013. *Volumul de lemn exploatat în anul 2012 (in Romania)*. Bucuresti.
- Institutul de Cercetări și Amenajări Silvice București, 1951. *Varghis Forest Management*. Ministry of Forestry.
- Institutul de Cercetări și Amenajări Silvice București, 2006. *Varghis Forest Managemnt*.
- Institutul de Cercetări și Amenajări Silvice București, 2008. *Manastirea Humorului Forest Management*.
- Institutul de Cercetări și Amenajări Silvice București, 2015. *Inventarul Forestier National - Evaluarea resurselor forestiere din Romania [WWW Document]*. URL <http://roifn.ro/site/ce-este-ifn/> (accessed 1.1.15).
- Institutul National de Statistica, 1994. *Anuarul statistic al Romaniei*. Editura Stiintifica, Bucharest.
- Institutul National de Statistica, 2006. *Anuarul statistic al Romaniei*. Editura Stiintifica, Bucharest.

- Institutul National de Statistica, 2010. Anuarul statistic al Romaniei. Editura Stiintifica, Bucharest.
- Institutul National de Statistica, 2015a. Suprafata terenurilor din fondul forestier parcursa cu taieri, pe categorii de taieri: Selectati criteriile de interogare [WWW Document]. URL <http://statistici.insse.ro/shop/> (accessed 1.1.15).
- Institutul National de Statistica, 2015b. Statistica Economica- Silvicultura [WWW Document]. URL <http://statistici.insse.ro/shop/> (accessed 1.1.15).
- Ioras, F., Abrudan, I. V., 2006. The Romanian forestry sector: privatisation facts. *Int For Rev* 8, 361–367.
- Isciuc, D., 2010. Cercetari privind instalarea artificiala a bradului (*Abies alba*) pe valea Tarcaului.
- Jin, S., Sader, S.A., 2006. Effects of forest ownership and change on forest harvest rates, types and trends in northern Maine. *For Ecol Manage* 228, 177–186.
- Keeton, W.S., Chernyavskyy, M., Gratzner, G., Main-Knorn, M., Shpylchak, M., Bihun, Y., 2010. Structural characteristics and aboveground biomass of old-growth spruce–fir stands in the eastern Carpathian mountains, Ukraine. *Plant Biosyst* 144, 148–159.
- Kittredge, D.B., Finley, A.O., Foster, D.R., 2003. Timber harvesting as ongoing disturbance in a landscape of diverse ownership. *For Ecol Manage* 180, 425–442.
- Knorn, J., Kuemmerle, T., Radeloff, V.C., Keeton, W.S., Gancz, V., Biriş, I.-A., Svoboda, M., Griffiths, P., Hagatis, A., Hostert, P., 2012a. Continued loss of temperate old-growth forests in the Romanian Carpathians despite an increasing protected area network. *Environ Conserv* 40, 182–193.
- Knorn, J., Kuemmerle, T., Radeloff, V.C., Szabo, A., Mindrescu, M., Keeton, W.S., Abrudan, I. V., Griffiths, P., Gancz, V., Hostert, P., 2012b. Forest restitution and protected area effectiveness in post-socialist Romania. *Biol Conserv* 146, 204–212.
- Kuemmerle, T., Chaskovskyy, O., Knorn, J., Radeloff, V.C., Kruhlov, I., Keeton, W.S., Hostert, P., 2009a. Forest cover change and illegal logging in the Ukrainian Carpathians in the transition period from 1988 to 2007. *Remote Sens Environ* 113, 1194–1207.
- Kuemmerle, T., Hostert, P., Radeloff, V.C., Perzanowski, K., Kruhlov, I., 2007. Post-socialist forest disturbance in the Carpathian border region of Poland, Slovakia, and Ukraine. *Ecol Appl* 17, 1279–95.
- Kuemmerle, T., Kozak, J., Radeloff, V.C., Hostert, P., 2009b. Differences in forest disturbance among land ownership types in Poland during and after socialism. *J Land Use Sci* 4, 73–83.
- Kuemmerle, T., Oloffson, P., Chaskovskyy, O., Baumann, M., Ostapowicz, K., Woodcock, C.E., Houghton, R.A., Hostert, P., Keeton, W.S., Radeloff, V.C., 2011. Post-Soviet farmland

- abandonment, forest recovery, and carbon sequestration in western Ukraine. *Glob Chang Biol* 17, 1335–1349.
- L'Office Central de Statistique Du Royaume Hongrie, 1904. *Annuaire statistique hongrois*. Imprimerie de la Societe Anonyme Athenaeum, Budapest.
- Lambin, E.F., Turner, B.L., Geist, H.J., Agbola, S.B., Angelsen, A., Bruce, J.W., Coomes, O.T., Dirzo, R., Fischer, G., Folke, C., George, P.S., Homewood, K., Imbernon, J., Leemans, R., Li, X., Moran, E.F., Mortimore, M., Ramakrishnan, P.S., Richards, J.F., Skånes, H., Steffen, W., Stone, G.D., Svedin, U., Veldkamp, T.A., Vogel, C., Xu, J., 2001. The causes of land-use and land-cover change: moving beyond the myths. *Glob Environ Chang* 11, 261–269.
- Levers, C., Verkerk, P.J., Müller, D., Verburg, P.H., Butsic, V., Leitão, P.J., Lindner, M., Kuemmerle, T., 2014. Drivers of forest harvesting intensity patterns in Europe. *For Ecol Manage* 315, 160–172.
- Marea Adunare Nationala, 1976. LEGE nr. 2 din 15 aprilie 1976 privind adoptarea “Programului national pentru conservarea si dezvoltarea fondului forestier in perioada 1976-2010.”
- Masek, J.G., Cohen, W.B., Leckie, D., Wulder, M. a., Vargas, R., De Jong, B., Healey, S., Law, B., Birdsey, R., Houghton, R. a., Mildrexler, D., Goward, S., Smith, W.B., 2011. Recent rates of forest harvest and conversion in North America. *J Geophys Res Biogeosciences* 116, 1–22.
- Mather, A.S., 1998. The forest transition : a theoretical basis. *Area* 30, 117–124.
- Max Planck Institute for Demographic Research, Chair for Geodesy and Geoinformatics, U. of R., 2015. MPIDR Population History GIS Collection – Europe.
- Meyfroidt, P., Lambin, E.F., 2011. Global Forest Transition: Prospects for an End to Deforestation. *Annu Rev Environ Resour* 36, 343–371.
- Ministere de L'Agriculture du Commerce de L'Industrie ed des Domaines, 1900. *Notices sur les forêts du royaume de Roumanie*. Imprimerie Dreptatea, Bucharest.
- Ministerul Agriculturii și Domeniilor, 1924. *Statistica padurilor de la 1924*. State Publishing House, Chisinau.
- Ministry of Agriculture and Forestry, 1945. *Grozesti - Harja Forest Management*.
- Munteanu, C., Kuemmerle, T., Boltiziar, M., Butsic, V., Gimmi, U., Kaim, D., Király, G., Konkoly-Gyuró, É., Kozak, J., Lieskovský, J., Mojses, M., Müller, D., Ostafin, K., Ostapowicz, K., Shandra, O., Štych, P., Walker, S., Radeloff, V.C., 2014. Forest and agricultural land change in the Carpathian region—A meta-analysis of long-term patterns and drivers of change. *Land use policy* 38, 685–697.
- Munteanu, C., Kuemmerle, T., Keuler, N.S., Müller, D., Balazs, P., Dobosz, M., Griffiths, P., Halada, L., Kaim, D., Király, G., Konkoly-Gyuró, É., Kozak, J., Lieskovský, J., Ostafin,

- K., Ostapowicz, K., Shandra, O., Radeloff, V.C., 2015. Legacies of 19th century land use shape contemporary forest cover Article. *Glob Environ Chang* 34, 83–94.
- National Institute of Statistics, 2012. Romanian Statistical Yearbook.
- Newbold, T., Hudson, L.N., Hill, S.L.L., Contu, S., Lysenko, I., Senior, R.A., Börger, L., Bennett, D.J., Choimes, A., Collen, B., Day, J., De Palma, A., Díaz, S., Echeverria-Londoño, S., Edgar, M.J., Feldman, A., Garon, M., Harrison, M.L.K., Alhusseini, T., Ingram, D.J., Itescu, Y., Kattge, J., Kemp, V., Kirkpatrick, L., Kleyer, M., Correia, D.L.P., Martin, C.D., Meiri, S., Novosolov, M., Pan, Y., Phillips, H.R.P., Purves, D.W., Robinson, A., Simpson, J., Tuck, S.L., Weiher, E., White, H.J., Ewers, R.M., Mace, G.M., Scharlemann, J.P.W., Purvis, A., 2015. Global effects of land use on local terrestrial biodiversity. *Nature* 520, 45–50.
- Nicolau-Barlad, G., 1938. Die Photogrammetrie im Forstwesen dargestellt an der Waldwirtschaft der Karpathen- und Balkanländer. Wichmann Verl. Berlin, Berlin.
- Nicolau-Barlad, G., 1944. Problema planurilor topografice. *Ann For Res*.
- Öllerer, K., 2014. The ground vegetation management of wood-pastures in Romania - Insights in the past for conservation management in the future. *Appl Ecol Environ Res* 12, 549–562.
- Plieninger, T., Schaich, H., Kizos, T., 2010. Land-use legacies in the forest structure of silvopastoral oak woodlands in the Eastern Mediterranean. *Reg Environ Chang* 11, 603–615.
- Plue, J., Dupouey, J.-L., Verheyen, K., Hermy, M., 2009. Forest seed banks along an intensity gradient of ancient agriculture. *Seed Sci Res* 19, 103.
- Popa, I., 2000. Doborâturi Produse De Vânt - Factor De Risc În Ecosistemele. *Analele ICAS* 48, 3–28.
- Popa, I., 2001. Analiza dendroecologică a regimului perturbărilor în pădurile din nordul Carpaților Orientali 17–30.
- Potapov, P., Turubanova, S., Hansen, M.C., 2011. Regional-scale boreal forest cover and change mapping using Landsat data composites for European Russia. *Remote Sens Environ* 115, 548–561.
- Potapov, P.V., Turubanova, S. a., Tyukavina, a., Krylov, a. M., McCarty, J.L., Radeloff, V.C., Hansen, M.C., 2014. Eastern Europe's forest cover dynamics from 1985 to 2012 quantified from the full Landsat archive. *Remote Sens Environ* 159, 28–43.
- Prishchepov, A. V., Radeloff, V.C., Baumann, M., Kuemmerle, T., Müller, D., 2012. Effects of institutional changes on land use: agricultural land abandonment during the transition from state-command to market-driven economies in post-Soviet Eastern Europe. *Environ Res Lett* 7, 024021.
- Rautiainen, A., Saikku, L., Kauppi, P.E., 2010. Carbon gains and recovery from degradation of forest biomass in European Union during 1990–2005. *For Ecol Manage* 259, 1232–1238.

- Romanian Church Forest Administration, 1926. Manastirea Humorului Forest Management.
- Rudel, T.K., Coomes, O.T., Moran, E.F., Achard, F., Angelsen, A., Xu, J., Lambin, E.F., 2005. Forest transitions: towards a global understanding of land use change. *Glob Environ Chang* 15, 23–31.
- Rus, D.-I., 2014. Conflict and consensus regarding resource policies in a borderland of the Habsburg Empire: Perspectives on and perceptions of forests and the environment in Transylvania at the end of the eighteenth century and in the nineteenth century. *Ekonomika i Ekohist* X, 233–245.
- Rusu, E., Cojinoșchi, A.-D., 2014. Structural Dynamics of Romanian Forests after 1990. *Present Environ Sustain Dev* 8.
- Sabau, V., 1957. Evoluția economiei forestiere în România. Editura Societății „Progresul silvic”, București.
- Schelhaas, M.-J., Nabuurs, G.-J., Schuck, A., 2003. Natural disturbances in the European forests in the 19th and 20th centuries. *Glob Chang Biol* 9, 1620–1633.
- Schmithuesen, F., Hirsch, F., 2010. Private Forest Ownership in Europe.
- Schulze, E.D., Bouriaud, L., Bussler, H., Gossner, M., Walentowski, H., Hessenmöller, D., Bouriaud, O., Gadow, K. V., 2014. Forest management and biodiversity. *Web Ecol* 14, 3–10.
- Stinghe, V.N., 1939. Amenajarea Padurilor. Editura Societății „Progresul silvic”, Bucharest.
- Thompson, J.R., Carpenter, D.N., Cogbill, C. V, Foster, D.R., 2013. Four centuries of change in northeastern United States forests. *PLoS One* 8, e72540.
- Timár, G., Biszak, S., Székely, B., Molnár, G., 2010. Digitized Maps of the Habsburg Military Surveys – Overview of the Project of ARCANUM Ltd. (Hungary), in: Jobst, M. (Ed.), *Preservation in Digital Cartography, Lecture Notes in Geoinformation and Cartography*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 273–283.
- Veen, P., Fanta, J., Raev, I., Biriș, I.-A., Smidt, J., Maes, B., 2010. Virgin forests in Romania and Bulgaria: results of two national inventory projects and their implications for protection. *Biodivers Conserv* 19, 1805–1819.
- Vilén, T., Gunia, K., Verkerk, P.J., Seidl, R., Schelhaas, M.-J., Lindner, M., Bellassen, V., 2012. Reconstructed forest age structure in Europe 1950–2010. *For Ecol Manage* 286, 203–218.
- Vliet, J. Van, Groot, H.L.F. De, Rietveld, P., Verburg, P.H., 2015. Landscape and Urban Planning Manifestations and underlying drivers of agricultural land use change in Europe. *Landsc Urban Plan* 133, 24–36.
- Wallin, D.O., Swanson, F.J., Marks, B., 1994. Landscape Pattern Response to Changes in Pattern Generation Rules: Land-Use Legacies in Forestry. *Ecol Appl* 4, 569–580.

Wirth, C., Messier, C., Bergeron, Y., Frank, D., Fankhänel, A., 2009. Old-Growth Forest Definitions: a Pragmatic View, in: Wirth, C., Gleixner, G., Heimann, M. (Eds.), Old-Growth Forests, Ecological Studies. Springer, Berlin, Heidelberg, pp. 11–33.

Figures and tables

Table 3.1 The three case studies, including name, area, historic region, type of management and forest ownership

Case study	Area (ha)	Historic region	Historical management	Historical ownership	Contemporary ownership
Humor	3500	Bucovina	Austrian	Church	State
Oituz	9000	Moldova	Romanian	Private	Private
Madaras	1400	Transylvania	Hungarian	Private	Private

Table 3.2 Data sources for forest extent, composition, age classes and disturbances for three time periods and at two spatial scales

	Historical (1924-1945)	Contemporary (2000-2014)
Spatial scale: Romania, at county level		
Disturbance occurrence	(Ministerul Agriculturii și Domeniilor, 1924)	(Hansen et al., 2013; Institutul National de Statistica, 2015b)
Forest ownership	(Ministerul Agriculturii și Domeniilor, 1924)	(Institutul de Cercetări și Amenajări Silvice București, 2015)
Age class distribution	(Ministerul Agriculturii și Domeniilor, 1924)(Nicolau-Barlad, 1938)	(Institutul de Cercetări și Amenajări Silvice București, 2015)
Species composition	(Ministerul Agriculturii și Domeniilor, 1924)	(Brus et al., 2011; Institutul de Cercetări și Amenajări Silvice București, 2015)
Spatial scale: Forest management unit: Humor, Oituz, Madaras		
Disturbance occurrence	(Institutul de Cercetări și Amenajări Silvice București, 1951; Ministry of Agriculture and Forestry, 1945; Romanian Church Forest Administration, 1926)	(Forest Design, 2010; Institutul de Cercetări și Amenajări Silvice București, 2008, 2006)
Age class distribution	(Institutul de Cercetări și Amenajări Silvice București, 1951; Ministry of Agriculture and Forestry, 1945; Romanian Church Forest Administration, 1926)	(Forest Design, 2010; Institutul de Cercetări și Amenajări Silvice București, 2008, 2006)
Species composition	(Institutul de Cercetări și Amenajări Silvice București, 1951; Ministry of Agriculture and Forestry, 1945; Romanian Church Forest Administration, 1926)	(Forest Design, 2010; Institutul de Cercetări și Amenajări Silvice București, 2008, 2006)

Table 3.3 Data sets used in our analysis and comparison to values from other sources such as forestry literature, statistical yearbooks and remote sensing estimates for historical (a) and contemporary (b) time periods

(a) Historical				
Compared content	Value used in Analysis	Cross-reference in literature	Extent of comparison	Difference in %
Percentage of forest in Romania	25.472 % (Ministerul Agriculturii și Domeniilor, 1924)	25.875 % (L'Office Central de Statistique Du Royaume Hongrie, 1904; Ministere de L'Agriculture du Commerce de L'Industrie ed des Domaines, 1900)	Country level	-0.403%
Forest species composition 1900	Coniferous – 25% Oaks – 23 % Deciduous – 52% (Ministerul Agriculturii și Domeniilor, 1924)	Coniferous – 21% Oaks – 26 % Deciduous – 53% (L'Office Central de Statistique Du Royaume Hongrie, 1904; Ministere de L'Agriculture du Commerce de L'Industrie ed des Domaines, 1900)	Country level	Conif +4% Oaks – 3 % Decid – 1%
Forest disturbance (ha)	524 698 (Ministerul Agriculturii și Domeniilor, 1924)	590 327 (Ministere de L'Agriculture du Commerce de L'Industrie ed des Domaines, 1900)	Valahia and Moldavia State and Communal Forest	-12.5%
Local Oituz Forest area (ha)	9008 (Forest Design, 2010)	9275 (Ministerul Agriculturii și Domeniilor, 1924)	forest management unit	-2.96%

(b) Contemporary

	Data used in analysis	Cross-checked with data	Extent of comparison	R-squared
Percentage of county covered by forest	(Institutul National de Statistica, 2015b)	(Hansen et al., 2013)	N=42 counties	0.97
Area covered by major species	(Brus et al., 2011; Institutul National de Statistica, 2015b)	(Institutul de Cercetări și Amenajări Silvice București, 2015)	N=15 (5 species, 3 regions)	0.81
Disturbance area	(Hansen et al., 2013; INS, 2013)	(Griffiths et al., 2014; INS, 2013; Potapov et al., 2014)	N=28 counties 91% of total forest cover	0.98

Figure 3.1 Location of study area in Europe, imperial boundaries from 1900s, the location of 3 cases studies in the Carpathian Mountains and county borders for 1924 and 2015




Figure 3.2 Overview of the evolution of forest cover, species composition, disturbance and ownership patterns in Romania between 1870s and 2010, in the context of major land tenure changes




Figure 3.3. Percentage forest cover and forest disturbance in Romania historically (n= 58 counties) and contemporary (n=42 counties)




Figure 3.4 Historical and contemporary forest disturbance at forest stand level in three case studies: Humor (3500ha), Oituz (9000ha) and Madaras (1500ha)




Figure 3.5 Percentage of major coniferous species (*Picea abies*, *Abies alba*) and major deciduous species (*Fagus sylvatica*, *Quercus* sp) within forest cover of Romanian regions in 1924 (n= 58 regions) and in 2014 (n=42 regions)




Figure 3.6 Species composition at local level in Humor (3500ha), Oituz (9000ha) and Madaras (1500ha) at the beginning of the 20th century and the beginning of the 21st century. Stands with species cover higher than 50% are represented in the graphic.





Figure 3.7 Forest age distribution in Romania in at (a) national level for the years 1924, 1964 and 2014 and at local case study level in (b) Oituz (9000ha) between 1930 and 2013, (c) Humor (3500ha) between 1926 and 2006 and (d) Madaras (1500ha) between 1945 and 2006.



Chapter 4. Land use legacies of historic political regimes for contemporary land abandonment

Introduction

Contemporary global environmental change reflects centuries of socio-ecological interactions. One of the major components of global change is land use change (Fuchs et al., 2014; Hurtt et al., 2006; Schelhaas et al., 2003), which is driven by complex direct and indirect driving forces such as environmental conditions, accessibility and market forces (Lambin and Meyfroidt, 2011; Meyfroidt, 2015; Meyfroidt et al., 2013). In addition to these proximate and underlying drivers, which have immediate effects, historical land uses can affect ecosystem structure and functioning for centuries into the future (Foster et al., 2003; Perring et al., 2016; Plue et al., 2009), yet the effect of land use legacies on environmental change remains poorly understood (Perring et al., 2016).

Worldwide, many landscapes bear marks of historical land use (DeFries et al., 2004; Foley et al., 2011). From an ecological perspective, land use legacies affect soils, water content, seed source, vegetation composition, and species establishment and dispersal (Brudvig et al., 2013; Ficetola et al., 2010; Foster et al., 2003), and include, for example, the chemical and biological alterations imposed on soils by past agricultural activity, as far back as Roman Empire agriculture (Plue et al., 2008). The reason why such legacies can persist for centuries is that changes in the soil structure favor the establishment of non-native species (Ficetola et al., 2010) or constrain vegetation succession and productivity (Foster et al., 2003) due to changes in the seed bank, seed dispersal, or soil nutrients (Plue et al., 2009, 2008). Agricultural legacies also affect beta-diversity in forest understory suggesting that the understanding of the interactions between current land use activities and past land use decisions is essential for understanding ecosystem

functioning (Mattingly et al., 2014). Indeed, past land use decisions are often affecting rates of biodiversity loss with a time-lag, contributing to extinction debt (Dullinger et al., 2013; Essl et al., 2015).

From a land use perspective, land use patterns and change are path-dependent processes (Lambin and Geist, 2006; Meyfroidt, 2015) that are at least partly driven by historical land uses (Munteanu et al., 2015a). Path dependencies have been clearly shown for urban dynamics (Lambin and Geist, 2006; Seto et al., 2012), but historical land use pattern may affect all types of land use in the future (Meyfroidt, 2015; Verburg et al., 2004). For instance, whether areas that were historically non-forested had a 50% higher chance of contemporary harvests and natural disturbances compared to areas forested historically (Munteanu et al., 2015a). In swidden-fallow systems, path dependence shapes the farmers land use choices: insufficient land holdings induce low productivity land use traps for poor farmers (Coomes et al., 2011) and in Tanzania agricultural intensification is a path-dependent and self-reinforcing leading to further intensification (Börjeson, 2007). However, while there is some local empirical evidence of path dependency, effects of land use legacies on recent land change have rarely been quantified.

Land use theory explains land use choices and subsequent land changes based on factors that relate environmental suitability and other spatial determinants of land change to the broader social, political, economic, technological, and cultural context (Geist and Lambin, 2002; Meyfroidt, 2015). Immediate land use choices are most often made based on spatial determinants such as environmental suitability or accessibility (Müller et al., 2013; Nagendra et al., 2003). For example, from a land rent perspective, agricultural land with favorable environmental conditions will be preferentially farmed (Ricardo, 1821), and subsequent to land use specialization, marginal land will be abandoned (Müller et al., 2013). If indeed path dependencies and land use

history do affect recent rate of change, then even after accounting for the spatial determinants, legacies will remain important predictors of change.

The effect of land use legacies on contemporary land change will be most clear during times of rampant land change. Most often, rampant land change is caused by major shifts in social, political, economic, technological, or cultural factors that in turn affect land use decisions (Geist and Lambin, 2002; Meyfroidt, 2015). For instance, Stalins' Great Plan for the Transformation of nature led to forceful reforestation to protect farmland in Southern Russia's steppes, but also caused later desertification (Brain, 2010). Similarly, agricultural expansion during Soviet Virgin Lands Campaign in Northern Kazakhstan still affects land abandonment today (Kraemer et al., 2015). In Socialist Romania, war repayments to the Soviet Union led to institutionalized overexploitation of forests, causing a major shift in species composition and age structure (Munteanu et al., 2016). Last but not least, the transition to market economies of former Soviet states led to widespread land abandonment across Europe and Asia (Alcantara et al., 2013; Prishchepov et al., 2012). Eastern Europe has a long land use history and experienced several major shifts in political and institutional regimes, such as the rise and fall of, first, the Habsburg Empire, and, second, of Socialism. These political changes caused major changes in land management and affected land use patterns over a relatively short period of time, providing a great 'natural experiment' for the study of legacies.

Agricultural abandonment is widespread today throughout both temperate and tropical biomes (Munroe et al., 2013), and well suited to study legacy effects, because post-agricultural landscapes are likely to bear the marks of historical land use decisions for many years (Plieninger, 2014; Plieninger et al., 2010). There are several potential processes via which historical land uses may affect abandonment today. For example, the removal of historic

subsidies or changes in the policy context can lead to rapid abandonment once those incentives are removed (Brain, 2010; Jepsen et al., 2015; Kraemer et al., 2015). Environmental conditions, in combination with intensive agricultural practices may cause an alteration of soil suitability for agriculture (Matteucci et al., 2016) potentially leading to more abandonment than if the land was less intensively farmed. In Europe alone, an area the size of Spain was abandoned since 2001 (Estel et al., 2015) and the drivers of land abandonment have been widely studied (Estel et al., 2015; MacDonald et al., 2000; Prishchepov et al., 2013; Rey Benayas, 2007) and include environmental, financial and socio-economic factors (McConnell and Keys, 2005). However, it is unclear if land use legacies also determine contemporary land abandonment. Much of the European land abandonment in past decades was concentrated in the former Eastern Bloc (Griffiths et al., 2013; Munteanu et al., 2014) following political and economic shock of the collapse of the Soviet Union (Baumann et al., 2011; Estel et al., 2015; Munteanu et al., 2014), making this region the perfect case study for investigating the long term effect of land use legacies on contemporary abandonment.

Our overarching goal was to assess the role of long-term land-use legacies on contemporary agricultural land use, using the Carpathian region as our study site. Here, we define land-use legacies as the effects of historical land uses on contemporary agricultural abandonment, especially the transition between tilled annual or perennial crops to any other land cover type. Our specific objectives were to quantify:

- (1) How agricultural land use changed in the Carpathians since 1860s and to what extent agro-ecological conditions explained agricultural land use patterns during each of the major political regimes?

- (2) If there were land use legacies once agro-ecological variation was controlled for and whether their effect diminished with time?
- (3) Whether legacies of different historical regimes differed?

Methods

Study area

We studied an area of approximately 265,000 km² covering the Carpathians and adjacent parts of the Pannonian Plains, with elevations ranging between 400 and 2,500 m. The study region includes all of Slovakia, large parts of Hungary and Romania, the mountainous parts of Poland and Ukraine, and the historic province of Moravia in the Czech Republic (Figure 4.2).

Temporally, we captured agricultural land use at six points in time: during the Habsburg era (1860s), Interwar era (1930s, socialist era (1960s, 1985) and during post-socialism (2000, 2010) (Table 4.1).

The contemporary land cover pattern in the study region consists of a mosaic of agricultural fields, grasslands, and forests at higher elevation, and predominantly agricultural fields and grasslands at lower elevations in the Pannonian plain (Kozak et al., 2013b; Munteanu et al., 2015b). Particularly in the lowlands of Hungary, fields are sometimes intermixed with forest plantations (Munteanu et al., 2014). Overall, agricultural land covers around 23% of the Carpathian region (Griffiths et al., 2013) (Figure 4.2). The main food crops are cereals (wheat, corn, barley) and legumes (potatoes, sugar beets), with energy crops such as rapeseed occupying a smaller proportion of the arable land (Griffiths et al., 2013).

Historical land ownership in the Habsburg Empire was highly concentrated, with noble families owning large tracts of land and a smaller share of the land being owned by institutions,

communities and small landholders (Berger, 2006). During socialism, the majority of the land was under collective state management, with the exception of Poland and some isolated mountainous areas, where private ownership persisted (Kozak et al., 2013a; Lerman et al., 2004). Following land reorganization and restitution after the collapse of the Soviet Union, land ownership in the Carpathian countries has been split between state and private and the patterns vary from small subsistence family farms to large private landholdings (Griffiths et al., 2013).

Historically, agricultural expansion was widespread during the time of the Habsburg Empire, partly due to Trade Unions which removed agricultural tariffs for the Eastern provinces of the Empire (Alix-Garcia and Walker, 2016). The socialist era was characterized by pressure to increase agricultural production, particularly in Romania and Ukraine. However, despite policies driving agricultural expansion and intensification, agricultural abandonment was already during socialism widespread in mountain regions due to the spatial reorganization of agriculture (Jepsen et al., 2015; Munteanu et al., 2014). After the collapse of the Soviet Union, abandonment intensified and has been the prevalent land change process in the Carpathian region.

Approximately 24% of the total cropland in 1985 has been abandoned until 2000 and another 9% until 2010, largely due to socio-demographic and political factors (Griffiths et al., 2013; Munteanu et al., 2014).

Datasets

We reconstructed agricultural land use for six points in time: 1860s, 1930s, 1960s, 1985, 2000 and 2010. Here, we define agricultural land as tilled areas used for annual or perennial crops, and delineated as such in historic maps or in remote sensing data (this definition does not include managed or unmanaged pastures and grasslands) (Table 4.1). We classified land use based on a

regular 2x2 km point sampling grid according to the 2007 INSPIRE directive (Infrastructure for Spatial Information in the European Community) and LUCAS (Land Use and Cover Area frame Survey, (Gallego and Delincé', 2010). For the years 1860, 1930 and 1960, we extracted agricultural land use from historic topographic maps, and for the years 1985, 2000 and 2010 from 30-m resolution Landsat TM/ETM+ image composites that were automatically classified with an accuracy >80% (Griffiths et al., 2013) (Table 4.2).

Our study area included a total number of 70,947 points. For the years 1860, 1930 and 1960 we assigned binary land use classes (agriculture or non-agriculture) to each point. To ensure consistency in point location across map sets, we employed a back-dating approach in which the location of the digitized point was verified in the subsequent time layers in relation to nearby landmarks (Kaim et al., 2014). This approach was employed for all points located in Slovakia, Czech Republic, and Poland. Due to low accuracy of historic maps in parts of Romania and Hungary, for a total of 3,409 points, we could not clearly distinguish agricultural from grassland use of the land for the 1860s time layer. For these points, we assigned the land use class according to the subsequent map dataset from 1930s. In order to check for potential errors, we ran the subsequent analysis with and without these points, but our results did not change substantially, so we retained the points in the analysis. For the years 1985, 2000 and 2010, we automatically extracted the agricultural use class from Landsat image classifications (Griffiths et al., 2013).

Based on the binary classifications (agriculture vs. non-agriculture), we mapped agricultural abandonment for the socialist and the post-socialist eras, using two 25 years' time period for comparability: socialist abandonment (1960-1985) and post-socialist abandonment (1985-2010) (Table 4.1). For the socialist abandonment we analyzed a total of 20,501 points that were used as

agriculture in 1960 and either were abandoned or not by 1985. To map post-socialist abandonment (between 1985 and 2010), we analyze a total of 13419 points, all of which were agriculture in 1985 and that were either abandoned or not by 2000 or by 2010.

Data analysis

Agricultural dynamics and agro-ecological conditions

In order to address our first objective of quantifying the agricultural change its relation to agro-ecological conditions, we compared percentage of agricultural cover over the six time periods for which data was available (1860s, 1930s, 1960s, 1985, 2000, and 2010). We mapped change trajectories between binary agriculture vs. non-agriculture classes. In order to understand the spatial determinants of agricultural dynamics in different time periods, we selected from the sample (70,947 points) only those points that were (a) used for agriculture during the Habsburg era (i.e., 1860s, 31,106 points), (b) converted to agriculture from other land uses during the socialist era (i.e., 1960s, 6,488 points) or (c) used for agriculture during the post-socialist era (i.e., 2010, 15,722 points). Because we were interested to what extent land use decisions during these distinct political and socio-economic time-periods were based on agro-ecological conditions, we modeled the occurrence of agricultural land as a function of seven agro-ecological variables: elevation, slope, distance to nearest river, average annual temperature, average annual precipitation, crop suitability index and length of the growing season using multiple logistic regression models (Hosmer et al., 2013). We checked model performance using receiver operating curves (ROC, (Freeman and Moisen, 2008)) and calculated the area under the ROC curve (AUC) in order to assess how well agro-ecological variables explain the farming choices in the three periods (Hosmer et al., 2013). We also mapped socialist (20,501 points) and post-

socialist abandonment (13,419 points) to better understand the spatial distribution of abandonment.

Persistence of land use legacies

To address our second objective regarding the persistence of land use legacies over time, we calculated and compared the effect of Habsburg land use legacies on socialist (1960-1985) and post-socialist abandonment (1985-2010). We fitted multiple logistic regression models (Hosmer and Lemeshow, 1980) that controlled for agro-ecological, accessibility and socio-political variation, to explain abandonment in relation to the land use legacy of the Habsburg era (1860s) (Table 4.1). Our models included a total sample of 20,501 points for the socialist abandonment models and a total 13,419 points for post-socialist abandonment (Appendix 2.1).

In addition to the agro-ecological variables used in Objective 1 (6 variables), we controlled for accessibility to markets and ease of transport (7 variables), and socio-political variation (2 variables) (Table 4.3), because data on accessibility and socio-political variation was available for recent time periods. We estimated the effect of Habsburg legacy (Tab 1) on socialist and post-socialist abandonment by relying on the odds ratio, which represents the exponential values of the model coefficients (Hosmer et al., 2013). For each of the two abandonment time periods we fitted an overall model for the Carpathian region and six country-specific models (Müller et al., 2009), in order to capture socio-economic and institutional diversity of the region. In total, we thus fitted fourteen models, seven for each abandonment period (Appendix 2.1).

We performed best-subsets variable selection using an exhaustive search (Hosmer et al., 2013) based on the Akaike Information Criterion (AIC). To ensure model parsimony, we restricted the maximum number of variables accepted in the model to six (for the country models) and to seven

for the overall model (including a country dummy). We always retained the best performing model, and in cases where the best performing model did not include the land use legacy (4 of 14 models), we refitted the best performing model adding the land use legacies, because we were interested in estimating and comparing the effect of past agricultural cover on current abandonment across countries.

We calculated the relative rates of agricultural abandonment in areas that were not farmed in 1860s (i.e. late Habsburg) comparing to areas that were farmed then (i.e. early Habsburg), based on the odds ratio. The odds ratio represents the exponential value of the model coefficients in logistic regression (Hosmer et al., 2013). We transformed the odds ratio to percentage points, where values higher than 0 indicated how much more likely abandonment is in areas that were not farmed historically, compared to areas farmed then, hence quantifying the land use legacy effect. For the remaining variables, we interpreted the sign of model coefficients, in order to understand how agro-ecological conditions and accessibility influenced agricultural abandonment. We did not calculate significance levels or confidence intervals in our analysis because our data represent a full census of historical and recent land cover and because our estimate of the effect that we observed is independent of sample size (Lohr, 2010; Munteanu et al., 2015a). We checked model performance using receiver operating curves (ROC, (Freeman and Moisen, 2008)) and evaluated model utility by calculating the area under the ROC curve (AUC). AUC values varied between 0.79 for the overall socialist abandonment model and 0.82 for the overall post-socialist abandonment model.

Strength of land use legacies

In order to assess if the strength of the legacies changed with political regimes, we compared the effect of legacies from three historic time periods (Habsburg era, Interwar era, Socialist era) for post-socialist agricultural abandonment (1985-2010). We modelled post-socialist abandonment using multiple logistic regression models (Hosmer and Lemeshow, 1980) that controlled for agro-ecological, accessibility, and socio-political variation (same 15 variables as in Objective 2, Table 4.3). We applied same model selection criteria as in Objective 2, retained the best-performing models and in cases where the best-performing model did not include the land use legacy (8 of 21 models), we refitted the best performing model including the land use legacies, because we were interested in estimating and comparing the effect of past land use on current abandonment. We interpreted the odds ratio to estimate the legacy effect on post-socialist abandonment. The legacy of the Habsburg era (hereafter Habsburg legacy, Table 4.1) captures differences in abandonment between land already farmed prior to 1860s (early Habsburg) compared to land farmed after 1860s (late Habsburg). The legacy of the socialist era (hereafter socialist legacy, Table 4.1) captures differences in abandonment between land expanded for agriculture in the Socialist era prior to 1960s (early socialist), compared to after 1960s (late socialist). Finally, the legacy of the Interwar era (hereafter Habsburg vs. Socialist legacy, Table 4.1) captures differences between land farmed during the Habsburg era and land expanded during the Socialist era. Given that the Interwar period captured major shifts in political regimes, and that agricultural cover was mostly stable during this period (Munteanu et al., 2014), we used this period to assess differences in abandonment depending on agricultural use during two major political regimes: Habsburg and socialist (Table 4.1).

In total, we fitted twenty one models (one overall model and six country-specific models for each of the three periods). Overall sample sizes varied between 3,424 for the socialist legacy model and 16,843 for Habsburg vs. socialist model (Appendix 4.2). We evaluated model performance and utility using ROC statistics (Freeman and Moisen, 2008). For overall models, AUC values ranged from 0.80 (socialist legacy) to 0.83 (Habsburg vs. Socialist legacy). For the country models, AUC was lowest for Ukraine in the Habsburg vs. Socialist model (AUC= 0.64) and highest for Poland in the Socialist legacy model (AUC=0.91) (Appendix 4.3). For all models, we converted the odds ratio to percentage points, where values higher than 0 indicated higher odds of abandonment on areas not farmed in the specific time period, compared to areas farmed then.

Results

We found strong land use legacy effects on land abandonment in our study region. As expected, the strength of the legacies diminished with time, but the differences in land abandonment in areas farmed under different political regimes were greatest. In the Carpathian region, agricultural land expanded until the 1960s, but after the 1930s, this expansion was to a large extent in in unsuitable areas. Abandonment was already strong in the socialist era, and continued during post-socialism. The effect of Habsburg land use legacies were stronger on socialist versus post-socialist abandonment, but we found the strongest legacies for post-socialist abandonment when comparing areas farmed during the Habsburg versus the socialist era.

Agricultural dynamics and their drivers

Agricultural use peaked in the study region in the 1960s, when agricultural land covered 38% of the study region (Figure 4.3). In 1860s, roughly 31% of the study area was arable, and this area increased during the time of the Habsburgs in all countries. Romania and Ukraine had the highest

percentage of land in agriculture in 1960s with 32% and 25% of their territory respectively. By 2010, only 20% of the study region was in agriculture (Figure 4.3).

Agricultural abandonment started in the Carpathians already during the late socialist era, when 34% of the agricultural land was abandoned, and continued throughout the post-socialist period, when 30% of the remaining agricultural land was abandoned. Between 1960 and 1985 the highest rates of abandonment occurred in the Polish Carpathians and the Southern Romanian Carpathians. Between 1985 and 2010 the highest land abandonment rates occurred in Ukraine, northern Romania and the southern Romanian Carpathians, where over 50% of the total agricultural land was abandoned by 2010. In some parts of Romania and Ukraine, virtually all agricultural land was abandoned. In Poland abandonment was substantially higher in the socialist period (68%) than in the post-socialist time (33%) (Figure 4.4).

When analyzing the relationship between agricultural land use and agro-ecological conditions, we found that socialist agricultural expansion occurred predominantly in areas that were less suitable, while both Habsburg and the remaining post-socialist agriculture were concentrated where agro-ecological conditions are favorable (AUC=0.82 and respectively AUC=0.89) (Figure 4.5a and c). Areas at low elevations, flatter slopes, closer to rivers and with higher precipitation and better crop suitability were more likely to be farmed in the Habsburg era. Conversely, areas of new agriculture during socialism, i.e., points converted to agriculture by either 1960s or 1985, were less well explained by agro-ecological conditions (AUC=0.69) (Figure 4.5b), and agricultural expansion happened mostly in areas of higher elevation and with lower crop suitability. Finally, points used for agriculture either in 2000 or 2010 were very well explained by agro-ecological factors (AUC 0.89) (Figure 4.5c). Areas at low elevations and flatter slopes,

closer to rivers, and with high crop suitability were most likely to remain in agriculture (Figure 4.5c).

Persistence of legacies

Clearly, agro-ecological conditions and accessibility are important spatial determinants of agricultural land use patterns. Furthermore, the spatial determinants affected both historical and recent land use, which means that just using historical land use to predict current land use patterns would greatly overestimate legacy effects, because historical land use would function as a proxy for agro-ecological conditions in such a univariate model. However, we found that even after controlling for agro-ecological, accessibility and socio-political variation, land use legacies effect were clearly evident in our data, persisted for as long as a century, and that their effect diminished slowly over time. When modelling socialist and post-socialist abandonment, we found that effects of Habsburg legacies existed for both abandonment periods. Agricultural abandonment was more likely in areas that were not farmed by the Habsburgs in 1860s (late Habsburg), than in areas farmed then (early Habsburg). The odds of socialist abandonment were 65% higher in areas converted to agriculture by Habsburgs after 1860s, compared to areas farmed before 1860s and the relationship was strong across all countries in the study region. The legacy effect was, however, smaller for post-socialist abandonment (46% higher odds), compared to the socialist abandonment (65%). In Poland (163%) and the Czech Republic (104%), the odds of socialist abandonment were especially high if land was farmed after 1860s. The relationship was weakest in Slovakia, where the odds of abandonment were only 39% higher for land farmed after 1860s compared to land farmed before 1860s (Figure 4.6a). Aside from legacy effects, we found that socialist abandonment was concentrated closer to settlement, on steeper slopes, and in areas with low crop suitability (Appendix 4.4).

When comparing the Habsburg legacy for socialist abandonment (65% higher odds) with the Habsburg legacy for post-socialist abandonment (46% higher odds), we found that land use legacies diminished over time. The odds of post-socialist abandonment were 46% higher in areas converted to agriculture after 1860s, compared to areas that were farmed then. However, the Habsburg legacy on post-socialist abandonment was strong for Romania (86% higher odds of abandonment) and Hungary (62%). In Slovakia and Czech Republic, the Habsburg legacy was practically absent (10% and 9% higher odds of abandonment) (Fig 6b). In addition to the legacy effects, our models showed that post-socialist abandonment was concentrated in areas with steep slopes, high precipitation, and low crop suitability (Appendix 4.4).

In summary, our results showed that even when accounting for agro-ecological and accessibility variation, legacy effects existed, but their effect diminished over time. In other words, the Habsburg legacy was stronger after 100 years than after 125 years.

Strength of land use legacies

To understand how the strength of land use legacies may be affected by underlying factors such as political regimes we compared the post-socialist legacy effect of three periods, the Habsburg era, the Interwar and the socialist era. Overall, we found that areas that were later converted to agricultural use had higher odds of abandonment compared to areas farmed earlier (Figure 4.7). Of all the areas abandoned by 2010, the most abandonment happened on land farmed by the Habsburgs (before 1930s), but overall socialist agriculture was abandoned at a higher rate than Habsburg agriculture (Appendix 4.2). Legacy effects were strong even when accounting for agro-ecological, accessibility and socio-political factors. All models were consistent in their variable selection: abandonment was predominantly on steeper slopes, and in areas with less

suitable soils and more precipitation. The socialist legacy models also indicated higher chance of abandonment in more accessible areas and closer to rivers (Appendix 4.5).

When comparing the legacy effects of difference in land use patterns among the three time periods, Habsburg, Interwar, and socialist, we found greatest differences for the Habsburg vs. socialist legacy, i.e., when comparing abandonment on lands farmed during the Habsburg era with agricultural expansion during the socialist era. Land farmed during the socialist era was 91% more likely to be abandoned compared to land farmed during the Habsburg era. This relationship was strong for all countries and the odds of abandonment were more than double for Hungary (158%), Romania (125%) and Czech Republic (122%). The legacy effect was weakest in Ukraine (20%). Our Habsburg vs. socialist models captured land use decisions of two eras with highly different land management strategies, the Habsburg era and the socialist era, and hence the greatest transformation in the underlying forces of land change (Figure 4.7).

We also found that the legacy of differences between early and late Habsburg land use patterns, i.e., the Habsburg legacy, on post-socialist abandonment was stronger than difference between early and late socialism, i.e., the socialist legacy. Land converted to agriculture late in the Habsburg era, i.e., after the 1860s, was 46% more likely to be abandoned compared to land farmed prior to the 1860s (Obj 1) (Figure 4.7). In contrast, land converted to agriculture late during socialism was only 23% more likely to be abandoned than land converted to agriculture early during socialism. The odds of abandonment were higher in Slovakia and Hungary (57% higher) and the legacy effect was very weak for Ukraine (12% higher odds of abandonment) and the Czech Republic (14% higher odds of abandonment). The sample size for Poland was too small to parameterize the model (total of 43 observations for this period).

Discussion

Our results showed that land use legacies can be strong in agricultural systems and that their strength differs in magnitude between historical political regimes. This means that considering historical land uses can add explanatory power in models of land change patterns, and contemporary land use decisions may affect environmental change for centuries into the future. We selected the Carpathian area as our study region because of its long land use history (Kozak et al., 2013a; Munteanu et al., 2014) and multiple institutional transformations (Bičík et al., 2001; Butsic et al., 2015), and found that historic land use strongly affected the rates of contemporary agricultural abandonment even after accounting for agro-ecological suitability and accessibility. We showed that underlying historic political and institutional contexts may result in legacy effects, i.e., that patterns of contemporary abandonment depend on whether land was farmed during different historical periods. Our findings support that land use legacies can shape all aspects of global environmental change (Foster et al., 2003; MacDonald et al., 2012; Perring et al., 2016).

In addition to prior studies that provide evidence for the existence of legacies in forest ecosystems (Dupouey et al., 2002; Munteanu et al., 2015a), agricultural and post-agricultural landscapes (Brudvig et al., 2013; Foster et al., 2003; Plieninger et al., 2006) and in species distribution patterns (Ficetola et al., 2010), we show here that different political regimes affect the strength of the observed legacies, even after accounting for environmental and accessibility factors. Our results advance land use modelling and highlight the importance of century-long effect human-environment interactions for contemporary environmental change. Maybe even more importantly, our result highlight the need making farsighted land management and

conservation decisions because these decisions will affect environmental change for centuries into the future.

Agricultural dynamics and their drivers

In order to better understand legacies for land abandonment, we asked first what the agricultural dynamics in the region were and which environmental and accessibility factors determined farming choices under different historic political regimes. We found that historically the choice of which land to farm was based on agro-ecological suitability and economic profitability (AUC=0.82), in line with land rent theories of Ricardo (Ricardo, 1821). Furthermore, environmental conditions explained well the distribution of the remaining agricultural land in the post-socialist era (AUC=0.89), likely a result of agricultural specialization, increasing land use efficiency and displacement of land use to areas outside Europe (Foley et al., 2011; Kastner et al., 2014; Meyfroidt et al., 2010). Less favorable areas, which were farmed during the socialist era, were abandoned first after the collapse of socialism. Our results also suggest that the high rates of post-socialist abandonment may be partly due to the socialist agricultural expansion on mostly unsuitable land for crops. When modelling agricultural expansion of the Soviet era, we found that the explanatory power of agro-ecological conditions was low (AUC=0.69) and that agricultural expansion was concentrated in areas at high elevations, on low suitability soils and farther away from rivers. These statistical results support an extensive body of anecdotal evidence that socialist agricultural expansion was driven by political goals and disregarded environmental conditions (Bičík et al., 2001; Štych et al., 2012). The expansion on unsuitable land, followed by economic hardship and collapse of the centralized system likely contributed to the high land abandonment rates observed both during and after the collapse of the Soviet Union (Baumann et al., 2011; Munteanu et al., 2015a). This is perhaps unsurprising, because if areas

with low agricultural suitability were farmed during the Soviet era, then these areas should have high rates of post-socialist abandonment according to land rent theory. Overall, spatial reorganization of agriculture and the industrialization factors are key in explaining land abandonment across other parts of Europe and the former Eastern Bloc (Jepsen et al., 2015). Our results regarding agricultural dynamics in the Carpathian region confirm regional trends that were previously only documented in local case studies from eastern and central European countries (Gerard et al., 2010; Kozak, 2003; Mojses and Petrovič, 2013). In the Carpathians agricultural land expanded during the Habsburg and Interwar time periods, and land abandonment was widespread during the socialist and the post-socialist eras. During the socialist era, we observed particularly high abandonment rates in southern Poland, a process likely related to the political decision to displace the Ruthenian population from the Polish territory, immediately after WW II (Woś, 2005). Abandonment rates were also high in the Southern Romanian Carpathians, where abandonment in mountain regions may be related to the forced industrialization policies of the Ceausescu regime, which displaced farmers to industrial centers (Ban, 2012). Between 1985 and 2010 about 0.84% of the cultivated land was abandoned annually in the Carpathians, compared to 0.72% across Europe for 2001-2012 (Estel et al., 2015). The widespread abandonment during the post-socialist period is likely linked to institutional changes and restructuring of property rights following the collapse of the Soviet Union in 1990 (Estel et al., 2015; Jepsen et al., 2015; Levers et al., 2014). Abandonment in the post-socialist period was particularly high in the Ukrainian Carpathians and in northern Romania, likely due to restructuring of economies (Baumann et al., 2011; Griffiths et al., 2013; Hartvigsen, 2014). Economic reforms and the opening of the borders caused large numbers of

people to seek better economic opportunities in urban areas or abroad, leading to abandonment of agricultural activities (Baumann et al., 2011; Munteanu et al., 2015b).

Last but not least, our results represent the first long-term cross-border assessment of agricultural dynamics in the Carpathian region and provide evidence for an ‘agricultural transition’, with the region as a whole experiencing the highest point in agricultural cover during the Soviet era. This transition point, occurred with a time lag compared to the forest transition in the region, which occurred during the Interwar period (Munteanu et al., 2015a, 2014).

Persistence of legacies

When modelling agricultural abandonment as function as historical land uses and environmental factors, we found that similar to forest legacies from a century ago (Munteanu et al., 2015a) agricultural legacies can persist for as long as a century in the land system, but their effect generally diminishes over time. Even after controlling for other spatial determinants of land change (Meyfroidt, 2015), we found that historical land use patterns had strong explanatory power in our land abandonment models. This finding provides quantitative evidence for path dependency in agricultural systems, suggesting that continued agricultural use is more likely in areas that were used for agriculture in the past. Our result is consistent with prior finding on path dependency showing that once a land use type is established, land change is less likely (Coomes et al., 2011; Seto et al., 2011; Verburg et al., 2004). Despite the overall trends, country differences were stark though, indicating that national shifts land use and land management policies may affect the strength or occurrence of the legacy effect.

Strength of land use legacies

Political and socio-economic conditions, which varied considerably between the Habsburg and the Socialist eras, greatly affected the land use legacies by causing drastic historical land change, which in turn was reflected in contemporary land use patterns. Specifically, we found highest differences in the odds of agricultural abandonment when comparing areas farmed during the Habsburg time period with those where agricultural expanded during socialism (91% higher odds of abandonment on Socialist vs. Habsburg agriculture even when accounting for agro-ecological and accessibility variation). We suggest that this observed legacy effect is related to differences in land management strategies during the two politically distinct time periods. In the Habsburg time period, in addition to agriculture being concentrated in agro-ecologically suitable areas (see Obj 1), agriculture was dominated by large land holdings by the nobility (Good, 1984), who made land use choices based on land productivity. Furthermore, the Habsburg empire promoted the homogenization of economic activities in least developed parts of empire, by supporting agriculture and industrial development in parts of Romania and Hungary (Good, 1984). In contrast, the socialist era land reforms included nationalization and collectivization, expansion of agriculture under heavy subsidies, economies of scale, and the heavy use of machinery and fertilizers (Bičík et al., 2001; Lerman et al., 2004), which proved unsustainable after 1990. Along with the environmental factors, these differences in land use policies between the two regimes most likely caused higher rates of abandonment on new socialist agriculture compared to Habsburg agriculture.

Land use legacies were also present when comparing odds of abandonment on late and early Habsburg agriculture (46% higher odds) and when comparing odds of abandonment on late and early Socialist agriculture (23% higher odds). However, these differences were not as stark as

when comparing legacies of different political regimes (Habsburg vs. Socialist). We suggest that this Habsburg legacy effect is explained by a change in policy and land management around the 1850s, aimed at homogenization of economic disparities between the Eastern and Western parts of the Habsburg Empire (Good, 1984). The empire conducted major reforms around the 1850s, which fostered agricultural production in Hungary and the Carpathian Basin, including adjustments in land ownership, liberation of the serfs (1948), and the removal of agricultural tariffs for Eastern Empire provinces (1850) (Alix-Garcia and Walker, 2016; Good, 1984). Furthermore, in the last third of the 19th century, the use of machinery and new agricultural tools skyrocketed, causing an increase in agricultural production. We suggest that these policy and technological changes might be related to the observed differences in abandonment between areas farmed before and after 1860s in the Habsburg era. Our explanation is supported by the fact that the legacy effect was strongest in Hungary and Romania, the two countries most affected by post-1860s Habsburg agricultural policies.

Legacies of the differences of land use patterns in the early versus the late Socialist era on post-socialist abandonment were small. Land converted to agriculture early in the Socialist era (after 1930s but before 1960s) was only 23% more likely to be abandoned than land converted to agriculture later in the Socialist era (between 1960s and 1985). Following WW II, governments throughout Eastern Europe adopted the agrarian philosophy of the Soviet Union, including collectivization of agriculture, economies of scale, increased application of fertilizers, use of heavy machinery, and agricultural reforms, albeit to varying degrees (Bezák and Mitchley, 2014; Lerman et al., 2004; Nelson, 1993). Despite national differences, the outcome on agricultural land was similar before and after the 1960s, in both periods there was forceful expansion and intensification, which led ultimately to abandonment of most of the land farmed during socialism

after the collapse of the Soviet Union. In other words, we did not strong differences in the odds of abandonment on early versus late Socialist agriculture, because socialist agricultural expansion was ecologically and economically unfeasible, regardless of when it occurred.

While our overall results revealed very clear patterns, we caution that our analysis is based on binary agricultural data and that the definition of agriculture here does not include grassland dynamics. We considered any transition from agriculture to non-agriculture as abandonment, but different definition of abandonment (e.g., only cropland-grassland conversions) may have yielded different results. We also caution that although the length of the abandonment periods that we studied as the same (25 years), the legacy eras have different durations. The nature of the data used to map socialist abandonment (Landsat based agricultural cover in 1985, 2000 and 2010), did not allow us to separate the land abandonment that occurred between 1985 and 1990 (the year that the Soviet Union collapsed). As a consequence, our estimation of ‘socialist abandonment’ might be conservative and the difference in legacy effects of the two periods (socialist and post-socialist) could be even greater than indicated by our results. We suspect that the weak legacy effect observed for the socialist period is partly related to the short period of socialist agricultural expansion considered here compared to the Habsburg period. We further caution that the historical maps used in our analysis depict land use, whereas remote sensing data depicts land cover. In terms of arable areas, these two datasets are comparable, because annual and perennial crops are depicted as such in land-use maps, and they are identified in satellite imagery by tilled soil in the early growing season.

Our study confirmed and reinforced the importance of land use legacies for contemporary and future global land change. Moreover, different political regimes may be associated with greatest legacy effects, suggesting that areas in where multiple socio-economic changes occurred over

time legacies may persist for a very long time into the future. In a scientific context, the consideration of past land uses as spatial determinants of change could enhance the performance of land use models at regional and global scale, and can improve the prediction of future land changes. In a land management context, we stress the importance of considering the effects that contemporary land use decisions have on centuries to come, which highlight the responsibility our generation has for future generations when making land management and political decisions.

References

- Alcantara, C., Kuemmerle, T., Baumann, M., Bragina, E. V., Griffiths, P., Hostert, P., Knorn, J., Müller, D., Prishchepov, A. V., Schierhorn, F., Sieber, A., Radeloff, V.C., 2013. Mapping the extent of abandoned farmland in Central and Eastern Europe using MODIS time series satellite data. *Environ Res Lett* 8, 035035.
- Alix-Garcia, J., Walker, S., 2016. Tariffs and Trees : The effects of the Austro-Hungarian customs union on specialization and land use change. *Eur Econ Rev* in review.
- Ban, C., 2012. Sovereign Debt, Austerity, and Regime Change: The Case of Nicolae Ceausescu's Romania. *East Eur Polit Soc* 26, 743–776.
- Baumann, M., Kuemmerle, T., Elbakidze, M., Ozdogan, M., Radeloff, V.C., Keuler, N.S., Prishchepov, A. V., Kruhlov, I., Hostert, P., 2011. Patterns and drivers of post-socialist farmland abandonment in Western Ukraine. *Land use policy* 28, 552–562.
- Berger, S. (Ed.), 2006. *A Companion to Nineteenth-Century Europe: 1789-1914*. Blackwell Publishing, Malden, Oxford, Carlton.
- Bezák, P., Mitchley, J., 2014. Drivers of change in mountain farming in Slovakia: from socialist collectivisation to the Common Agricultural Policy. *Reg Environ Chang* 14, 1343–1356.
- Bičík, I., Jeleček, L., Štěpánek, V., 2001. Land-use changes and their social driving forces in Czechia in the 19th and 20th centuries. *Land use policy* 18, 65–73.
- Börjeson, L., 2007. Boserup backwards? Agricultural intensification as “its own driving force” in the Mbulu Highlands, Tanzania. *Geogr Ann Ser B Hum Geogr* 89, 249–267.
- Brain, S., 2010. The Great Stalin Plan for the Transformation of Nature. *Environ Hist Durh N C* 15, 670–700.
- Brudvig, L.A., Grman, E., Habeck, C.W., Orrock, J.L., Ledvina, J.A., 2013. Strong legacy of agricultural land use on soils and understory plant communities in longleaf pine woodlands. *For Ecol Manage* 310, 944–955.
- Butsic, V., Munteanu, C., Griffiths, P., Knorn, J., Radeloff, V.C., Kuemmerle, T., 2016. The impact of protected areas on forest harvesting in the Carpathian Mountains 1985-2010 *Cons Biol*, in review.
- Coomes, O.T., Takasaki, Y., Rhemtulla, J.M., 2011. Land-use poverty traps identified in shifting cultivation systems shape long-term tropical forest cover. *Proc Natl Acad Sci USA* 108, 13925–30.
- DeFries, R.S., Foley, J.A., Asner, G.P., 2004. Land-use choices : balancing human needs and ecosystem function In a nutshell : *Frontiers Ecol* 2, 249–257.
- Dullinger, S., Essl, F., Rabitsch, W., Erb, K.-H., Gingrich, S., Haberl, H., Hülber, K., Jarosík, V., Krausmann, F., Kühn, I., Pergl, J., Pyšek, P., Hulme, P.E., 2013. Europe's other debt crisis caused by the long legacy of future extinctions. *Proc Natl Acad Sci USA* 110, 7342–7.

- Dupouey, J.L., Dambrine, E., Laffite, J.D., Moares, C., 2002. Irreversible Impact of Past Land Use on Forest Soils and Biodiversity. *Ecology* 83, 2978–2984.
- Essl, F., Dullinger, S., Rabitsch, W., Hulme, P.E., Pyšek, P., Wilson, J.R.U., Richardson, D.M., 2015. Historical legacies accumulate to shape future biodiversity in an era of rapid global change. *Divers Distrib* 21, 534–547.
- Estel, S., Kuemmerle, T., Alcántara, C., Levers, C., Prishchepov, A., Hostert, P., 2015. Mapping farmland abandonment and recultivation across Europe using MODIS NDVI time series. *Remote Sens Environ* 163, 312–325.
- Ficetola, G.F., Maiorano, L., Fallucci, A., Dendoncker, N., Boitani, L., Padoa-Schioppa, E., Miaud, C., Thullier, W., 2010. Knowing the past to predict the future: land-use change and the distribution of invasive bullfrogs. *Glob Chang Biol* 16, 528–537.
- Foley, J.A., Ramankutty, N., Brauman, K. a, Cassidy, E.S., Gerber, J.S., Johnston, M., Mueller, N.D., O’Connell, C., Ray, D.K., West, P.C., Balzer, C., Bennett, E.M., Carpenter, S.R., Hill, J., Monfreda, C., Polasky, S., Rockström, J., Sheehan, J., Siebert, S., Tilman, D., Zaks, D.P.M., 2011. Solutions for a cultivated planet. *Nature* 478, 337–42.
- Foster, D.R., Swanson, F.J., Aber, J., Burke, I., Brokaw, N., Tilman, D., Knapp, A., 2003. The importance of land-use legacies to ecology and conservation. *Bioscience* 53, 77–88.
- Freeman, E.A., Moisen, G.G., 2008. A comparison of the performance of threshold criteria for binary classification in terms of predicted prevalence and kappa. *Ecol Modell* 217, 48–58.
- Fuchs, R., Herold, M., Verburg, P.H., Clevers, J.G.P.W., Eberle, J., 2014. Gross changes in reconstructions of historic land cover/use for Europe between 1900 and 2010. *Glob Chang Biol* 21, 299–313.
- Gallego, J., Delincé, J., 2010. The European land use and cover area-frame statistical survey, in: Benedetti, R., Bee, M., Espa, G., Piersimoni, F. (Eds.), *Agricultural Survey Methods*. John Wiley & Sons, pp. 151–168.
- Geist, H.J., Lambin, E.F., 2002. Proximate Causes and Underlying Driving Forces of Tropical Deforestation. *Bioscience* 52, 143.
- Gerard, F., Petit, S., Smith, G., Thomson, A., Brown, N., Manchester, S., Wadsworth, R., Bugár, G., Halada, L., Bezak, P., Boltižiar, M., De badts, E., Halabuk, A., Mojses, M., Petrovič, F., Gregor, M., Hazeu, G., Mucher, C. a., Wachowicz, M., Huitu, H., Tuominen, S., Kohler, R., Olschofsky, K., Ziese, H., Kolar, J., Sustera, J., Luque, S., Pino, J., Pons, X., Roda, F., Roscher, M., Feranec, J., 2010. Land cover change in Europe between 1950 and 2000 determined employing aerial photography. *Prog Phys Geogr* 34, 183–205.
- Good, D.F., 1984. *The Economic Rise of the Habsburg Empire, 1750-1914*. University of California Press, Berkley and Los Angeles.
- Griffiths, P., Müller, D., Kuemmerle, T., Hostert, P., 2013. Agricultural land change in the Carpathian ecoregion after the breakdown of socialism and expansion of the European Union. *Environ Res Lett* 8, 045024.

- Hartvigsen, M., 2014. Land reform and land fragmentation in Central and Eastern Europe. *Land use policy* 36, 330–341.
- Hosmer, D.W., Jr., Lemeshow, S., Sturdivant, R.X., 2013. *Applied Logistic Regression*. John Wiley & Sons.
- Hurt, G.C., Frolking, S., Fearon, M.G., Moore, B., Shevliakova, E., Malyshev, S., Pacala, S.W., Houghton, R. a., 2006. The underpinnings of land-use history: three centuries of global gridded land-use transitions, wood-harvest activity, and resulting secondary lands. *Glob Chang Biol* 12, 1208–1229.
- Jepsen, M.R., Kuemmerle, T., Müller, D., Erb, K., Verburg, P.H., Haberl, H., Vesterager, J.P., Andrič, M., Antrop, M., Austrheim, G., Björn, I., Bondeau, A., Bürgi, M., Bryson, J., Caspar, G., Cassar, L.F., Conrad, E., Chromý, P., Daugirdas, V., Van Eetvelde, V., Elena-Rosselló, R., Gimmi, U., Izakovicova, Z., Jančák, V., Jansson, U., Kladnik, D., Kozak, J., Konkoly-Gyuró, E., Krausmann, F., Mander, Ü., McDonagh, J., Pärn, J., Niedertscheider, M., Nikodemus, O., Ostapowicz, K., Pérez-Soba, M., Pinto-Correia, T., Ribokas, G., Rounsevell, M., Schistou, D., Schmit, C., Terkenli, T.S., Tretvik, A.M., Trzepacz, P., Vadineanu, A., Walz, A., Zhllima, E., Reenberg, A., 2015. Transitions in European land-management regimes between 1800 and 2010. *Land use policy* 49, 53–64.
- Kaim, D., Kozak, J., Ostafin, K., Dobosz, M., Ostapowicz, K., Kolecka, N., Gimmi, U., 2014. Uncertainty in historical land-use reconstructions with topographic maps. *Quaest Geogr* 33, 55–63.
- Kastner, T., Erb, K.-H., Haberl, H., 2014. Rapid growth in agricultural trade: effects on global area efficiency and the role of management. *Environ Res Lett* 9, 034015.
- Kozak, J., 2003. Forest Cover Change in the Western Carpathians in the Past 180 Years. *Mt Res Dev* 23, 369–375.
- Kozak, J., Ostapowicz, K., Byterowicz, A., Wyzga, B., 2013a. The Carpathian Mountains: Challenges for the Central and Eastern European Landmark, in: *The Carpathians: Integrating Nature and Society Towards Sustainability*. Springer, Berlin, Heidelberg, pp. 1–11.
- Kozak, J., Ostapowicz, K., Bytnerowicz, A., Wyżga, B. (Eds.), 2013b. *The Carpathians: Integrating Nature and Society Towards Sustainability*, Environmental Science and Engineering. Springer Berlin Heidelberg, Berlin, Heidelberg.
- Kraemer, R., Prishchepov, A. V., Müller, D., Kuemmerle, T., Radeloff, V.C., Dara, A., Terekhov, A., Frühauf, M., 2015. Long-term agricultural land-cover change and potential for cropland expansion in the former Virgin Lands area of Kazakhstan. *Environ Res Lett* 10, 054012.
- Lambin, E.F., Geist, H.J. (Eds.), 2006. *Land-Use and Land-Cover Change - Local Processes and Global Impacts*, Global Cha. ed. Springer.
- Lambin, E.F., Meyfroidt, P., 2011. Global land use change, economic globalization, and the looming land scarcity. *Proc Natl Acad Sci USA* 108, 3465–3472.

- Lerman, Z., Csáki, C., Feder, G., 2004. *Agriculture in Transition: Land Policies and Evolving Farm Structures in Post-Soviet Countries*. Lexington Books.
- Levers, C., Verkerk, P.J., Müller, D., Verburg, P.H., Butsic, V., Leitão, P.J., Lindner, M., Kuemmerle, T., 2014. Drivers of forest harvesting intensity patterns in Europe. *For Ecol Manage* 315, 160–172.
- Lohr, S., 2010. *Sampling: Design and Analysis*, 2nd ed. Brooks/Cole, Boston MA.
- MacDonald, D., Crabtree, J., Wiesinger, G., Dax, T., Stamou, N., Fleury, P., Gutierrez Lazpita, J., Gibon, a, 2000. Agricultural abandonment in mountain areas of Europe: Environmental consequences and policy response. *J Environ Manage* 59, 47–69.
- MacDonald, G.K., Bennett, E.M., Taranu, Z.E., 2012. The influence of time, soil characteristics, and land-use history on soil phosphorus legacies: a global meta-analysis. *Glob Chang Biol* 18, 1904–1917.
- Matteucci, S.D., Totino, M., Arístide, P., 2016. Ecological and social consequences of the Forest Transition Theory as applied to the Argentinean Great Chaco. *Land use policy* 51, 8–17.
- Mattingly, W.B., Orrock, J.L., Collins, C.D., Brudvig, L. a, Damschen, E.I., Veldman, J.W., Walker, J.L., 2014. Historical agriculture alters the effects of fire on understory plant beta diversity. *Oecologia* 177, 507–518.
- McConnell, W.J., Keys, E., 2005. Meta-analysis of agricultural change, in: Moran, E.F., Ostrom, E. (Eds.), *Seeing the Forest and the Trees. Human-Environment Interactions in Forest Ecosystems*. MIT Press, Cambridge, pp. 325–354.
- Meyfroidt, P., 2015. Approaches and terminology for causal analysis in land systems science. *J Land Use Sci* 4248, 1–27.
- Meyfroidt, P., Lambin, E.F., Erb, K.-H., Hertel, T.W., 2013. Globalization of land use: distant drivers of land change and geographic displacement of land use. *Curr Opin Environ Sustain* 5, 438–444.
- Meyfroidt, P., Rudel, T.K., Lambin, E.F., 2010. Forest transitions, trade, and the global displacement of land use. *Proc Natl Acad Sci USA* 107, 20917–22.
- Mojšes, M., Petrovič, F., 2013. Land use changes of historical structures in the agricultural landscape at the local level - Hriňova case study. *Ekológia Int J Ecol Probl Biosph* 32, 1–12.
- Müller, D., Leitão, P.J., Sikor, T., 2013. Comparing the determinants of cropland abandonment in Albania and Romania using boosted regression trees. *Agric Syst* 117, 66–77.
- Munroe, D.K., van Berkel, D.B., Verburg, P.H., Olson, J.L., 2013. Alternative trajectories of land abandonment: causes, consequences and research challenges. *Curr Opin Environ Sustain* 1–6.
- Munteanu, C., Kuemmerle, T., Boltiziar, M., Butsic, V., Gimmi, U., Kaim, D., Király, G., Konkoly-Gyuró, É., Kozak, J., Lieskovský, J., Mojšes, M., Müller, D., Ostafin, K.,

- Ostapowicz, K., Shandra, O., Štych, P., Walker, S., Radeloff, V.C., 2014. Forest and agricultural land change in the Carpathian region—A meta-analysis of long-term patterns and drivers of change. *Land use policy* 38, 685–697.
- Munteanu, C., Kuemmerle, T., Keuler, N.S., Müller, D., Balazs, P., Dobosz, M., Griffiths, P., Halada, L., Kaim, D., Király, G., Konkoly-Gyuró, É., Kozak, J., Lieskovský, J., Ostafin, K., Ostapowicz, K., Shandra, O., Radeloff, V.C., 2015a. Legacies of 19th century land use shape contemporary forest cover. *Glob Environ Chang* 34, 83–94.
- Munteanu, C., Nita, M.-D., Abrudan, I. V., Radeloff, V.C., 2016. Historical forest management in Romania is imposing strong legacies on contemporary forests and their management. *For Ecol Manage* 361, 179–193.
- Munteanu, C., Radeloff, V., Griffiths, P., Halada, L., Kaim, D., Knorn, J., Kozak, J., Kuemmerle, T., Lieskovsky, J., Müller, D., Ostapowicz, K., Shandra, O., Stych, P., 2015b. Land change in the Carpathian Region before and after major institutional changes, in: Gutman, G., Radeloff, V.C. (Eds.), *Land Use and Land Cover Change in Eastern Europe after the Collpase of Socialism*. Springer, p. in press.
- Nagendra, H., Southworth, J., Tucker, C., 2003. Accessibility as a determinant of landscape transformation in western Honduras: linking pattern and process. *Landsc Ecol* 18, 141–158.
- Nelson, G.C., 1993. Agricultural Policy Reform in Eastern Europe: Discussion. *Am J Agric Econ* 75, 857.
- Perring, M.P., De Frenne, P., Baeten, L., Maes, S.L., Depauw, L., Blondeel, H., Carón, M.M., Verheyen, K., 2016. Global environmental change effects on ecosystems: the importance of land-use legacies. *Glob Chang Biol* 22, 1361–1371.
- Plieninger, T., 2014. Socialist and postsocialist land use legacies determine farm and woodland composition and structure: lessons form Eastern Germany. *Eur J For Res* 133, 597–610.
- Plieninger, T., Höchtl, F., Spek, T., 2006. Traditional land-use and nature conservation in European rural landscapes. *Environ Sci Policy* 9, 317–321.
- Plieninger, T., Schaich, H., Kizos, T., 2010. Land-use legacies in the forest structure of silvopastoral oak woodlands in the Eastern Mediterranean. *Reg Environ Chang* 11, 603–615.
- Plue, J., Dupouey, J.L., Verheyen, K., Hermy, M., 2009. Forest seed banks along an intensity gradient of ancient agriculture. *Seed Sci Res* 19, 103–114.
- Plue, J., Hermy, M., Verheyen, K., Thuillier, P., Saguez, R., Decocq, G., 2008. Persistent changes in forest vegetation and seed bank 1,600 years after human occupation. *Landsc Ecol* 23, 673–688.
- Prishchepov, A. V., Müller, D., Dubinin, M., Baumann, M., Radeloff, V.C., 2013. Determinants of agricultural land abandonment in post-Soviet European Russia. *Land use policy* 30, 873–884.

- Prishchepov, A. V., Radeloff, V.C., Baumann, M., Kuemmerle, T., Müller, D., 2012. Effects of institutional changes on land use: agricultural land abandonment during the transition from state-command to market-driven economies in post-Soviet Eastern Europe. *Environ Res Lett* 7, 024021.
- Rey Benayas, J., 2007. Abandonment of agricultural land: an overview of drivers and consequences. *CAB Rev Perspect Agric Vet Sci Nutr Nat Resour* 2, 0–14.
- Ricardo, D., 1821. *On the Principles of Political Economy and Taxation*, Third Edit. ed. John Murray, Albemarle Street, London.
- Schelhaas, M.-J., Nabuurs, G.-J., Schuck, A., 2003. Natural disturbances in the European forests in the 19th and 20th centuries. *Glob Chang Biol* 9, 1620–1633.
- Seto, K.C., Fragkias, M., Güneralp, B., Reilly, M.K., 2011. A meta-analysis of global urban land expansion. *PLoS One* 6, e23777.
- Seto, K.C., Reenberg, A., Boone, C.G., Fragkias, M., Haase, D., Langanke, T., Marcotullio, P., Munroe, D.K., Olah, B., Simon, D., 2012. Urban land teleconnections and sustainability. *Proc Natl Acad Sci USA* 109, 7687–92.
- Štych, P., Bičík, I., Stonáček, V., Bláha, J., 2012. Change of Land Use Patterns 1827-2005, in: Bičík, I., Himiyama, Y., Feranec, J., Štych, P. (Eds.), *Land Use/Cover Changes in Selected Regions in the World. Volume VII. IGU-LUCC*, Prague, pp. 37–41.
- Verburg, P.H., Eck, J.R.R. Van, Nijs, T.C.M. De, Dijst, M.J., Schot, P., 2004. Determinants of land-use change patterns in the Netherlands. *Environ Plan B Plan Des* 31, 125–150.
- Woś, B., 2005. Zmiany pokrycia terenu w wybranych gminach Beskidów w drugiej połowie XX w. na podstawie analizy zdjęć lotniczych (in Polish). *Teledetekcja Środowiska* 35, 1–114.

Tables and figures

Table 4.1 Paper terminology

Agricultural land	We define agricultural land here as tilled areas, covered by annual or perennial crops. Our definition of agricultural land does not include pastures or grasslands.
Habsburg era	In this manuscript, the time from 1800s to 1914, during which the most of the study area was part of the Habsburg Empire (except the southern ridge of the Carpathian Mountains in Romania). We refer to early Habsburg as the period before the 1860s and to late Habsburg thereafter.*
Interwar era	1914-1945, time of major political transformations due to the two World Wars, and the Great Depression. The period is marked at both ends by the Habsburg and the Socialist political regimes.
Socialist era	In this manuscript, 1945-1985, during which all countries in the study area were influenced by Moscow politics. We refer to early socialist as the period before the 1960s and to late socialist thereafter.*
Post-socialist era	In this manuscript, 1985-2010, during which most countries in the study region changed to market economies and accessed the EU (except Ukraine).*
Socialist abandonment	Land abandonment that occurred between the 1960s and 1985, mapped using military topographic maps (1960s) and Landsat image composites (1985). Note that this does not capture the entire socialist era, but a period of 25 years, to ensure comparability with post-socialist abandonment.
Post-socialist abandonment	Land abandonment that occurred between 1985 and 2010, mapped using Landsat images for 1985, 2000 and 2010.*
Land use legacy	The effects of historical land uses and land use decisions on contemporary land use change, once other spatial determinants of change are accounted for.
Habsburg legacy	The effect of whether or not land was farmed prior to the 1860s on subsequent agricultural abandonment. This legacy captures the effect of early versus late Habsburg agriculture on abandonment.
Habsburg vs. Soviet legacy	The effect of the whether or not land was farmed prior to the 1930s on subsequent abandonment. This legacy captures the differences between land farmed during the Habsburg era and agricultural expansion during the socialist era.**
Socialist legacy	The effect of whether or not land was farmed prior to the 1960s on subsequent abandonment. This legacy captures the effect of early versus late Soviet agricultural expansion on abandonment.

* We note that the historical periods are not fully captured by our analysis period. We use the same names for simplification.

** The Interwar era represented a time of major political and socio-economic changes with largely stable agricultural dynamics throughout the period (Munteanu et al., 2014). We use this point in time to capture effects of two major land management systems that mark the interwar period: Habsburg and Socialist.

Table 4.2 Map and remote sensing data sources used to map agricultural land cover in the Carpathians


Time layer	Data range of maps	Map scale/ resolution	Map source/ description
1860s	1819-1873	1:28.000	Second Austrian Military Survey
1930s	1919-1939	1:100.000	Maps of Wojskowy Instytut Geograficzny (WIG)
	1923-1938	1:20.000	Preliminary Beneš maps, Definitive Křovák maps
	1923-1945	1:25.000	Revised Third Military Survey, German topographic maps (“Messtischblatt”, Karte der Slowakei, Karte der Tsechoslowakei)
1960s	1949-1983	1:50.000 and 1:25.000	Soviet and National Military Maps from the Cold War period
1985	1982-1987	30m	Landsat TM composite
2000	2003-2007	30m	Landsat TM/ ETM+ composite
2010	2008-2012	30m	Landsat TM/ ETM+ composite

Table 4.3 List of predictor used in logistic regression models, including land use legacies, agro-ecological conditions, accessibility and socio-political variables.

	Description	Source	Unit	Spatial Resolution
Response	Agricultural abandonment between 1960 and 1985	mapped, Griffiths et al, 2013	Yes/No	30 m
	Agricultural abandonment between 1985 and 2010	Griffiths et al, 2013	Yes/No	30 m
Historic land use	Habsburg agriculture (1860)	mapped		vector
	Interwar agriculture (1930)	mapped		vector
	Soviet agriculture (1960)	mapped		vector
Agro-ecological	Elevation	Farr et al, 2007	m	90 m
	Slope	Farr et al, 2007	°	90 m
	Annual Mean Temperature	Hijmans et al, 2005	C° * 10	~1 km
	Annual Precipitation in mm\	Hijmans et al, 2005	mm	~1 km
	Crop suitability index	FAO (GAEZ), 2014	%	~8 km
	Length of growing season	FAO (GAEZ), 2014	days	~8 km
Accessibility	Travel time to the nearest town with 50,000 inhabitants	Nelson, 2008	minutes	~1 km
	Distance to nearest major city	ESRI, 2008	km	vector
	Distance to nearest settlement	EEA, 2013	km	vector
	Distance to nearest road	CIESIN & ITOS, 2013	km	vector
	Distance to nearest current border	calculated	km	vector
	Distance to nearest railroad	ESRI, 2008	km	vector
	Distance to nearest main river	Vogt et al 2007	km	vector
Socio-political	Country	ESRI, 2008	N/A	vector
	Population count 1990	CIESIN, FAO & CIAT, 2005	No.	~5 km

Figure 4.1 Study objectives: persistence and strength of land use legacies. For Obj. 1 we developed two models (A,B), both of which compared abandonment of 1930s agricultural land that was already farmed in 1860 with land that was not farmed then, and we modelled the effect of early versus late Habsburg agriculture on both A) socialist and B) post-socialist land abandonment.. For Obj. 2, we modelled the legacy effect of three periods, i.e., the C) Habsburg, D) Interwar and E) Socialist period, on post-socialist abandonment. Models C is identical to model B for objective 2. Model D compares land farmed at any point during the Habsburg era with land farmed at any time during the Socialist period. Model E refers to socialist agricultural expansion and compares abandonment on land farmed prior to 1960, i.e., during the early Socialist period, with land farmed after 1960 (i.e. late Socialist). Please see Table 1 for definitions of terms and time periods

Obj.2: Persistence of legacies over time



Obj.3: Strength of legacies in relation to political regime




Figure 4.2 Study area. Country codes: CZ: Czech Republic, HU: Hungary PL: Poland, RO: Romania, SK: Slovakia, UA: Ukraine




Figure 4.3 Agricultural dynamics in the Carpathian region and by country.




Figure 4.4 Agricultural cover between 1860 and 2010, socialist and post-socialist land abandonment in the Carpathian region (see Table 4.1 for definitions).




Figure 4.5 Model performance for distribution of agricultural land as a function of environmental variables during the Habsburg, socialist and post-socialist eras. Model performance is reported as area under the receiver operator curve (AUC), where values close to 1 indicate high model performance.




Figure 4.6 Persistence of Habsburg land use legacies. Odds of abandonment (in %) in areas that were not farmed by the Habsburgs in 1860s, compared to areas farmed then. The odds of abandonment on land not farmed in 1860s were higher and the relationship was consistent across countries. For example, in Romania, the odds of socialist abandonment (a) were 63% higher in areas farmed in 1860s, compared to areas not farmed then and the odds of post-socialist abandonment (b) were 86% higher.





Figure 4.7 Strength of land use legacies of political regimes. Odds of abandonment (in %) in areas that were not farmed in a given historic period, compared to areas farmed then. The odds of abandonment on land not farmed historically were higher and the relationship was consistent across countries. For example, in Romania, the odds of abandonment were 86% higher in areas farmed in 1860s, compared to areas not farmed then. Legacy effects were strongest for the political transformation legacy, which captures differences in political regimes between the Habsburg and Socialist periods.



Appendix

Appendix 4.1 Sample sizes for legacy persistence modelling

Socialist abandonment (1960-1985)			Post-socialist abandonment (1985-2010)		
	1860 Non- AG	1860 AG		1860 Non- AG	1860 AG
Overall		20501	Overall		13419
not abandoned	3013	10406	not abandoned	1988	7350
abandoned	2002	5080	abandoned	1025	3056
ROMANIA		6013	ROMANIA		4251
not abandoned	924	3327	not abandoned	355	1749
abandoned	577	1185	abandoned	569	1578
HUNGARY		4114	HUNGARY		2849
not abandoned	893	1956	not abandoned	702	1666
abandoned	506	759	abandoned	191	290
SLOVAKIA		4586	SLOVAKIA		3143
not abandoned	736	2407	not abandoned	653	1988
abandoned	282	1161	abandoned	83	419
UKRAINE		1101	UKRAINE		547
not abandoned	169	378	not abandoned	40	91
abandoned	243	311	abandoned	129	287
CZECH REPUBLIC		2678	CZECH REPUBLIC		1998
not abandoned	248	1750	not abandoned	213	1461
abandoned	160	520	abandoned	35	289
POLAND		2009	POLAND		631
not abandoned	43	588	not abandoned	25	395
abandoned	234	1144	abandoned	18	193

Appendix 4.2 Sample sizes for strength of legacies in relation to political regimes

Habsburg legacy		
	1860 Non-AG	1860 AG
Overall		13419
not		
abandoned	1988	7350
abandoned	1025	3056
ROMANIA		4251
not		
abandoned	355	1749
abandoned	569	1578
HUNGARY		2849
not		
abandoned	702	1666
abandoned	191	290
SLOVAKIA		3143
not		
abandoned	653	1988
abandoned	83	419
UKRAINE		547
not		
abandoned	40	91
abandoned	129	287
CZECH REPUBLIC		1998
not		
abandoned	213	1461
abandoned	35	289
POLAND		631
not		
abandoned	25	395
abandoned	18	193

Habsburg vs. Socialist legacy		
	1930 Non-AG	1930 AG
Overall		16843
not		
abandoned	1496	9338
abandoned	1928	4081
ROMANIA		5821
not		
abandoned	372	2104
abandoned	1198	2147
HUNGARY		3416
not		
abandoned	396	2368
abandoned	171	481
SLOVAKIA		3748
not		
abandoned	451	2641
abandoned	154	502
UKRAINE		901
not		
abandoned	76	131
abandoned	278	416
CZECH REPUBLIC		2283
not		
abandoned	181	1674
abandoned	104	324
POLAND		674
not		
abandoned	20	420
abandoned	23	211

Socialist legacy		
	1960 Non-AG	1960 AG
Overall		3424
not		
abandoned	934	562

Socialist legacy (cont)		
	1960 Non-AG	1960 AG
UKRAINE		354

abandoned	1084	844
ROMANIA		1570
not abandoned	153	219
abandoned	586	612
HUNGARY		567
not abandoned	228	168
abandoned	112	59
SLOVAKIA		605
not abandoned	340	111
abandoned	135	19

not abandoned	40	36
abandoned	144	134
CZECH REPUBLIC		285
not abandoned	159	22
abandoned	89	15
POLAND		43
not abandoned	14	6
abandoned	18	5

Appendix 4.3 Model utility and performance (AUC) for three legacy models for the Carpathian region as a whole and by country

	Habsburg legacy	Habsburg vs. Socialist legacy	Soviet legacy
Overall	0.82	0.83	0.80
RO	0.76	0.75	0.68
HU	0.77	0.76	0.71
SK	0.81	0.81	0.80
UA	0.76	0.64	0.68
CZ	0.81	0.81	0.75
PL	0.71	0.72	0.91

Appendix 4.4 Variables selected in socialist abandonment models and sign of the coefficient indicating the direction of the effect

	Socialist abandonment (1960-1985)	Post-socialist abandonment (1985-2010)
Intercept	(-)	(-)
Historic agriculture	(-)	(-)
Slope	(+)	(+)
Precipitation	(+)	(+)
Crop suitability	(-)	(-)
Length of growing season	()	(+)
Distance to border	(-)	(-)
Distance to settlement	(-)	()
Country	(***)	(***)
AUC	0.787	0.82

Appendix 4.5 Variables selected in post-socialist abandonment models and sign of the coefficient indicating the direction of the effect.

	Habsburg legacy	Habsburg vs. Socialist legacy	Socialist legacy
Intercept	(-)	(-)	(-)
Historic agriculture	(-)	(-)	(-)
Slope	(+)	(+)	(+)
Precipitation	(+)	(+)	(+)
Crop suitability	(-)	(-)	(-)
Length of growing season	(+)	(+)	
Accessibility			(+)
Distance to border	(-)		
Distance to river			(+)
Population (1990)			
Country	(***)	(***)	(***)

Chapter 5. Conservation recommendations for the protection of bird habitat in the Carpathian Ecoregion in light long-term land use trends

Introduction

European landscapes are changing rapidly due to agricultural intensification, reforestation and land abandonment (Estel et al., 2015; Levers et al., 2014; Munteanu et al., 2014). The resulting habitat loss (Newbold et al., 2015), landscape fragmentation (Jaeger et al., 2011; Jongman, 2002), make these changes a primary concern for biodiversity conservation (Balmford et al., 2002; Green et al., 2005). At European scale, bird populations are declining (Donald et al., 2001; Gregory et al., 2007). In agricultural systems the decline is caused by intensification and changes in the Common Agricultural Policy (Donald et al., 2002, 2001). Forest birds are in decline due to loss of nesting habitat and reduced forest health (Gregory et al., 2007). Grassland birds declined due to agricultural activity, invasive species and habitat alteration (Busche, 1994; Skorka et al., 2010). However, there is also an opposite trend, where land use is becoming less intensive in some landscapes (Alcantara et al., 2013; Baumann et al., 2011) offering opportunities for habitat recovery (Ceașu et al., 2015a; Navarro and Pereira, 2012), and local increases of some forest birds have occurred in Eastern Europe (Reif et al., 2008). When land use becomes less intensive, the question for conservation is whether it is better to push for a full rewilding (Navarro and Pereira, 2012), or for the conservation of low-intensity land use practices, which can have conservation benefits in their own right (Fischer et al., 2012). This trade-off among different conservation goals is particularly relevant in regions with widespread agricultural land abandonment.

Globally, most land use changes are due to intensifying land use, and two conservation strategies have been proposed to balance conservation and food production: land sharing (or wildlife

friendly farming), where food and biodiversity are sustained in the same places at low intensity; and land sparing, where high-yield farming in one place allows land elsewhere to be retained as natural habitat (Fischer et al., 2014; Grau et al., 2013; Phalan et al., 2011). Areas where low-intensity agriculture has been practiced for extended periods of time are considered ideal for wildlife-friendly farming (Hartel et al., 2013; Mikulcak et al., 2013; Plieninger et al., 2006), because these landscapes preserve high levels of farmland diversity (Kleijn et al., 2009; Tschamntke et al., 2012). On the other hand: land sparing relies on separating land for conservation from land for crops, with high-yield farming in one place facilitating the protection of remaining natural habitats such as contiguous tracts of forest elsewhere (Phalan et al., 2011). In most cases a combination of sparing and sharing may be most appropriate where land use is intensifying.

In areas where land use is becoming less intensive, the arguments in favor of land sparing may not be relevant though. It may be highly beneficial for conservation to pay farmers to forgo intensive land use practices, and retain traditional farming practices. However, retaining such practices may not be best for conservation when the alternative is rewilding. Indeed, remote regions of Europe, such as the Carpathian Mountains, have been suggested for rewilding, due to their high rate of agricultural abandonment, contiguous forest ecosystems, high mammal and avian biodiversity and unique presence of large carnivore and herbivore species (Ceașu et al., 2015a, 2015b; Navarro and Pereira, 2012). The question is though whether such a rewilding, which could greatly benefit forest species, should be the goal for conservation given the trade-off of a likely loss of habitat for farmland and grassland species.

The question of what conservation goals are effective in places with decreasing land use intensity is not a theoretical one. The European Union supports substantial programs to address

biodiversity conservation, and they include both preserving low-intensity, high-natural-value farmland via agri-environmental payments (€4.44 bln/ year between 2007-2012) (European Court of Auditors, 2011) and protecting species and habitats under the Natura 2000 conservation program (estimated €5.80 bln/ year in 2012) (The European Commission, 2014). Yet these efforts are rarely allocated considering the long-term land changes in a particular region or the responsibility of geographic areas to the overall persistence and survival of a given species (Schmeller et al., 2012, 2008). Historical land uses can affect ecosystem structure, functioning and ultimately biodiversity levels (Brudvig et al., 2013; Dullinger et al., 2013; Plue et al., 2009). Furthermore, land use legacies can restrict the pace at which contemporary land change is possible (Munteanu et al., 2015), which may reduce the effectiveness of agri-environmental payment schemes. Given a species' proportional distribution, relative abundance or conservation status, certain geographic or administrative regions may carry higher conservation responsibility than others for that species global population (Keller and Bollman, 2004; Keller and Bollmann, 2001). This is why it is essential that, for instance agri-environmental payments are directed towards regions which carry high conservation responsibility for farmland species.

Empirical evidence suggest that bird populations in Eastern Europe are declining at slower rates than in Western Europe (Gregory et al., 2007; Skorka et al., 2010; Verhulst et al., 2004), likely due to lower intensity land uses compared to the rest of Europe (Gregory et al., 2007). This means that Eastern Europe may be a future hotspot for bird diversity conservation in Europe. Despite local evidence on which species could benefit most from a land sharing or rewilding, the historical landscape context and the conservation responsibility of the Carpathians at broad spatial scales remain largely unexplored, leading to potentially ill-informed conservation actions.

Our overarching goal was to provide conservation recommendations for the Carpathian Mountains, based on historical and recent habitat evolution for species of highest conservation responsibility at European level. Specifically, we asked:

- 1) How did habitat for different bird species change in the Carpathian Mountains over the past 150 years?
- 2) For which of the bird species do the Carpathians carry the highest conservation responsibility at the European level?
- 3) What bird conservation strategies should be pursued, and what future land use trends would be most desirable for conservation, given the major threats to different species, their past habitat trends and the conservation responsibility of the Carpathians?

Methods

Study area

We studied the Carpathian Ecoregion (207,309 km²) which is the largest mountain range and a conservation hotspot in Europe due to its valuable forest ecosystems (Knorn et al., 2012), diverse cultural landscapes (Kozak et al., 2013) and high levels of biodiversity (Akeroyd and Page, 2011; Kuemmerle et al., 2010; Pereira and Navarro, 2015). The Carpathians are part of six European countries, and experienced numerous shifts in land use over the last century, due to major changes in socio-economic and political conditions (Munteanu et al., 2015), increasing forest cover and agricultural abandonment being the most prominent changes (Munteanu et al., 2014). The contemporary land cover consists of forests (58%), agricultural fields (9%), grasslands (30%), and scattered settlements, but differs considerably from historical land use patterns. For example, forest cover experienced its lowest extent during the 1930s, when only 46% of the Carpathians were forested and agricultural use experienced a high during the 1960s,

when 27% of the region was arable land (Figure 5.1). The main agricultural crops are cereals (wheat, corn, barley) and legumes (potatoes, sugar beets), and they are mostly scattered across the landscape as small subsistence farms (Griffiths et al., 2013). Grasslands consist mostly of pastures, hay meadows and wooded pastures, and preserve a high level of biodiversity due to their low intensity use (Akeroyd and Page, 2011; Halada et al., 2008; Hartel et al., 2013). Despite the fact that the region has experienced high forest disturbance followed by spruce plantings during and after the Socialist regime (Griffiths et al., 2014; Munteanu et al., 2015), tree species diversity is high in the Carpathians. At low elevation, deciduous woodlands (*Quercus* sp, *Fagus sylvatica*, *Carpinus betulus*, *Populus* sp, and *Robinia pseudoacacia*) are common, while at high elevations, coniferous forests are dominant (*Pinus* sp, *Picea abies*, *Abies alba*) (Munteanu et al., 2015). The Carpathians have an increasing network of protected areas, some of which are aimed at conserving habitat for bird species (under the NATURA 2000 initiative), with highly variable effectiveness (Butsic et al., 2015).

Data

We analyzed species range maps for 252 bird species (BirdLife International and NatureServe, 2014), as well as their attributes including IUCN Red List status, population trend at European level from 1990 to 2000, European level threats to the populations survival, and major habitat (i.e., the primary habitat used by the species for breeding or feeding; (Birdlife International, 2016). Habitat data was available for 170 species.

We reconstructed forest, grassland and agricultural land use throughout the Carpathian Ecoregion for six points in time: 1860s, 1930s, 1960s, 1985, 2000 and 2010. We used a regular 2-km point sampling grid according to the 2007 INSPIRE directive (Infrastructure for Spatial

Information in the European Community) and LUCAS (Land Use and Cover Area frame Survey, (Gallego and Delince', 2010) and assigned one of the three cover types to each grid point at each time point. We obtained land use data for the 1860s, 1930s and 1960s from historical topographic maps (Munteanu et al., 2016a, 2015) and land cover data for the years 1985, 2000 and 2010 from classified Landsat TM/ETM+ image composites (Griffiths et al., 2014, 2013). In total, we had 51,648 data points. Due to low accuracy of historic maps in parts of Romania and Hungary, for a total of 3,409 points, we could not distinguish agricultural from grassland land use for the 1860s. For these points, we assigned the 1930s land use class for the 1860s as well (Munteanu et al., 2016a). For comparability, we reclassified the major habitat data for each species (BirdLife International and NatureServe, 2014; Birdlife International, 2016) into three categories that matched our long term land cover data: agriculture, grasslands and forests.

Analysis

To determine the potential habitat change for each bird species, we analyzed data points that matched the spatial extent of each species' range and the major habitat types used by each species. The resulting points represented the 'potential habitat' for each species, for the years 1860s, 1930s, 1960s, 1985, 2000 and 2010. Based on this, we mapped and analyzed the dynamics of forest, agricultural and grassland change between 1860 and 2010 within the range of all species present in the Carpathians. Although several species use other habitats as their major habitat (such as wetlands or open water), this data was not available for all time periods, which is why we did not include these habitat types in our analysis.

In order to determine the conservation responsibility of the Carpathians for each species, we considered three criteria: a) if a species had a high proportion of its European range in the

Carpathians, b) if a species had an IUCN conservation status of vulnerable (VU), near threatened (NT), endangered (EN) or critically endangered (CR) and c) if a species' population declined at the European level from 1990 to 2000. We assumed that the proportional distribution of a species is a proxy for the relative importance of the region for the species viability (Keller and Bollman, 2004; Schmeller et al., 2008). Because the Carpathian Mountains make up 3.106% of the total European landmass (including European Russia) we considered values less than 3.106% as low, values higher than 3.106% as medium and values over 6.212% as a high proportion of the species European range being in the Carpathians. We chose the value of 6.212% as the double of the Carpathians fraction of Europe, following prior studies for the Alps (Keller and Bollman, 2004; Keller and Bollmann, 2001). We defined four classes of conservation responsibility for the Carpathians:

HHCD: High conservation responsibility for the Carpathians because the species has: **high** percentage of the species range in the Carpathians ($> 6.212\%$), IUCN status of **concern** (VU, NT, EN, CR) and population at European level in **decline** or fluctuating between 1990 and 2000.

HHLC: High conservation responsibility for the Carpathians because the species has: **high** percentage of the species range in the Carpathians ($> 6.212\%$), IUCN status of **least concern** (LC) and population at European level either in decline or increasing.

HMCD: High conservation responsibility for the Carpathians because the species has: **medium** percentage of the species range in the Carpathians ($> 3.106\%$ and $< 6.212\%$), IUCN status of **concern** (VU, NT, EN, CR) and population at European level in **decline**.

LLLC: Low conservation responsibility for the Carpathians because the species has: **low** percentage of the species range in the Carpathians ($< 3.106\%$) or medium percentage of the

species range in the Carpathians (>3.106 and $< 6.212\%$) and IUCN status of least concerned (LC).

To understand which habitat types are important for species for which the Carpathians have high conservation responsibility (classes HHCD, HHLC and HMCD), we analyzed the percentage of species' European range in the Carpathians by species major habitats and in relation to the thresholds of 3.106% and 6.212% (medium and high percentage of species range situated in Carpathians). If a species uses more than one habitat type, we double counted. Finally, for species of high conservation responsibility (classes HHCD, HHLC and HMCD) we summarized all points of major habitat within their range, and analyzed potential habitat dynamics over time.

To understand what conservation strategies could be pursued in the Carpathians, we reviewed the overall species threats at European level in relation to species past habitat trends and the conservation responsibility of the Carpathians. For the species of high conservation responsibility, we analyzed the major threat classes for each species as described by IUCN Red List and Birdlife International (BirdLife International and NatureServe, 2014). If a listed threat would affect habitat and would operate at European level, but was not present in the Carpathians, then we assumed that region could make a substantial contribution to the conservation of that species. For example, if agricultural expansion represents the major threat to a species at European level, but in the Carpathians agricultural land was in decline, this may indicate high conservation potential for that species in the Carpathians. We counted the type of threats in each IUCN threat class and compared them to the potential habitat dynamics in order to identify land management options that could minimize the number of threats for most of the Carpathians species.

Results

We found that forest and grassland habitat increased substantially since 1860 within the ranges of all Carpathian species, while agricultural habitat declined. Overall, the Carpathians have a high conservation responsibility for species that use forests and grasslands as their major habitat and only low conservation responsibility for birds that rely on agriculture. Furthermore, several species of high conservation responsibility would benefit from a mosaic of forest and grassland landscapes. The main threats at European level for the species of high conservation responsibility were agricultural intensification and natural system modification. Because agricultural land declined substantially in our study region since the 1960s, we suggest that there is a high potential for conservation of these species exists in the Carpathians. We identified a list of 29 species for which the Carpathian ecoregion could become a conservation hotspot and suggest that land management should focus on providing suitable habitat for these species in the Carpathians.

When analyzing the evolution of potential habitat for all species between 1860 and 2010, we found a general increase in grassland and forest habitats, and a decrease in agricultural habitat (Figure 5.2). For a species whose range coincided with the Carpathian ecoregion, the forest made up 58% of its Carpathian range in 2010. The lowest forest extent occurred in the 1930s, and substantial increases were recorded thereafter in southern Poland, northern Slovakia and Ukraine (Figure 5.2). Agricultural area increased until the 1960s, when agriculture made up 27%, but abandonment rates were high afterwards so that by 2010, only 9% of the territory was farmed. Grassland expansion was widespread since 1960s, reaching the maximum extent in 2000 (32%) and decreasing by 2010 by 1%, due to reforestation and some agricultural recultivation. Of the species for which habitat data was available (170), 39% used forest as their major habitat, 27%

grasslands, and only 2% (4 species) agriculture (Figure 5.2). When mapping major habitat dynamics for all species, we found that for all species agricultural habitats declined since 1860, while forest and grassland habitat increased. By 2010, 63% of the land farmed in 1860 was abandoned, grasslands increased by 38% and forest by 19%.

The Carpathian Mountains represent 3.106% of the European territory (including European Russia). However, for 169 of the 252 species present in the Carpathians, the region made up more than 3.106% of their range, indicating that the Carpathians carry conservation responsibility at the European level for 67% of the species present there. Furthermore, for 7% of all species (19 species) the Carpathians made up more than 6.212 % of their European range. For five species, the Carpathians represented more than 10% of their European range: corn bunting (*Miliaria calandria*, 20.6%), Ural owl (*Strix uralensis*, 12.8%), collared flycatcher (*Ficedula albicollis*, 11.7%), wallcreeper (*Tichodroma muraria*, 11.6%) and spotted nutcracker (*Nucifraga caryocatactes*, 10.1%). Most species with a high or medium proportion of their range in the Carpathians were either forest or grassland species, and none were agricultural species (Figure 5.3).

Of the 252 bird species present in the Carpathians, 16 species were IUCN Red listed as either vulnerable (VU), near threatened (NT), critically endangered (CR), or endangered (EN). Of these, the Eastern Imperial Eagle (*Aquila heliaca*) and the lesser white-fronted goose (*Anser erythropus*) had particularly high portions of their ranges in the Carpathians (8.1 and 6.4% respectively).

When analyzing the European level population dynamics between 1990 and 2000 for the 252 species, we found that 33.7% (85 species) of the species declined at the European level and

further 10% missed data on their population trend. Of the 85 species, 55 had a high percentage of their range in the Carpathians, and 12 were of conservation concern (VU or NT, Table 5.1).

Overall, we identified 29 species of high conservation responsibility according to the proportion of their habitat in the Carpathians, their population trend, and their IUCN red list status (i.e., they belonged to either HHCD, HHLC, HMCD). The HHCD responsibility class included two species with IUCN status of concern and a high percentage of their range in the Carpathians: the Eastern imperial eagle and the lesser white-fronted goose. The HHLC responsibility class included a total of 17 species with varying population trends and the HMCD responsibility class included 10 species of conservation concern that had a moderate proportion of their range in the Carpathians (Table 5.1 and Appendix 5.1).

Most of the 29 high conservation responsibility species require forest habitats (13 species), followed by grasslands (11), and other habitats (including wetlands, water and settlement areas, 9 species). This result is consistent with the habitat requirements of all species in the Carpathians. For four species habitat requirements were unknown: Syrian woodpecker, corn bunting, wallcreeper, and red-footed falcon, and eight species required more than one habitat. For example, the Eastern imperial eagle relies on forest for nesting and perching and on open wetlands for hunting (Table 5.2).

When analyzing potential habitat dynamics of the species of high conservation responsibility, we found for all species an overall increase of their potential habitat since 1860, and particularly fast increases after 1960. A few species registered a small decline in potential habitat since 2000, due to grassland conversions (Figure 5.4). Of the species of high conservation responsibility in the Carpathians, lesser spotted eagle (*Clanga pomarina*), collared flycatcher (*Ficedula albicollis*),

ring ouzel (*Turdus torquatus*), and red kite (*Milvus milvus*) require forest and grasslands habitat for their survival. All species in the HHLC conservation responsibility category are major forest and/or grasslands species (except the corn crake, which is also a wetland species). In contrast, most of the HMCD conservation responsibility species require grassland or wetland habitat. The Carpathians carry minimal responsibility for species that use agriculture though because no species of high conservation responsibility have agricultural land as their major habitat.

When analyzing the major threats at the European level to the species of high conservation responsibility in the Carpathians, the most frequent threats fell in two categories: either agriculture and aquaculture, or natural system disturbance (Figure 5.5). Farming, grazing and wood extraction were the most common issues for the 29 species. Twelve species were threatened by agricultural intensification and herbicide use, two by wood harvesting, and for thirteen species, no information on threats was available (Table 5.3). Lesser grey shrike (*Lanius minor*), snow bunting (*Plectrophenax nivalis*) and ring ouzel were more affected by climate change than by habitat change (Figure 5.5).

Discussion

Regions where land use intensity is decreasing, such as the Carpathians may offer opportunities for conservation, but the question is whether conservation efforts should aim for rewilding or for the maintenance of low-intensity land use, and traditional agricultural practices. The answer to this question requires to consider long term habitat dynamics, the conservation responsibility of a given region to a species' population at large (Keller and Bollmann, 2001; Schmeller et al., 2012), and the threats to this species. Here, we assessed long-term habitat dynamics for Carpathian bird species and identified for which of those species the Carpathians carry a high conservation responsibility. We found the highest conservation responsibility for forest and

grassland species. None of the species which use agricultural habitats are of conservation responsibility in this region. The potential habitat of species of high conservation responsibility increased in the Carpathians since the 1860s, and many of the European level threats such as agricultural intensification are not affecting species here.

Our results suggest that conservation action should be directed towards forest specialists and forest generalist species (classes HHCD and HHLC), because they are of high conservation responsibility and their habitat has increased since 1860s in the Carpathians. This includes woodpeckers, Ural owl, and hazel grouse, all of which require structurally heterogeneous forests for survival. Ural owls prefer large contiguous tracts of forest, with high proportion of deciduous and old trees (Kajtoch et al., 2015). Hazel grouse require dense understory and prefer alder (Åberg et al., 2003; Schaublin and Bollmann, 2011). Three toed woodpecker and middle spotted woodpecker require old trees and snags (Löhmus, 2003; Roberge et al., 2008). In general, woodpeckers are great indicators for forest bird diversity in Eastern Europe (Mikusiński et al., 2001), suggesting that conserving species in the HHLC class will benefit many species in the Carpathians.

We caution though that the conservation of forest birds will depend greatly on the type of forests, and their structure. The Carpathians are one of the last strongholds of old-growth forests in Europe (Kozak et al., 2013; Munteanu et al., 2015), and potential habitat for forest species has increased since 1860s. However, intense forest harvest in the Carpathians has diminished old-growth forests both in the early 20th century (Munteanu et al., 2016b) and in recent decades (Knorn et al., 2012). Furthermore, forest composition shifted towards coniferous forests since the 1860s due to large-scale clear-cuts and spruce and pine monoculture (Griffiths et al., 2014; Munteanu et al., 2015). These changes very likely affect HHCD and HHLC responsibility

species. Forest management should thus aim to restore tree species diversity and structural forest diversity. Although IUCN threat data is missing for many forest specialist species in our analysis, the lack of nest sites, large trees, and understory vegetation may negatively affect populations of woodpeckers, hazel grouse, and Ural owls (Åberg et al., 2003; Lõhmus, 2003; Roberge et al., 2008).

Several of the species in our analysis require both woodland plus either grassland or wetlands. One example is the high conservation responsibility (HHCD) Eastern imperial eagle, which prefers wooded grasslands, forest edges and open wetland habitat (Hallmann, 1996; Kovács et al., 2008). The imperial eagle population in the Carpathians was very small in the 19th and early 20th century (Salmen, 1980), and restricted to high altitudes, but increased since 2000, especially in Slovakia, Hungary, and the Czech Republic (Demerdzhiev et al., 2011), which may be related to habitat increases since the 1860s (Figure 5.4). The biggest European-level threats to the eastern imperial eagle are agricultural intensification and natural habitat loss, which caused substantial population declines in Greece (Hallmann, 1996). However, in the Carpathians agricultural abandonment is widespread and large-scale agriculture uncommon, which is why we suggest that the region is well suited for the protection this species. Furthermore, red kites, lesser spotted eagles, and collared flycatchers have similar habitat requirements and threats as the imperial eagle. All of these species inhabit a combination of woodlands and open spaces with large trees, and some also require cavities for nesting (such as the flycatcher) (Salmen, 1980). Our results indicated that both forest and grassland habitat for these species increased greatly since the 1960s in the Carpathians. The Transylvanian plain in particular could represent highly suitable landscape for these species due to its mosaic of woodlands, grasslands, and agricultural patches, whereas the more contiguous forests of the Carpathian range are probably less suitable.

Our results also suggest that grassland birds should be a focus of conservation actions in the Carpathians because the region has high conservation responsibility for many grassland species, and witnessed generally an increase in grassland habitats over the past century, but this trend was reversed after 2000. Most of the grassland species of high conservation responsibility (classes HMCD and HHLC) are declining across Europe, such as corn crake, lesser grey shrike, and ring ouzel. Corn crakes use managed and unmanaged grasslands as well as riverine floodplains (Elts, 1997), and its populations declined in the Carpathians after the early 1900s when agricultural expansion started in Transylvania (Salmen, 1980). The decline persisted until the middle of the 20th century (Salmen, 1980) consistent with our potential habitat trends (Figure 5.4). The lesser grey shrike is also a grassland specialist, but requires scattered groups of trees and small patches of woodlands for nesting (Lovász et al., 2000). At the beginning of the 20th century, lesser grey shrike was the most numerous shrike species in Eastern Europe, but European populations decreased after the 1970s (Lovász et al., 2000) due to intensifying agriculture (Kvist et al., 2011). We found increasing of potential habitat since 1985 for the lesser grey shrike. The ring ouzel uses grassland and shrubby vegetation for nesting (Sim et al., 2007). We found an increase in the ring ouzel's potential habitat in the Carpathians due to agricultural abandonment. We caution though that due to EU agricultural policies, some cropland fields have been recently re-cultivated in the Carpathians (Griffiths et al., 2013) potentially posing threats to grassland species such as pesticide use, and loss of nesting habitat.

One reason why the Carpathians provide more habitat for so many forest and grassland species is that agricultural land declined in the Carpathians since 1985, due to the collapse of the collective farming system and removal of subsidies received under socialist regimes (Griffiths et al., 2013; Hartvigsen, 2014; Kuemmerle et al., 2008). At the European level, the majority of the threats to

the Carpathians' species of conservation responsibility are related to agricultural intensification, and herbicide use. Agricultural expansion, intensification or pesticide use are not widespread in the Carpathians though, with the exception of some agricultural re-cultivation in the most fertile areas (Griffiths et al., 2013).

Overall, three broad conservation messages emerge from our analysis: 1) The Carpathians carry the greatest conservation responsibility for forest and grassland species, and hence, conservation efforts should focus on these species and their habitats. 2) The conservation responsibility of the Carpathians for agricultural species is very low, which is why we suggest that cropland abandonment is a positive trend in the Carpathians, as long as grasslands persist. 3) The observed increase in potential habitat provides a great opportunity for conservation, but conservation efforts need ensure that forest diversity, and structural elements such as dead wood and snags, are maintained.

Ultimately, we suggest that maintaining both forested areas and grasslands in the Carpathians would be best for bird conservation in the Carpathians, but supporting row-crop agriculture is not. Spatially, the forested areas should be located at higher elevations and rewilding would benefit species in the mountainous region (Pereira and Navarro, 2015). Grasslands should be maintained in Transylvania and in the larger river valleys, where wildlife friendly livestock farming could contribute to the conservation of grassland species (Hartel et al., 2013; Mikulcak et al., 2013). In addition to grassland birds, protecting grasslands would also benefit other taxa, including plants and butterflies, which can exhibit high species richness in Carpathian grasslands (Cremene et al., 2005; Loos et al., 2014; Schmitt and Rákossy, 2007). Furthermore, the large river valleys may offer opportunities for wetland restoration, and hence benefit wetland species (Moga and Ollerer, 2007).

Conclusions

The conservation goals that arise from our results, i.e., to focus on forest and grassland bird species, could be achieved by allowing for natural succession on all land that was forested in 1860s or thereafter, and allowing the remaining agricultural lands to convert to grasslands. If such management was implemented, the Carpathians would consist of 66% forest cover and 32% grassland, which would benefit all species of conservation concern. Furthermore, this management plan would require a 8% increase in forest cover and a 2% increase in grasslands, rates of change well within historic ranges of change in the Carpathians, and consistent with recent land use trends (see Appendix 5.2 for visualization of this landscape).

The current trend of agricultural abandonment in the Carpathian Mountains is positive for avian diversity, and we see little justification for EU subsidies for row-crop farming from a conservation perspective. The Transylvanian plains were historically mostly un-managed grassland landscapes. We suggest that in those areas, low-intensity livestock management could enhance conservation outcomes.

Decreasing land use intensity provides great opportunities for conservation in the Carpathians and elsewhere. However, it is important to evaluate if conservation actions designed for regions where land use is intensifying, such as land sharing, are best for regions where land use intensity is declining, and efforts to maintain land sharing may prevent rewilding. The approach that we developed here for one region of high conservation importance where land use intensity has declined substantially could easily be applied elsewhere.

References

- Åberg, J., Swenson, J.E., Angelstam, P., 2003. The habitat requirements of hazel grouse (*Bonasa bonasia*) in managed boreal forest and applicability of forest stand descriptions as a tool to identify suitable patches. *For Ecol Manage* 175, 437–444.
- Akeroyd, J.R., Page, J.N., 2011. Conservation of High Nature Value (HNV) grassland in a farmed landscape in Transylvania, Romania. *Contrib Bot* XLVI, 57–71.
- Alcantara, C., Kuemmerle, T., Baumann, M., Bragina, E. V., Griffiths, P., Hostert, P., Knorn, J., Müller, D., Prishchepov, A. V., Schierhorn, F., Sieber, A., Radeloff, V.C., 2013. Mapping the extent of abandoned farmland in Central and Eastern Europe using MODIS time series satellite data. *Environ Res Lett* 8, 035035.
- Balmford, A., Bruner, A., Cooper, P., Costanza, R., Farber, S., Green, R.E., Jenkins, M., Jefferiss, P., Jessamy, V., Madden, J., Munro, K., Myers, N., Naeem, S., Paavola, J., Rayment, M., Rosendo, S., Roughgarden, J., Trumper, K., Turner, R.K., 2002. Economic reasons for conserving wild nature. *Science* 297, 950–3.
- Baumann, M., Kuemmerle, T., Elbakidze, M., Ozdogan, M., Radeloff, V.C., Keuler, N.S., Prishchepov, A. V., Kruhlov, I., Hostert, P., 2011. Patterns and drivers of post-socialist farmland abandonment in Western Ukraine. *Land use policy* 28, 552–562.
- Birdlife International, 2016. Birdlife Data Zone. Terms and definitions [WWW Document]. URL <http://www.birdlife.org/datazone/info/spchabalt> (accessed 1.1.16).
- BirdLife International and NatureServe, 2014. Bird species distribution maps of the world.
- Brudvig, L.A., Grman, E., Habeck, C.W., Orrock, J.L., Ledvina, J.A., 2013. Strong legacy of agricultural land use on soils and understory plant communities in longleaf pine woodlands. *For Ecol Manage* 310, 944–955.
- Busche, G., 1994. The decline of wet-meadow birds in Schleswig-Holstein/Germany from 1950 to 1992. *J fur Ornithol* 135, 167–177.
- Butsic, V., Munteanu, C., Griffiths, P., Knorn, J., Radeloff, V.C., Lieskovský, J., Müller, D., Kuemmerle, T., 2016. The impact of protected areas on forest disturbance in the Carpathian Mountains 1985-2010. *Conserv Biol* (in review).
- Ceaușu, S., Carver, S., Verburg, P.H., Kuechly, H.U., Hölker, F., Brotons, L., Pereira, H.M., 2015a. Rewilding European Landscapes 25–46.
- Ceaușu, S., Hofmann, M., Navarro, L.M., Carver, S., Verburg, P.H., Pereira, H.M., 2015b. Mapping opportunities and challenges for rewilding in Europe. *Conserv Biol*. 00, 1–11
- Cremene, C., Groza, G., Rakosy, L., Schilenyko, A.A., Baur, A., Erhardt, A., Baur, B., 2005. Alterations of Steppe-Like Grasslands in Eastern Europe: a Threat to Regional Biodiversity Hotspots. *Conserv Biol* 19, 1606–1618.
- Demerdzhiev, D., Horváth, M., Kovács, A., Stoychev, S., Karyakin, I., 2011. Status and

- population trend of the eastern imperial eagle (*Aquila heliaca*) in Europe in the period 2000–2010. *Acta Zool Bulg* 63, 5–14.
- Donald, P.F., Gree, R.E., Heath, M.F., 2001. Agricultural intensification and the collapse of Europe's farmland bird populations. *Proc Biol Sci* 268, 25–9.
- Donald, P.F., Pisano, G., Rayment, M.D., Pain, D.J., 2002. The Common Agricultural Policy, EU enlargement and the conservation of Europe's farmland birds. *Agric Ecosyst Environ* 89, 167–182.
- Dullinger, S., Essl, F., Rabitsch, W., Erb, K.-H., Gingrich, S., Haberl, H., Hülber, K., Jarosik, V., Krausmann, F., Kühn, I., Pergl, J., Pysek, P., Hulme, P.E., 2013. Europe's other debt crisis caused by the long legacy of future extinctions. *Proc Natl Acad Sci USA* 110, 7342–7.
- Eltis, J., 1997. Studies of the Corncrake in Estonia in 1995. *Vogelwelt* 118, 236–238.
- Estel, S., Kuemmerle, T., Alcántara, C., Levers, C., Prishchepov, A., Hostert, P., 2015. Mapping farmland abandonment and recultivation across Europe using MODIS NDVI time series. *Remote Sens Environ* 163, 312–325.
- European Court of Auditors, 2011. Is agri-environment support well designed and managed?, Reproduction.
- Fischer, J., Abson, D.J., Butsic, V., Chappell, M.J., Ekroos, J., Hanspach, J., Kuemmerle, T., Smith, H.G., von Wehrden, H., 2014. Land sparing versus land sharing: moving forward. *Conserv Lett* 7.
- Fischer, J., Hartel, T., Kuemmerle, T., 2012. Conservation policy in traditional farming landscapes. *Conserv Lett* 5, 167–175.
- Gallego, J., Delincé, J., 2010. The European land use and cover area-frame statistical survey, in: Benedetti, R., Bee, M., Espa, G., Piersimoni, F. (Eds.), *Agricultural Survey Methods*. John Wiley & Sons, pp. 151–168.
- Grau, R., Kuemmerle, T., Macchi, L., 2013. Beyond “land sparing versus land sharing”: environmental heterogeneity, globalization and the balance between agricultural production and nature conservation. *Curr Opin Environ Sustain* 5, 477–483.
- Green, R.E., Cornell, S.J., Scharlemann, J.P.W., Balmford, A., 2005. Farming and the fate of wild nature. *Science* 307, 550–5.
- Gregory, R.D., Vorisek, P., Van Strein, A., Gmelig Meyling, A., Jiguet, F., Fornasari, L., Reif, J., Chylatrecki, P., Burfield, I.J., 2007. Population trends of widespread woodland birds in Europe. *Ibis (Lond 1859)* 149, 78–97.
- Griffiths, P., Kuemmerle, T., Baumann, M., Radeloff, V.C., Abrudan, I. V., Lieskovský, J., Munteanu, C., Ostapowicz, K., Hostert, P., 2014. Forest disturbances, forest recover, and changes in forest types across the Carpathian ecoregion from 1985 to 2010 based on Landsat image composites. *Remote Sens Environ* 151, 72–88.

- Griffiths, P., Müller, D., Kuemmerle, T., Hostert, P., 2013. Agricultural land change in the Carpathian ecoregion after the breakdown of socialism and expansion of the European Union. *Environ Res Lett* 8, 045024.
- Halada, L., Ružičková, H., David, S., Halabuk, A., 2008. Semi-natural grasslands under impact of changing land use during last 30 years: *Trollio-Cirsietum* community in the Liptov region (N Slovakia). *Community Ecol* 9, 115–123.
- Hallmann, B., 1996. The decline of the Imperial Eagle *Aquila heliaca* in Greece. *Eagle Stud* 439–442.
- Hartel, T., Dorresteyn, I., Klein, C., Máthé, O., Moga, C.I., Öllerer, K., Roellig, M., von Wehrden, H., Fischer, J., 2013. Wood-pastures in a traditional rural region of Eastern Europe: Characteristics, management and status. *Biol Conserv* 166, 267–275.
- Hartvigsen, M., 2014. Land reform and land fragmentation in Central and Eastern Europe. *Land use policy* 36, 330–341.
- Jaeger, J.A.G., Soukup, T., Madriñán, L.F., Schwick, C., Kienast, F., 2011. Landscape fragmentation in Europe., EEA Report Series. European Environment Agency, Brussels.
- Jongman, R.H.G., 2002. Homogenisation and fragmentation of the European landscape: ecological consequences and solutions. *Landsc Urban Plan* 58, 211–221.
- Kajtoch, Ł., Żmihorski, M., Wieczorek, P., 2015. Habitat displacement effect between two competing owl species in fragmented forests. *Popul Ecol* 57, 517–527.
- Keller, V., Bollman, K., 2004. From Red List to species of Conservation Concern. *Conserv Biol* 18, 1636–1644.
- Keller, V., Bollmann, K., 2001. Fuer welche Vogelarten traegt die Schweiz eine besondere Verantwortung? *Ornithol Beobachter* 98, 323–340.
- Kleijn, D., Kohler, F., Báldi, A., Batáry, P., Concepción, E.D., Clough, Y., Díaz, M., Gabriel, D., Holzschuh, A., Knop, E., Kovács, A., Marshall, E.J.P., Tschamntke, T., Verhulst, J., 2009. On the relationship between farmland biodiversity and land-use intensity in Europe. *Proc Biol Sci* 276, 903–9.
- Knorn, J., Kuemmerle, T., Radeloff, V.C., Keeton, W.S., Gancz, V., Biriş, I.-A., Svoboda, M., Griffiths, P., Hagatis, A., Hostert, P., 2012. Continued loss of temperate old-growth forests in the Romanian Carpathians despite an increasing protected area network. *Environ Conserv* 40, 182–193.
- Kovács, A., Demeter, I., Fatér, I., Bagyura, J., Nagy, K., Szitta, T., Firmánszky, G., Horváth, M., 2008. Current Efforts to Monitor and Conserve the Eastern Imperial Eagle *Aquila heliaca* in Hungary. *AMBIO A J Hum Environ* 37, 457–459.
- Kozak, J., Ostapowicz, K., Bytnerowicz, A., Wyżga, B. (Eds.), 2013. The Carpathians: Integrating Nature and Society Towards Sustainability, Environmental Science and

- Engineering. Springer Berlin Heidelberg, Berlin, Heidelberg.
- Kuemmerle, T., Hostert, P., Radeloff, V.C., van der Linden, S., Perzanowski, K., Kruhlov, I., 2008. Cross-border Comparison of Post-socialist Farmland Abandonment in the Carpathians. *Ecosystems* 11, 614–628.
- Kuemmerle, T., Perzanowski, K., Chaskovskyy, O., Ostapowicz, K., Halada, L., Bashta, A.-T., Kruhlov, I., Hostert, P., Waller, D.M., Radeloff, V.C., 2010. European Bison habitat in the Carpathian Mountains. *Biol Conserv* 143, 908–916.
- Kvist, L., Giralt, D., Valera, F., Hoi, H., Kristin, A., Darchiashvili, G., Lovaszi, P., 2011. Population decline is accompanied by loss of genetic diversity in the Lesser Grey Shrike *Lanius minor*. *Ibis (Lond 1859)* 153, 98–109.
- Levers, C., Verkerk, P.J., Müller, D., Verburg, P.H., Butsic, V., Leitão, P.J., Lindner, M., Kuemmerle, T., 2014. Drivers of forest harvesting intensity patterns in Europe. *For Ecol Manage* 315, 160–172.
- Löhmus, A., 2003. Do Ural owls (*Strix uralensis*) suffer from the lack of nest sites in managed forests? *Biol Conserv* 110, 1–9.
- Loos, J., Dorresteyn, I., Hanspach, J., Fust, P., Rakosy, L., Fischer, J., 2014. Low-intensity agricultural landscapes in Transylvania support high butterfly diversity: implications for conservation. *PLoS One* 9, e103256.
- Lovászi, P., Bártol, I., Moskát, C., 2000. Nest-site selection and breeding success of the Lesser grey shrike (*Lanius minor*) in Hungary. *Ring* 22, 157–164.
- Mikulcak, F., Newig, J., Milcu, A.I., Hartel, T., Fischer, J., 2013. Integrating Rural Development and Biodiversity Conservation in Central Romania. *Environ Conserv* 40, 129–137.
- Mikusiński, G., Gromadzki, M., Chylarecki, P., 2001. Woodpeckers as Indicators of Forest Bird Diversity. *Conserv Biol* 15, 208–217.
- Moga, C.I., Öllerer, K., 2007. The importance of the riparian forest habitat for bird species richness in the Tarnava Mare Valley (Transylvania, Romania). *Transylvanian Rev Syst Ecol Res* 4, 179–186.
- Munteanu, C., Kuemmerle, T., Boltiziar, M., Butsic, V., Gimmi, U., Kaim, D., Király, G., Konkoly-Gyuró, É., Kozak, J., Lieskovský, J., Mojses, M., Müller, D., Ostafin, K., Ostapowicz, K., Shandra, O., Štych, P., Walker, S., Radeloff, V.C., 2014. Forest and agricultural land change in the Carpathian region—A meta-analysis of long-term patterns and drivers of change. *Land use policy* 38, 685–697.
- Munteanu, C., Kuemmerle, T., Boltiziar, M., Lieskovský, J., Mojses, M., Kaim, D., Konkoly-Gyuró, É., Mackovčin, P., Müller, D., Ostapowicz, K., Radeloff, V.C., 2016a. Land use legacies of historic political regimes for contemporary land abandonment. *Reg Environ Chang* in preparation.
- Munteanu, C., Kuemmerle, T., Keuler, N.S., Müller, D., Balazs, P., Dobosz, M., Griffiths, P.,

- Halada, L., Kaim, D., Király, G., Konkoly-Gyuró, É., Kozak, J., Lieskovský, J., Ostafin, K., Ostapowicz, K., Shandra, O., Radeloff, V.C., 2015. Legacies of 19th century land use shape contemporary forest cover. *Glob Environ Chang* 34, 83–94.
- Munteanu, C., Nita, M.-D., Abrudan, I. V., Radeloff, V.C., 2016b. Historical forest management in Romania is imposing strong legacies on contemporary forests and their management. *For Ecol Manage* 361, 179–193.
- Navarro, L.M., Pereira, H.M., 2012. Rewilding Abandoned Landscapes in Europe. *Ecosystems* 15, 900–912.
- Newbold, T., Hudson, L.N., Hill, S.L.L., Contu, S., Lysenko, I., Senior, R.A., Börger, L., Bennett, D.J., Choimes, A., Collen, B., Day, J., De Palma, A., Díaz, S., Echeverria-Londoño, S., Edgar, M.J., Feldman, A., Garon, M., Harrison, M.L.K., Alhusseini, T., Ingram, D.J., Itescu, Y., Kattge, J., Kemp, V., Kirkpatrick, L., Kleyer, M., Correia, D.L.P., Martin, C.D., Meiri, S., Novosolov, M., Pan, Y., Phillips, H.R.P., Purves, D.W., Robinson, A., Simpson, J., Tuck, S.L., Weiher, E., White, H.J., Ewers, R.M., Mace, G.M., Scharlemann, J.P.W., Purvis, A., 2015. Global effects of land use on local terrestrial biodiversity. *Nature* 520, 45–50.
- Pereira, H.M., Navarro, L.M. (Eds.), 2015. *Rewilding European Landscapes*. Springer.
- Phalan, B., Onial, M., Balmford, A., Green, R.E., 2011. Reconciling food production and biodiversity conservation: land sharing and land sparing compared. *Science* 333, 1289–1291.
- Plieninger, T., Höchtl, F., Spek, T., 2006. Traditional land-use and nature conservation in European rural landscapes. *Environ Sci Policy* 9, 317–321.
- Plue, J., Meuris, S., Verheyen, K., Hermy, M., 2009. The importance of artefacts of ancient land use on plant communities in Meerdaal forest, Belgium. *Belgian J Bot* 142, 16.
- Reif, J., Storch, D., Voříšek, P., Šťastný, K., Bejček, V., 2008. Bird-habitat associations predict population trends in central European forest and farmland birds. *Biodivers Conserv* 17, 3307–3319.
- Roberge, J.-M., Angelstam, P., Villard, M.-A., 2008. Specialised woodpeckers and naturalness in hemiboreal forests – Deriving quantitative targets for conservation planning. *Biol Conserv* 141, 997–1012.
- Salmen, H., 1980. *Die Ornithologie Siebenbürgens. Beiträge zu einer Monographie der Vogelwelt dieses Landes*. Bohlau Verlag.
- Schaublin, S., Bollmann, K., 2011. Winter habitat selection and conservation of Hazel Grouse (*Bonasa bonasia*) in mountain forests. *J Ornithol* 152, 179–192.
- Schmeller, D.S., Gruber, B., Budrys, E., Framsted, E., Lengyel, S., Henle, K., 2008. National responsibilities in European species conservation: A methodological review. *Conserv Biol* 22, 593–601.

- Schmeller, D.S., Maier, Evans, D., Henle, K., 2012. National responsibilities for conserving habitats – a freely scalable method. *Nat Conserv* 3, 21–44.
- Schmitt, T., Rákossy, L., 2007. Changes of traditional agrarian landscapes and their conservation implications: a case study of butterflies in Romania. *Divers Distrib* 13, 855–862.
- Sim, I.M.W., Burfield, I.J., Grant, M.C., Pearce-Higgins, J.W., Brooke, M. de L., 2007. The role of habitat composition in determining breeding site occupancy in a declining Ring Ouzel *Turdus torquatus* population. *Ibis (Lond 1859)* 149, 374–385.
- Skorka, P., Lenda, M., Tryjanowski, P., 2010. Invasive alien goldenrods negatively affect grassland bird communities in Eastern Europe. *Biol Conserv* 143, 856–861.
- The European Comission, 2014. Investing in the future of Natura 2000 Ensuring the future financing of the Natura 2000 Network. *Nat Biodivers Newsl* 15.
- Tscharntke, T., Clough, Y., Wanger, T.C., Jackson, L., Motzke, I., Perfecto, I., Vandermeer, J., Whitbread, A., 2012. Global food security, biodiversity conservation and the future of agricultural intensification. *Biol Conserv* 151, 53–59.
- Verhulst, J., Baldi, A., Kleijn, D., 2004. Relationship between land-use intensity and species richness and abundance of birds in Hungary. *Agric Ecosyst Environ* 104, 465–473.

Tables and figures

Table 5.1 Conservation responsibility classes for 252 bird species in the Carpathians

Carpathian contribution to European range	IUCN concern status	European population trend	No. species	Conservation responsibility class
high	C	decline	1	HHCD
high	C	stable/fluct	1	HHCD
high	LC	decline	3	HHLC
high	LC	stable/fluct	8	HHLC
high	LC	increase	2	HHLC
high	LC	NA	4	HHLC
medium	C	decline	10	HMCD
medium	LC	decline	41	LLLC
medium	LC	stable/fluct	66	LLLC
medium	LC	increase	23	LLLC
medium	LC	NA	10	LLLC
low	C	decline	2	LLLC
low	C	stable/fluct	1	LLLC
low	C	NA	1	LLLC
low	LC	decline	28	LLLC
low	LC	stable/fluct	24	LLLC
low	LC	increase	12	LLLC
low	LC	NA	15	LLLC
Total number of species			252	
% in classes of high conservation responsibility			67.06%	

Table 5.2 Major habitat types for species of high conservation responsibility. Highlighted species use more than one habitat type.

Major habitats considered in analysis				Major habitat not considered in analysis	
	Forest	Grassland	Agriculture	Wetland/Water/ Other	Unknown
HHCD	Aquila heliaca			Aquila heliaca Anser erythropus	
HHLC	Bombycilla garrulus Bonasa bonasia Clanga pomarina Ficedula albicollis Leiopicus medius Turdus torquatus Picoides tridactylus Picus canus Strix uralensis Sylvia nisorina Nucifraga caryocatactes	Crex crex Lanius minor Clanga pomarina Ficedula albicollis Plectrophenax nivalis Turdus torquatus		Crex crex	Dendrocopos syriacus Miliaria calandra Tichodroma muraria
HMCD	Milvus milvus	Milvus milvus Circus macrourus Coracias garrulus Gallinago media Limosa limosa		Acrocephalus paludicola Aythya nyroca Falco cherrug Gallinago media Limosa limosa Numenius arquata	Falco vespertinus
Total	13	11	0	9	4

Table 5.3 IUCN threats related to agricultural and forest systems for species of high conservation responsibility

Agricultural intensification and herbicides	Wood extraction and forestry	No threat information
Acrocephalus paludicola	Aquila heliaca	Bombycilla garrulus
Anser erythropus	Falco cherrug	Bonasa bonasia
Aythya nyroca		Clanga pomarina
Circus macrourus		Dendrocopos syriacus
Coracias garrulus		Ficedula albicollis
Crex crex		Leiopicus medius
Falco cherrug		Miliaria calandra
Falco vespertinus		Nucifraga caryocatactes
Gallinago media		Picoides tridactylus
Limosa limosa		Picus canus
Milvus milvus		Strix uralensis
Numenius arquata		Sylvia nisoria
		Tichodroma muraria

Figure 5.1 Study region: Carpathian Ecoregion in Eastern Europe.





Figure 5.2 (a) Evolution of potential major habitat (grassland, agriculture and forest) for a species whose range covers the entire Carpathians. (b) Evolution of agriculture, grassland and forest for all species that use these habitats. Double counting is possible if a species uses more than one habitat. This does not take into account dynamics of other habitat types.



b)




Figure 5.3 Percentage of European range in the Carpathians for 170 species, by major habitat type




Figure 5.4 Potential habitat dynamics for species of high conservation responsibility. The color of the line represents the conservation responsibility class. Red: HHCD Blue: HHLC Green: HMCD






Figure 5.5 IUCN threats and number of times they occur listed for the 29 species of conservation responsibility



IUCN Threat Classification Scheme, Level 1



Appendix

Appendix 5.1 Species of high conservation responsibility for the Carpathians and their species attributes

Scientific name	Common name	Family	Order	Conservation responsibility class
<i>Acrocephalus paludicola</i>	Aquatic Warbler	Sylviidae	PASSERIFORMES	HMCD
<i>Anser erythropus</i>	Lesser White-fronted Goose	Anatidae	ANSERIFORMES	HHCD
<i>Aquila heliaca</i>	Eastern Imperial Eagle	Accipitridae	ACCIPITRIFORMES	HHCD
<i>Aythya nyroca</i>	Ferruginous Duck	Anatidae	ANSERIFORMES	HMCD
<i>Bombycilla garrulus</i>	Bohemian Waxwing	Bombycillidae	PASSERIFORMES	HHLC
<i>Bonasa bonasia</i>	Hazel Grouse	Phasianidae	GALLIFORMES	HHLC
<i>Circus macrourus</i>	Pallid Harrier	Accipitridae	ACCIPITRIFORMES	HMCD
<i>Clanga pomarina</i>	Lesser Spotted Eagle	Accipitridae	ACCIPITRIFORMES	HHLC
<i>Coracias garrulus</i>	European Roller	Coraciidae	CORACIIFORMES	HMCD
<i>Crex crex</i>	Corncrake	Rallidae	GRUIFORMES	HHLC
<i>Dendrocopos syriacus</i>	Syrian Woodpecker	Picidae	PICIFORMES	HHLC
<i>Falco cherrug</i>	Saker Falcon	Falconidae	FALCONIFORMES	HMCD
<i>Falco vespertinus</i>	Red-footed Falcon	Falconidae	FALCONIFORMES	HMCD
<i>Ficedula albicollis</i>	Collared Flycatcher	Muscicapidae	PASSERIFORMES	HHLC
<i>Gallinago media</i>	Great Snipe	Scolopacidae	CHARADRIIFORMES	HMCD
<i>Lanius minor</i>	Lesser Grey Shrike	Laniidae	PASSERIFORMES	HHLC
<i>Leiopicus medius</i>	Middle Spotted Woodpecker	Picidae	PICIFORMES	HHLC
<i>Limosa limosa</i>	Black-tailed Godwit	Scolopacidae	CHARADRIIFORMES	HMCD
<i>Miliaria calandra</i>	Corn Bunting	Emberizidae	PASSERIFORMES	HHLC
<i>Milvus milvus</i>	Red Kite	Accipitridae	ACCIPITRIFORMES	HMCD
<i>Nucifraga caryocatactes</i>	Spotted Nutcracker	Corvidae	PASSERIFORMES	HHLC
<i>Numenius arquata</i>	Eurasian Curlew	Scolopacidae	CHARADRIIFORMES	HMCD

<i>Picoides tridactylus</i>	Three-toed Woodpecker	Picidae	PICIFORMES	HHLC
<i>Picus canus</i>	Grey-faced Woodpecker	Picidae	PICIFORMES	HHLC
<i>Plectrophenax nivalis</i>	Snow Bunting	Emberizidae	PASSERIFORMES	HHLC
<i>Strix uralensis</i>	Ural Owl	Strigidae	STRIGIFORMES	HHLC
<i>Sylvia nisoria</i>	Barred Warbler	Sylviidae	PASSERIFORMES	HHLC
<i>Tichodroma muraria</i>	Wallcreeper	Sittidae	PASSERIFORMES	HHLC
<i>Turdus torquatus</i>	Ring Ouzel	Turdidae	PASSERIFORMES	HHLC

(Appendix 5.1 cont)

Scientific name	European population trend	Percent of range in Carpathians	IUCN Red List Status	Forest dependency	Average mass (g)	Min elevation	Max elevation
<i>Acrocephalus paludicola</i>	decline	5.37%	VU	Non-forest	15	0	200
<i>Anser erythropus</i>	decline	6.39%	VU	Non-forest	1761	0	700
<i>Aquila heliaca</i>	stable/fluct	8.09%	VU	Medium	3215	0	1400
<i>Aythya nyroca</i>	decline	5.31%	NT	Non-forest	530	NA	NA
<i>Bombycilla garrulus</i>	stable/fluct	6.54%	LC	High	57	NA	NA
<i>Bonasa bonasia</i>	increase	8.22%	LC	High	415	1600	1800
<i>Circus macrourus</i>	decline	6.16%	NT	Low	389	0	4000
<i>Clanga pomarina</i>	NA	6.78%	LC	High	1370	0	2200
<i>Coracias garrulus</i>	decline	5.70%	NT	Low	141	NA	2400
<i>Crex crex</i>	stable/fluct	6.85%	LC	Non-forest	156	NA	3000
<i>Dendrocopos syriacus</i>	decline	8.53%	LC	Medium	77	NA	2700
<i>Falco cherrug</i>	decline	3.74%	EN	Medium	966	0	4700
<i>Falco vespertinus</i>	decline	4.85%	NT	Low	153	0	1500
<i>Ficedula albicollis</i>	increase	11.73%	LC	Low	13	NA	NA
<i>Gallinago media</i>	decline	4.36%	NT	Non-forest	171	NA	2400
<i>Lanius minor</i>	decline	6.82%	LC	Non-forest	47	NA	3400

<i>Leiopicus medius</i>	NA	6.77%	LC	High	68	NA	2300
<i>Limosa limosa</i>	decline	3.42%	NT	Non-forest	291	NA	NA
<i>Miliaria calandra</i>	decline	20.56%	LC	Non-forest	49	NA	500
<i>Milvus milvus</i>	decline	6.13%	NT	Medium	1080	0	800
<i>Nucifraga caryocatactes</i>	stable/fluct	10.13%	LC	High	178	NA	NA
<i>Numenius arquata</i>	decline	3.20%	NT	Low	806	NA	NA
<i>Picoides tridactylus</i>	NA	8.01%	LC	High	66	360	2700
<i>Picus canus</i>	stable/fluct	7.27%	LC	Medium	152	NA	1700
<i>Plectrophenax nivalis</i>	stable/fluct	6.60%	LC	Non-forest	42	NA	NA
<i>Strix uralensis</i>	stable/fluct	12.84%	LC	Medium	785	450	1600
<i>Sylvia nisoria</i>	NA	6.30%	LC	Medium	24	NA	NA
<i>Tichodroma muraria</i>	stable/fluct	11.62%	LC	Non-forest	15	NA	NA
<i>Turdus torquatus</i>	stable/fluct	9.45%	LC	Medium	114	300	3000

Appendix 5.2 Possible Carpathian landscape configuration. Left: habitat distribution in 2010. Right: potential future habitat distribution if all agricultural land would be abandoned, and forest succession would occur in all areas that were forested at any point after 1860s.

